

Full wwPDB EM Validation Report (i)

Mar 10, 2024 – 11:57 PM EDT

EMDB ID : EMD-20846

Title: Human HCN1 channel in a hyperpolarized conformation

Authors: Lee, C.-.H.; MacKinnon, R.

Deposited on : 2019-10-19

Resolution : 3.04 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at
https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70 Validation Pipeline (wwPDB-VP) : 2.36

1 Experimental information (i)

Property	Value	Source	
EM reconstruction method	SINGLE PARTICLE	Depositor	
Imposed symmetry	POINT, C4	Depositor	
Number of particles used	104860	Depositor	
Resolution determination method	FSC 0.143 CUT-OFF	Depositor	
CTF correction method	Not provided		
Microscope	FEI TITAN KRIOS	Depositor	
Voltage (kV)	300	Depositor	
Electron dose $(e^-/\text{Å}^2)$	75.0	Depositor	
Minimum defocus (nm)	Not provided		
Maximum defocus (nm)	Not provided		
Magnification	Not provided		
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor	
Maximum map value	4.166	Depositor	
Minimum map value	-2.670	Depositor	
Average map value	0.001	Depositor	
Map value standard deviation	0.088	Depositor	
Recommended contour level	0.27	Depositor	
Map size (Å)	329.59998, 329.59998, 329.59998	wwPDB	
Map dimensions	320, 320, 320	wwPDB	
Map angles (°)	90.0, 90.0, 90.0	wwPDB	
Pixel spacing (Å)	1.03, 1.03, 1.03	Depositor	




2 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-20846. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

2.1 Orthogonal projections (i)

2.1.1 Primary map



2.1.2 Raw map




The images above show the map projected in three orthogonal directions.




2.2 Central slices (i)


2.2.1 Primary map








Y Index: 160




Z Index: 160


2.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160

The images above show central slices of the map in three orthogonal directions.




2.3 Largest variance slices (i)


2.3.1 Primary map








Y Index: 163




Z Index: 186


2.3.2 Raw map



X Index: 160



Y Index: 160




Z Index: 187

The images above show the largest variance slices of the map in three orthogonal directions.




2.4 Orthogonal standard-deviation projections (False-color) (i)

2.4.1 Primary map



2.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




2.5 Orthogonal surface views (i)

2.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.27. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

2.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

2.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



3 Map analysis (i)

This section contains the results of statistical analysis of the map.


3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



3.2 Volume estimate (i)




The volume at the recommended contour level is $256~\mathrm{nm}^3$; this corresponds to an approximate mass of $231~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



3.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.329 $\rm \mathring{A}^{-1}$



4 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

4.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.329 $\rm \mathring{A}^{-1}$



4.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.04	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.75	4.29	3.79

^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.75 differs from the reported value 3.04 by more than 10 %

