



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:14 pm BST

PDB ID : 331D
Title : CRYSTAL STRUCTURE OF D(GCGCGCG) WITH 5'-OVERHANG G'S
Authors : Pan, B.; Ban, C.; Wahl, M.; Sundaralingam, M.
Deposited on : 1997-05-13
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 355 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*CP*GP*CP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	7	Total 142	C 67	N 29	O 40	P 6	0	0	0
1	B	7	Total 142	C 67	N 29	O 40	P 6	0	0	0

- Molecule 2 is COBALT HEXAMMINE(III) (three-letter code: NCO) (formula: $\text{CoH}_{18}\text{N}_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Co	N		
2	A	1	Total 7	Co 1	N 6	0	0
2	A	1	Total 7	Co 1	N 6	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	29	Total 29	O 29	0	0
3	B	28	Total 28	O 28	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: DNA (5'-D(*GP*CP*GP*CP*GP*CP*G)-3')

Chain A:  100%


G1 G2 G3 G4 G5 G6 G7

- Molecule 1: DNA (5'-D(*GP*CP*GP*CP*GP*CP*G)-3')

Chain B:  14% 86%


G8 G9 G10 G11 G12 G13 G14

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	20.41Å 29.65Å 51.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.65	Depositor
% Data completeness (in resolution range)	87.1 (8.00-1.65)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.184 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	355	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NCO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.58	1/159 (0.6%)	2.03	6/244 (2.5%)
1	B	1.59	1/159 (0.6%)	2.18	10/244 (4.1%)
All	All	1.59	2/318 (0.6%)	2.11	16/488 (3.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	DG	N9-C4	5.56	1.42	1.38
1	B	8	DG	C5'-C4'	5.03	1.56	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	11	DC	N1-C2-O2	8.20	123.82	118.90
1	B	11	DC	O4'-C1'-N1	7.94	113.56	108.00
1	B	8	DG	O4'-C1'-N9	7.17	113.02	108.00
1	A	4	DC	N1-C2-O2	7.11	123.16	118.90
1	B	13	DC	O4'-C1'-N1	6.91	112.84	108.00
1	A	2	DC	N1-C2-O2	6.87	123.02	118.90
1	B	10	DG	O4'-C4'-C3'	6.45	109.87	106.00
1	A	2	DC	O4'-C1'-N1	6.26	112.38	108.00
1	B	9	DC	O4'-C1'-N1	6.25	112.37	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	11	DC	N3-C2-O2	-5.96	117.73	121.90
1	B	14	DG	O4'-C4'-C3'	5.65	109.39	106.00
1	B	10	DG	O4'-C1'-C2'	5.47	110.28	105.90
1	B	9	DC	N1-C2-O2	5.34	122.11	118.90
1	A	7	DG	N3-C4-C5	-5.23	125.99	128.60
1	A	6	DC	O4'-C4'-C3'	5.21	109.12	106.00
1	A	5	DG	O4'-C4'-C3'	5.02	109.01	106.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	DG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	142	0	79	0	0
1	B	142	0	79	0	0
2	A	14	0	0	0	0
3	A	29	0	0	0	0
3	B	28	0	0	0	0
All	All	355	0	158	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NCO	A	15	-	6,6,6	4.75	6 (100%)	-		
2	NCO	A	16	-	6,6,6	5.11	6 (100%)	-		

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	16	NCO	CO-N6	5.31	2.15	1.96
2	A	16	NCO	CO-N1	5.25	2.15	1.96
2	A	15	NCO	CO-N6	5.22	2.15	1.96
2	A	16	NCO	CO-N2	5.13	2.15	1.96
2	A	16	NCO	CO-N4	5.12	2.14	1.96
2	A	15	NCO	CO-N2	5.04	2.14	1.96

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	16	NCO	CO-N3	4.98	2.14	1.96
2	A	15	NCO	CO-N4	4.86	2.14	1.96
2	A	16	NCO	CO-N5	4.83	2.13	1.96
2	A	15	NCO	CO-N1	4.78	2.13	1.96
2	A	15	NCO	CO-N5	4.55	2.12	1.96
2	A	15	NCO	CO-N3	3.95	2.10	1.96

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.