



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 16, 2023 – 08:06 pm GMT

PDB ID : 4A0B
Title : Structure of hsDDB1-drDDB2 bound to a 16 bp CPD-duplex (pyrimidine at D-1 position) at 3.8 Å resolution (CPD 4)
Authors : Scrima, A.; Fischer, E.S.; Iwai, S.; Gut, H.; Thoma, N.H.
Deposited on : 2011-09-08
Resolution : 3.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1212 (4.00-3.60)
Clashscore	141614	1288 (4.00-3.60)
Ramachandran outliers	138981	1243 (4.00-3.60)
Sidechain outliers	138945	1237 (4.00-3.60)
RSRZ outliers	127900	1121 (4.00-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1159	
1	C	1159	
2	B	382	
2	D	382	
3	G	15	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	15	
4	H	16	
4	J	16	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 23938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA DAMAGE-BINDING PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1090	8543	5423	1443	1631	46	0	0	0
1	C	1095	8582	5447	1449	1639	47	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	expression tag	UNP Q16531
A	-17	HIS	-	expression tag	UNP Q16531
A	-16	HIS	-	expression tag	UNP Q16531
A	-15	HIS	-	expression tag	UNP Q16531
A	-14	HIS	-	expression tag	UNP Q16531
A	-13	HIS	-	expression tag	UNP Q16531
A	-12	HIS	-	expression tag	UNP Q16531
A	-11	VAL	-	expression tag	UNP Q16531
A	-10	ASP	-	expression tag	UNP Q16531
A	-9	GLU	-	expression tag	UNP Q16531
A	-8	ASN	-	expression tag	UNP Q16531
A	-7	LEU	-	expression tag	UNP Q16531
A	-6	TYR	-	expression tag	UNP Q16531
A	-5	PHE	-	expression tag	UNP Q16531
A	-4	GLN	-	expression tag	UNP Q16531
A	-3	GLY	-	expression tag	UNP Q16531
A	-2	GLY	-	expression tag	UNP Q16531
A	-1	GLY	-	expression tag	UNP Q16531
A	0	ARG	-	expression tag	UNP Q16531
A	224	SER	GLU	engineered mutation	UNP Q16531
C	-18	MET	-	expression tag	UNP Q16531
C	-17	HIS	-	expression tag	UNP Q16531
C	-16	HIS	-	expression tag	UNP Q16531
C	-15	HIS	-	expression tag	UNP Q16531
C	-14	HIS	-	expression tag	UNP Q16531

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-13	HIS	-	expression tag	UNP Q16531
C	-12	HIS	-	expression tag	UNP Q16531
C	-11	VAL	-	expression tag	UNP Q16531
C	-10	ASP	-	expression tag	UNP Q16531
C	-9	GLU	-	expression tag	UNP Q16531
C	-8	ASN	-	expression tag	UNP Q16531
C	-7	LEU	-	expression tag	UNP Q16531
C	-6	TYR	-	expression tag	UNP Q16531
C	-5	PHE	-	expression tag	UNP Q16531
C	-4	GLN	-	expression tag	UNP Q16531
C	-3	GLY	-	expression tag	UNP Q16531
C	-2	GLY	-	expression tag	UNP Q16531
C	-1	GLY	-	expression tag	UNP Q16531
C	0	ARG	-	expression tag	UNP Q16531
C	224	SER	GLU	engineered mutation	UNP Q16531

- Molecule 2 is a protein called DNA DAMAGE-BINDING PROTEIN 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	354	2839	1804	498	526	11	0	0	0
2	D	355	2843	1806	499	527	11	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	76	MET	-	expression tag	UNP Q2YDS1
B	77	HIS	-	expression tag	UNP Q2YDS1
B	78	HIS	-	expression tag	UNP Q2YDS1
B	79	HIS	-	expression tag	UNP Q2YDS1
B	80	HIS	-	expression tag	UNP Q2YDS1
B	81	HIS	-	expression tag	UNP Q2YDS1
B	82	HIS	-	expression tag	UNP Q2YDS1
B	83	ARG	-	expression tag	UNP Q2YDS1
B	84	ARG	-	expression tag	UNP Q2YDS1
B	85	LEU	-	expression tag	UNP Q2YDS1
B	86	VAL	-	expression tag	UNP Q2YDS1
B	87	PRO	-	expression tag	UNP Q2YDS1
B	88	ARG	-	expression tag	UNP Q2YDS1
B	89	GLY	-	expression tag	UNP Q2YDS1
B	90	SER	-	expression tag	UNP Q2YDS1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	91	GLY	-	expression tag	UNP Q2YDS1
B	92	GLY	-	expression tag	UNP Q2YDS1
B	93	ARG	-	expression tag	UNP Q2YDS1
B	180	GLN	LEU	variant	UNP Q2YDS1
B	214	ARG	TRP	variant	UNP Q2YDS1
D	76	MET	-	expression tag	UNP Q2YDS1
D	77	HIS	-	expression tag	UNP Q2YDS1
D	78	HIS	-	expression tag	UNP Q2YDS1
D	79	HIS	-	expression tag	UNP Q2YDS1
D	80	HIS	-	expression tag	UNP Q2YDS1
D	81	HIS	-	expression tag	UNP Q2YDS1
D	82	HIS	-	expression tag	UNP Q2YDS1
D	83	ARG	-	expression tag	UNP Q2YDS1
D	84	ARG	-	expression tag	UNP Q2YDS1
D	85	LEU	-	expression tag	UNP Q2YDS1
D	86	VAL	-	expression tag	UNP Q2YDS1
D	87	PRO	-	expression tag	UNP Q2YDS1
D	88	ARG	-	expression tag	UNP Q2YDS1
D	89	GLY	-	expression tag	UNP Q2YDS1
D	90	SER	-	expression tag	UNP Q2YDS1
D	91	GLY	-	expression tag	UNP Q2YDS1
D	92	GLY	-	expression tag	UNP Q2YDS1
D	93	ARG	-	expression tag	UNP Q2YDS1
D	180	GLN	LEU	variant	UNP Q2YDS1
D	214	ARG	TRP	variant	UNP Q2YDS1

- Molecule 3 is a DNA chain called 5'-D(*DGP*GP*TP*GP*AP*AP*AP*(TTD)P*AP*GP*CP*AP*GP*DGP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	G	14	312	149	61	88	14	0	0	0
3	I	13	290	139	56	82	13	0	0	0

- Molecule 4 is a DNA chain called 5'-D(*CP*CP*TP*GP*CP*TP*CP*CP*TP*TP*TP*CP*AP*CP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	H	14	274	133	44	84	13	0	0	0

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
4	J	13	255	124	41	78	12	0	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA DAMAGE-BINDING PROTEIN 1






• Molecule 1: DNA DAMAGE-BINDING PROTEIN 1



V1132
R1138
I1139
H1140


● Molecule 2: DNA DAMAGE-BINDING PROTEIN 2

Chain B: 63% 26% 7%



● Molecule 2: DNA DAMAGE-BINDING PROTEIN 2

Chain D: 65% 24% 7%



● Molecule 3: 5'-D(*DGP*GP*TP*GP*AP*AP*AP*(TTD)P*AP*GP*CP*AP*GP*DGP)-3

Chain G: 47% 40% 7% 7%


D6
G2
G3
A6
A7
T8
T9
A11
G12
G15
G16

● Molecule 3: 5'-D(*DGP*GP*TP*GP*AP*AP*AP*(TTD)P*AP*GP*CP*AP*GP*DGP)-3

Chain I: 60% 20% 7% 13%



- Molecule 4: 5'-D(*CP*CP*TP*GP*CP*TP*CP*CP*TP*TP*TP*CP*AP*CP*CP*C)-3'



- Molecule 4: 5'-D(*CP*CP*TP*GP*CP*TP*CP*CP*TP*TP*TP*CP*AP*CP*CP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.10Å 145.90Å 224.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.72 – 3.80 47.72 – 3.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.72-3.80) 99.8 (47.72-3.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 3.77Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.243 , 0.319 0.240 , 0.313	Depositor DCC
R_{free} test set	1862 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	102.6	Xtrriage
Anisotropy	0.090	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 62.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	23938	wwPDB-VP
Average B, all atoms (Å ²)	126.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TTD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/8697	0.56	0/11777
1	C	0.37	0/8737	0.56	0/11831
2	B	0.37	0/2913	0.57	0/3957
2	D	0.37	0/2917	0.57	0/3962
3	G	0.63	0/306	1.13	2/470 (0.4%)
3	I	0.61	0/281	1.09	0/431
4	H	0.66	0/304	1.35	3/465 (0.6%)
4	J	0.65	0/283	1.17	1/433 (0.2%)
All	All	0.39	0/24438	0.61	6/33326 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	15	DG	P-O3'-C3'	5.89	126.77	119.70
4	J	12	DC	P-O3'-C3'	5.46	126.25	119.70
4	H	11	DT	C4-C5-C7	5.34	122.20	119.00
4	H	11	DT	C6-C5-C7	-5.21	119.77	122.90
4	H	12	DC	P-O3'-C3'	5.16	125.89	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8543	0	8538	184	0
1	C	8582	0	8588	177	0
2	B	2839	0	2785	63	0
2	D	2843	0	2788	66	0
3	G	312	0	171	8	0
3	I	290	0	160	4	0
4	H	274	0	160	6	0
4	J	255	0	149	0	0
All	All	23938	0	23339	494	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 494 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:GLU:HG2	1:A:54:GLU:HG3	1.44	0.97
1:A:864:LYS:HG3	1:A:899:LEU:HB2	1.51	0.93
1:A:204:LYS:HG2	1:A:205:GLY:H	1.33	0.91
1:C:391:ARG:HH21	1:C:711:HIS:HD2	1.18	0.90
2:D:102:GLN:HG3	2:D:107:HIS:HB3	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1076/1159 (93%)	960 (89%)	100 (9%)	16 (2%)	10	46
1	C	1081/1159 (93%)	953 (88%)	114 (10%)	14 (1%)	12	48
2	B	352/382 (92%)	307 (87%)	36 (10%)	9 (3%)	5	36
2	D	353/382 (92%)	318 (90%)	30 (8%)	5 (1%)	11	46

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2862/3082 (93%)	2538 (89%)	280 (10%)	44 (2%)	10	46

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	118	ILE
2	B	183	THR
2	B	194	ALA
1	C	707	ILE
1	C	855	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	957/1015 (94%)	873 (91%)	84 (9%)	10	38
1	C	962/1015 (95%)	870 (90%)	92 (10%)	8	34
2	B	313/335 (93%)	278 (89%)	35 (11%)	6	28
2	D	313/335 (93%)	284 (91%)	29 (9%)	9	35
All	All	2545/2700 (94%)	2305 (91%)	240 (9%)	8	35

5 of 240 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	421	ARG
2	D	202	GLN
1	C	244	LYS
2	D	185	PHE
2	D	393	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	412	ASN
2	D	370	GLN
1	C	261	HIS
2	D	202	GLN
1	C	845	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TTD	I	9	3	42,45,46	1.57	7 (16%)	62,74,77	2.72	16 (25%)
3	TTD	G	9	3	42,45,46	1.65	7 (16%)	62,74,77	2.44	17 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TTD	I	9	3	-	14/22/109/110	0/5/6/6
3	TTD	G	9	3	-	19/22/109/110	0/5/6/6

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	9	TTD	C1'-N1	4.68	1.51	1.45
3	G	9	TTD	C1R-N1T	4.61	1.51	1.45
3	I	9	TTD	C1R-N1T	4.18	1.51	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	9	TTD	C1'-N1	4.17	1.51	1.45
3	G	9	TTD	C2-N1	3.58	1.43	1.36

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	9	TTD	O4R-C1R-N1T	13.54	124.69	108.65
3	G	9	TTD	O4R-C1R-N1T	8.05	118.19	108.65
3	I	9	TTD	C5T-C4T-N3T	6.69	121.88	116.06
3	G	9	TTD	N3-C2-N1	6.61	123.56	116.69
3	I	9	TTD	C5-C4-N3	6.46	121.69	116.06

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	9	TTD	O4'-C4R-C5'-O5'
3	G	9	TTD	C3R-O3R-PB-O5P
3	G	9	TTD	C3R-O3R-PB-O4P
3	G	9	TTD	O4'-C1'-N1-C2
3	G	9	TTD	C5R-O5R-PB-O5P

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	9	TTD	1	0
3	G	9	TTD	5	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1090/1159 (94%)	0.12	67 (6%) 21 17	68, 109, 228, 265	0
1	C	1095/1159 (94%)	0.02	33 (3%) 50 40	70, 106, 215, 247	0
2	B	354/382 (92%)	-0.29	0 100 100	77, 98, 132, 144	0
2	D	355/382 (92%)	-0.22	0 100 100	74, 103, 147, 175	0
3	G	13/15 (86%)	-0.56	0 100 100	128, 164, 217, 229	0
3	I	12/15 (80%)	-0.34	0 100 100	167, 206, 232, 286	0
4	H	14/16 (87%)	-0.20	0 100 100	131, 154, 223, 232	0
4	J	13/16 (81%)	0.33	0 100 100	196, 230, 337, 374	0
All	All	2946/3144 (93%)	-0.01	100 (3%) 45 37	68, 106, 219, 374	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	584	GLY	6.4
1	A	583	GLY	6.3
1	A	497	ASN	4.8
1	A	548	ASP	4.7
1	A	547	GLY	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TTD	I	9	40/41	0.83	0.28	164,193,226,236	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TTD	G	9	40/41	0.89	0.23	128,141,161,167	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.