



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 10:48 pm BST

PDB ID : 5A2W  
Title : Crystal structure of mtPAP in complex with ATPgammaS  
Authors : Lapkouski, M.; Hallberg, B.M.  
Deposited on : 2015-05-26  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	 3% 79% 5% 16%
1	B	555	 4% 81% 16%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOCHONDRIAL PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3668	2358	606	681	23	0	2	0
1	B	467	3629	2335	593	680	21	0	0	0

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	14	MET	-	expression tag	UNP F1NBW0
A	15	HIS	-	expression tag	UNP F1NBW0
A	16	HIS	-	expression tag	UNP F1NBW0
A	17	HIS	-	expression tag	UNP F1NBW0
A	18	HIS	-	expression tag	UNP F1NBW0
A	19	HIS	-	expression tag	UNP F1NBW0
A	20	HIS	-	expression tag	UNP F1NBW0
A	21	SER	-	expression tag	UNP F1NBW0
A	22	SER	-	expression tag	UNP F1NBW0
A	23	GLY	-	expression tag	UNP F1NBW0
A	24	VAL	-	expression tag	UNP F1NBW0
A	25	ASP	-	expression tag	UNP F1NBW0
A	26	LEU	-	expression tag	UNP F1NBW0
A	27	GLY	-	expression tag	UNP F1NBW0
A	28	THR	-	expression tag	UNP F1NBW0
A	29	GLU	-	expression tag	UNP F1NBW0
A	30	ASN	-	expression tag	UNP F1NBW0
A	31	LEU	-	expression tag	UNP F1NBW0
A	32	TYR	-	expression tag	UNP F1NBW0
A	33	PHE	-	expression tag	UNP F1NBW0
A	34	GLN	-	expression tag	UNP F1NBW0
A	35	SER	-	expression tag	UNP F1NBW0
A	36	MET	-	expression tag	UNP F1NBW0
B	14	MET	-	expression tag	UNP F1NBW0
B	15	HIS	-	expression tag	UNP F1NBW0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	HIS	-	expression tag	UNP F1NBW0
B	17	HIS	-	expression tag	UNP F1NBW0
B	18	HIS	-	expression tag	UNP F1NBW0
B	19	HIS	-	expression tag	UNP F1NBW0
B	20	HIS	-	expression tag	UNP F1NBW0
B	21	SER	-	expression tag	UNP F1NBW0
B	22	SER	-	expression tag	UNP F1NBW0
B	23	GLY	-	expression tag	UNP F1NBW0
B	24	VAL	-	expression tag	UNP F1NBW0
B	25	ASP	-	expression tag	UNP F1NBW0
B	26	LEU	-	expression tag	UNP F1NBW0
B	27	GLY	-	expression tag	UNP F1NBW0
B	28	THR	-	expression tag	UNP F1NBW0
B	29	GLU	-	expression tag	UNP F1NBW0
B	30	ASN	-	expression tag	UNP F1NBW0
B	31	LEU	-	expression tag	UNP F1NBW0
B	32	TYR	-	expression tag	UNP F1NBW0
B	33	PHE	-	expression tag	UNP F1NBW0
B	34	GLN	-	expression tag	UNP F1NBW0
B	35	SER	-	expression tag	UNP F1NBW0
B	36	MET	-	expression tag	UNP F1NBW0

- Molecule 2 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (three-letter code: AGS) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	N			0	0	
			11	6	5					
2	A	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	15	Total 15	O 15	0	0
5	B	10	Total 10	O 10	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.64Å 96.70Å 192.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.07 – 2.50 38.04 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.07-2.50) 99.9 (38.04-2.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.226 , 0.253 0.228 , 0.253	Depositor DCC
$R_{free}$ test set	2036 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.1	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7405	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PEG, AGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/3761	0.58	4/5106 (0.1%)
1	B	0.30	0/3717	0.48	0/5055
All	All	0.30	0/7478	0.53	4/10161 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	129	GLU	CB-CA-C	-17.34	75.73	110.40
1	A	129	GLU	N-CA-C	9.35	136.23	111.00
1	A	269	PRO	CB-CA-C	-5.35	98.62	112.00
1	A	129	GLU	C-N-CA	5.24	134.79	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	129	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3668	0	3567	14	0
1	B	3629	0	3481	5	0
2	A	42	0	16	2	0
2	B	31	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	5	0	0
4	B	4	0	5	0	0
5	A	15	0	0	1	0
5	B	10	0	0	0	0
All	All	7405	0	7086	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1529:AGS:O3'	5:A:2011:HOH:O	1.99	0.81
1:A:180:ILE:HG22	1:A:184[A]:MET:HE3	1.73	0.69
1:A:180:ILE:HG22	1:A:184[A]:MET:CE	2.26	0.65
2:A:1529:AGS:H5'2	2:A:1529:AGS:PB	2.36	0.65
1:B:476:GLN:O	1:B:477:ASP:HB2	2.13	0.49
1:A:286:LEU:HD22	1:A:293:TYR:CE2	2.50	0.47
1:A:231:PHE:O	1:A:352:ARG:NH2	2.48	0.47
1:A:184[A]:MET:HE2	1:A:435:LEU:HD22	1.97	0.45
1:A:214:ALA:HB1	1:B:214:ALA:HB1	2.00	0.43
1:A:399:LEU:HB2	1:A:416:VAL:CG2	2.49	0.42
1:A:171:ILE:HB	1:A:172:PRO:HD3	2.01	0.42
1:A:180:ILE:CG2	1:A:184[A]:MET:CE	2.95	0.42
1:B:171:ILE:HB	1:B:172:PRO:HD3	2.01	0.42
1:B:312:GLN:N	1:B:313:PRO:CD	2.82	0.42
1:A:136:LYS:NZ	1:B:218:ASP:OD2	2.38	0.41
1:A:180:ILE:CG2	1:A:184[A]:MET:HE2	2.51	0.41
1:A:184[A]:MET:HE1	1:A:435:LEU:HB2	2.03	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:HIS:CG	1:A:130:HIS:O	2.75	0.40
1:A:312:GLN:N	1:A:313:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	466/555 (84%)	453 (97%)	12 (3%)	1 (0%)	47 68
1	B	463/555 (83%)	452 (98%)	10 (2%)	1 (0%)	47 68
All	All	929/1110 (84%)	905 (97%)	22 (2%)	2 (0%)	47 68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	130	HIS
1	B	269	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	401/497 (81%)	389 (97%)	12 (3%)	41 68
1	B	392/497 (79%)	382 (97%)	10 (3%)	46 72

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	793/994 (80%)	771 (97%)	22 (3%)	43 70

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	ILE
1	A	126	LYS
1	A	267	ARG
1	A	294	SER
1	A	312	GLN
1	A	324	ASN
1	A	328	ILE
1	A	428	THR
1	A	437	ASP
1	A	440	GLN
1	A	448	ARG
1	A	510	GLN
1	B	131	HIS
1	B	294	SER
1	B	324	ASN
1	B	328	ILE
1	B	347	LEU
1	B	401	ASP
1	B	402	GLU
1	B	423	LYS
1	B	437	ASP
1	B	440	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	312	GLN
1	A	510	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	A	1531	-	3,3,6	0.48	0	2,2,5	0.17	0
2	AGS	B	1528	3	26,33,33	4.56	9 (34%)	26,52,52	2.06	3 (11%)
2	AGS	A	1529	3	26,33,33	1.72	3 (11%)	26,52,52	1.55	7 (26%)
4	PEG	B	1530	-	3,3,6	0.46	0	2,2,5	0.10	0
2	AGS	A	1528	-	10,12,33	1.08	1 (10%)	8,17,52	1.65	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	1531	-	-	1/1/1/4	-
2	AGS	B	1528	3	-	4/17/38/38	0/3/3/3
2	AGS	A	1529	3	-	6/17/38/38	0/3/3/3
4	PEG	B	1530	-	-	1/1/1/4	-
2	AGS	A	1528	-	-	-	0/2/2/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1528	AGS	O4'-C1'	15.61	1.62	1.41
2	B	1528	AGS	C2'-C1'	-13.09	1.33	1.53
2	B	1528	AGS	O4'-C4'	-6.87	1.29	1.45
2	A	1529	AGS	PG-S1G	4.90	2.01	1.90
2	B	1528	AGS	C6-N6	3.93	1.48	1.34
2	B	1528	AGS	O2'-C2'	3.58	1.51	1.43
2	B	1528	AGS	C2-N1	3.37	1.40	1.33
2	B	1528	AGS	C2-N3	3.36	1.37	1.32
2	A	1529	AGS	PG-O3G	-3.10	1.44	1.54
2	A	1529	AGS	C2'-C1'	-3.05	1.49	1.53
2	B	1528	AGS	PA-O1A	2.38	1.59	1.50
2	A	1528	AGS	C5-C4	2.16	1.46	1.40
2	B	1528	AGS	O3'-C3'	-2.00	1.38	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1528	AGS	C5-C6-N6	6.58	130.34	120.35
2	B	1528	AGS	N3-C2-N1	-5.60	119.92	128.68
2	B	1528	AGS	N6-C6-N1	-4.27	109.72	118.57
2	A	1529	AGS	C3'-C2'-C1'	3.25	105.88	100.98
2	A	1528	AGS	N3-C2-N1	-3.21	123.66	128.68
2	A	1529	AGS	O2'-C2'-C3'	-2.69	103.12	111.82
2	A	1529	AGS	N3-C2-N1	-2.64	124.55	128.68
2	A	1529	AGS	O3G-PG-O3B	2.49	112.96	104.64
2	A	1529	AGS	C4-C5-N7	-2.20	107.11	109.40
2	A	1529	AGS	O3'-C3'-C2'	-2.17	104.81	111.82
2	A	1529	AGS	PA-O3A-PB	-2.16	125.41	132.83
2	A	1528	AGS	C4-C5-N7	-2.08	107.23	109.40

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1528	AGS	PB-O3B-PG-O2G
2	A	1529	AGS	PB-O3B-PG-O2G
2	A	1529	AGS	PB-O3B-PG-O3G
2	A	1529	AGS	O4'-C4'-C5'-O5'
2	A	1529	AGS	PA-O3A-PB-O1B
4	A	1531	PEG	O1-C1-C2-O2
4	B	1530	PEG	O1-C1-C2-O2
2	B	1528	AGS	PA-O3A-PB-O2B

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	1529	AGS	PA-O3A-PB-O2B
2	A	1529	AGS	C3'-C4'-C5'-O5'
2	B	1528	AGS	PB-O3B-PG-O3G
2	B	1528	AGS	PA-O3A-PB-O1B

There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1529	AGS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	468/555 (84%)	0.19	14 (2%) 50 53	50, 75, 115, 159	0
1	B	467/555 (84%)	0.36	22 (4%) 31 33	53, 75, 113, 153	0
All	All	935/1110 (84%)	0.28	36 (3%) 39 42	50, 75, 115, 159	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	128	ALA	7.1
1	B	127	ALA	6.9
1	B	152	ALA	6.1
1	B	130	HIS	6.0
1	A	130	HIS	4.3
1	A	410	GLY	4.2
1	A	128	ALA	4.2
1	A	477	ASP	3.8
1	B	153	GLU	3.6
1	A	245	HIS	3.5
1	A	411	TYR	3.4
1	A	129	GLU	3.4
1	B	513	ILE	3.2
1	B	516	GLU	3.1
1	B	129	GLU	3.1
1	B	401	ASP	3.1
1	B	465	SER	3.0
1	A	507	ASP	2.8
1	B	400	ALA	2.8
1	B	419	LEU	2.8
1	B	378	ILE	2.6
1	B	154	GLU	2.5
1	A	419	LEU	2.5
1	A	364	VAL	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	348	VAL	2.4
1	B	515	LYS	2.4
1	B	375	THR	2.3
1	A	176	HIS	2.3
1	A	363	SER	2.3
1	A	368	TRP	2.2
1	A	127	ALA	2.2
1	B	344	VAL	2.1
1	B	364	VAL	2.1
1	B	418	ASP	2.1
1	B	175	CYS	2.1
1	B	514	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	AGS	A	1528	11/31	0.73	0.27	133,137,139,139	0
3	MG	A	1530	1/1	0.78	0.05	65,65,65,65	0
4	PEG	B	1530	4/7	0.85	0.50	67,68,70,71	0
3	MG	B	1529	1/1	0.92	0.22	64,64,64,64	0
4	PEG	A	1531	4/7	0.93	0.28	64,65,65,65	0
2	AGS	A	1529	31/31	0.96	0.10	58,70,75,75	0
2	AGS	B	1528	31/31	0.97	0.15	54,63,66,67	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


**Electron density around AGS A 1528:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around AGS A 1529:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.