



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 28, 2024 – 05:44 am BST

PDB ID : 4A3G
Title : RNA Polymerase II initial transcribing complex with a 2nt DNA-RNA hybrid
Authors : Cheung, A.C.M.; Sainsbury, S.; Cramer, P.
Deposited on : 2011-09-30
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)
RNA backbone	3102	1002 (4.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1732	 55% 23% 18%
2	B	1224	 60% 26% 9%
3	C	318	 54% 26% 16%
4	D	221	 56% 18% 6% 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	215	 69% 28% .
6	F	155	 35% 17% . 46%
7	G	171	 64% 29% 7% .
8	H	146	 2% 60% 27% 5% 9%
9	I	122	 3% 70% 23% . .
10	J	70	 47% 34% 11% 7%
11	K	120	 % 70% 23% . .
12	L	70	 6% 36% 26% . 34%
13	N	14	 21% 50% 14% 36%
14	P	2	 50% 50%
15	T	27	 7% 48% 7% 44%

2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 31728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1422	11174	7037	1954	2121	62	0	0	0

- Molecule 2 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	1115	8859	5609	1554	1641	55	0	0	0

- Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	266	2095	1317	348	417	13	0	0	0

- Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	178	1434	887	257	288	2	0	0	0

- Molecule 5 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	214	1752	1111	309	321	11	0	0	0

- Molecule 6 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	84	679	434	115	127	3	0	0	0

- Molecule 7 is a protein called RPB7, DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	171	1340	861	222	249	8	0	0	0

- Molecule 8 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	133	1068	673	180	211	4	0	0	0

- Molecule 9 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	119	971	596	179	186	10	0	0	0

- Molecule 10 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	J	65	532	339	93	94	6	0	0	0

- Molecule 11 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	K	115	920	590	157	171	2	0	0	1

- Molecule 12 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	L	46	363	224	72	63	4	0	0	0

- Molecule 13 is a DNA chain called NON TEMPLATE DNA 5'-D(*GP*GP*CP*AP*CP*AP*AP*CP*TP*GP*CP *GP*CP*TP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
13	N	9	181	87	36	50	8	0	0	0

- Molecule 14 is a RNA chain called TRANSCRIPT RNA 5'-R(*CP*AP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
14	P	2	42	19	8	13	2	0	0	0

- Molecule 15 is a DNA chain called TEMPLATE DNA 5'-D(*AP*GP*CP*GP*CP*AP*GP*TP*TP*GP*TP*GP *CP*TP*AP*TP*GP*AP*BRUP*AP*TP*TP*TP*TP*TP*AP*D T)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	Br	C	N	O	P			
15	T	15	309	1	147	53	93	15	0	0	0

- Molecule 16 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	2	Total	Zn	0	0
			2	2		
16	B	1	Total	Zn	0	0
			1	1		
16	C	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		
16	J	1	Total	Zn	0	0
			1	1		
16	L	1	Total	Zn	0	0
			1	1		


- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	1	Total	Mg	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1







- Molecule 6: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2




- Molecule 7: RPB7, DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7



- Molecule 8: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3




- Molecule 9: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9




- Molecule 10: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5

Chain J: 



- Molecule 11: DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11

Chain K: 



ASP
ALA
PHE


- Molecule 12: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4

Chain L: 



- Molecule 13: NON TEMPLATE DNA 5'-D(*GP*GP*CP*AP*CP*AP*AP*CP*TP*GP*CP*GP*CP*TP)-3'

Chain N: 



- Molecule 14: TRANSCRIPT RNA 5'-R(*CP*AP)-3'

Chain P: 

C9
A10

- Molecule 15: TEMPLATE DNA 5'-D(*AP*GP*CP*GP*CP*AP*GP*TP*TP*GP*TP*GP*CP*TP*AP*TP*GP*AP*BRUP*AP*TP*TP*TP*TP*AP*DT)-3'

Chain T: 



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	220.67Å 394.13Å 282.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.27 – 3.50 49.27 – 3.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.27-3.50) 100.0 (49.27-3.50)	Depositor EDS
R_{merge}	0.96	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 3.48Å)	Xtrriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.166 , 0.194 0.187 , 0.214	Depositor DCC
R_{free} test set	3045 reflections (1.97%)	wwPDB-VP
Wilson B-factor (Å ²)	103.7	Xtrriage
Anisotropy	0.457	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 96.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.036 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.043 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	31728	wwPDB-VP
Average B, all atoms (Å ²)	120.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BRU, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	0/11374	0.85	14/15383 (0.1%)
2	B	0.52	0/9029	0.81	2/12171 (0.0%)
3	C	0.50	0/2133	0.78	0/2891
4	D	0.52	0/1444	0.85	2/1935 (0.1%)
5	E	0.47	0/1788	0.74	0/2406
6	F	0.61	0/691	0.83	0/933
7	G	0.52	0/1368	0.82	0/1844
8	H	0.50	0/1086	0.79	0/1470
9	I	0.46	0/989	0.76	0/1331
10	J	0.58	0/541	0.92	1/727 (0.1%)
11	K	0.49	0/938	0.73	0/1267
12	L	0.55	0/365	1.01	0/485
13	N	1.24	0/203	1.05	0/311
14	P	1.48	1/46 (2.2%)	0.90	0/69
15	T	1.19	0/323	1.01	0/497
All	All	0.54	1/32318 (0.0%)	0.83	19/43720 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	P	9	C	C1'-N1	6.17	1.58	1.48

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	LYS	C-N-CA	7.55	140.58	121.70
4	D	25	ALA	C-N-CA	7.43	140.28	121.70
1	A	399	HIS	N-CA-CB	6.90	123.03	110.60
1	A	56	PRO	C-N-CA	6.78	138.65	121.70
1	A	310	GLY	C-N-CA	5.91	136.46	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	34	LYS	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11174	0	11233	228	0
2	B	8859	0	8901	182	0
3	C	2095	0	2051	57	0
4	D	1434	0	1460	30	0
5	E	1752	0	1776	36	0
6	F	679	0	701	23	0
7	G	1340	0	1357	42	0
8	H	1068	0	1040	22	0
9	I	971	0	927	17	0
10	J	532	0	542	17	0
11	K	920	0	929	19	0
12	L	363	0	386	8	0
13	N	181	0	102	1	0
14	P	42	0	22	1	0
15	T	309	0	169	1	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	C	1	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	31728	0	31596	601	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 601 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:867:ILE:CG1	1:A:867:ILE:CD1	1.75	1.54
1:A:868:TYR:CE1	1:A:1064:VAL:HG11	1.90	1.05
2:B:114:PRO:HG3	2:B:181:LEU:HD11	1.47	0.95
6:F:109:VAL:HG11	6:F:123:LYS:HG2	1.57	0.86
2:B:766:ARG:HE	2:B:1020:ARG:HG2	1.39	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1414/1732 (82%)	1224 (87%)	137 (10%)	53 (4%)	3	26
2	B	1095/1224 (90%)	956 (87%)	101 (9%)	38 (4%)	3	27
3	C	264/318 (83%)	232 (88%)	28 (11%)	4 (2%)	10	45
4	D	174/221 (79%)	155 (89%)	10 (6%)	9 (5%)	2	18
5	E	212/215 (99%)	192 (91%)	17 (8%)	3 (1%)	11	46
6	F	82/155 (53%)	75 (92%)	7 (8%)	0	100	100
7	G	169/171 (99%)	152 (90%)	12 (7%)	5 (3%)	4	30
8	H	129/146 (88%)	101 (78%)	20 (16%)	8 (6%)	1	15
9	I	117/122 (96%)	94 (80%)	18 (15%)	5 (4%)	2	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
10	J	63/70 (90%)	53 (84%)	4 (6%)	6 (10%)	0	8
11	K	113/120 (94%)	108 (96%)	5 (4%)	0	100	100
12	L	44/70 (63%)	28 (64%)	14 (32%)	2 (4%)	2	21
All	All	3876/4564 (85%)	3370 (87%)	373 (10%)	133 (3%)	3	28

5 of 133 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	GLU
1	A	48	ALA
1	A	55	ASP
1	A	57	ARG
1	A	58	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1240/1519 (82%)	1042 (84%)	198 (16%)	2	14
2	B	966/1061 (91%)	822 (85%)	144 (15%)	3	17
3	C	234/274 (85%)	206 (88%)	28 (12%)	5	24
4	D	160/200 (80%)	132 (82%)	28 (18%)	2	10
5	E	196/197 (100%)	179 (91%)	17 (9%)	10	38
6	F	74/137 (54%)	67 (90%)	7 (10%)	8	34
7	G	152/152 (100%)	128 (84%)	24 (16%)	2	15
8	H	117/128 (91%)	107 (92%)	10 (8%)	10	39
9	I	113/116 (97%)	103 (91%)	10 (9%)	10	38
10	J	60/65 (92%)	47 (78%)	13 (22%)	1	5
11	K	99/102 (97%)	87 (88%)	12 (12%)	5	24
12	L	40/57 (70%)	28 (70%)	12 (30%)	0	2
All	All	3451/4008 (86%)	2948 (85%)	503 (15%)	3	18

5 of 503 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	343	ILE
7	G	152	SER
2	B	835	GLN
7	G	138	THR
10	J	28	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
7	G	10	ASN
9	I	11	ASN
11	K	29	ASN
1	A	1393	ASN
1	A	1265	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
14	P	1/2 (50%)	1 (100%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
14	P	10	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
15	BRU	T	22	15	18,21,22	1.99	4 (22%)	26,30,33	2.93	9 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	BRU	T	22	15	-	0/7/21/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	T	22	BRU	C6-C5	5.03	1.44	1.34
15	T	22	BRU	BR-C5	4.51	1.99	1.88
15	T	22	BRU	C6-N1	3.68	1.44	1.38
15	T	22	BRU	C1'-N1	2.47	1.54	1.48

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	T	22	BRU	O4'-C1'-N1	8.29	122.67	107.86
15	T	22	BRU	N3-C2-N1	5.22	121.82	114.89
15	T	22	BRU	C6-C5-C4	-5.11	115.49	120.67
15	T	22	BRU	C5-C4-N3	4.84	118.91	113.34
15	T	22	BRU	C6-N1-C2	-4.30	116.94	121.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	934:LYS	C	935:ARG	N	5.56
1	B	351:TYR	C	352:ALA	N	2.97

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1422/1732 (82%)	-0.25	9 (0%) 89 86	57, 107, 167, 228	0
2	B	1115/1224 (91%)	-0.15	12 (1%) 80 75	62, 118, 181, 206	0
3	C	266/318 (83%)	-0.32	0 100 100	78, 111, 152, 186	0
4	D	178/221 (80%)	-0.26	1 (0%) 89 86	78, 118, 174, 194	0
5	E	214/215 (99%)	-0.25	0 100 100	81, 139, 190, 203	0
6	F	84/155 (54%)	-0.47	0 100 100	58, 83, 118, 131	0
7	G	171/171 (100%)	-0.22	0 100 100	81, 102, 144, 164	0
8	H	133/146 (91%)	0.21	3 (2%) 60 54	123, 155, 187, 211	0
9	I	119/122 (97%)	-0.10	4 (3%) 45 40	111, 149, 190, 205	0
10	J	65/70 (92%)	-0.38	0 100 100	88, 107, 144, 156	0
11	K	115/120 (95%)	-0.31	1 (0%) 84 79	72, 105, 141, 157	0
12	L	46/70 (65%)	-0.09	4 (8%) 10 11	91, 138, 166, 184	0
13	N	9/14 (64%)	1.11	3 (33%) 0 0	216, 225, 263, 270	0
14	P	2/2 (100%)	0.36	0 100 100	185, 185, 185, 193	0
15	T	14/27 (51%)	0.62	2 (14%) 2 3	141, 173, 267, 268	0
All	All	3953/4607 (85%)	-0.20	39 (0%) 82 77	57, 114, 180, 270	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	883	LEU	4.8
11	K	115	ALA	4.5
1	A	194	ALA	3.8
2	B	339	THR	3.7
1	A	191	THR	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	BRU	T	22	20/21	0.78	0.21	174,184,191,192	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
17	MG	A	2458	1/1	0.90	0.20	287,287,287,287	0
16	ZN	I	1122	1/1	0.97	0.04	184,184,184,184	0
16	ZN	L	1071	1/1	0.98	0.09	144,144,144,144	0
16	ZN	I	1121	1/1	0.99	0.12	123,123,123,123	0
16	ZN	A	2456	1/1	0.99	0.07	137,137,137,137	0
16	ZN	A	2457	1/1	0.99	0.15	72,72,72,72	0
16	ZN	B	2225	1/1	0.99	0.22	90,90,90,90	0
16	ZN	C	1269	1/1	1.00	0.12	84,84,84,84	0
16	ZN	J	1066	1/1	1.00	0.24	93,93,93,93	0

6.5 Other polymers [i](#)

There are no such residues in this entry.