



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 22, 2022 – 05:04 pm BST

PDB ID : 7A3J
Title : Crystal structure of DPP8 in complex with a 4-oxo-b-lactam based inhibitor, A272
Authors : Ross, B.H.; Huber, R.
Deposited on : 2020-08-18
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.30
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	898	 2% 89% 8%
1	B	898	 2% 87% 10%
1	C	898	 % 88% 9%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 20097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	820	Total 6653	C 4275	N 1114	O 1238	S 26	0	0	0
1	A	827	Total 6716	C 4317	N 1124	O 1247	S 28	0	0	0
1	B	812	Total 6580	C 4229	N 1098	O 1226	S 27	0	0	0

- Molecule 2 is 2-ethyl-2-methanoyl- {N}-[3-[[4-(naphthalen-1-ylmethyl)piperazin-1-yl]methyl]phenyl]butanamide (three-letter code: QXK) (formula: C₂₉H₃₅N₃O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	C	1	Total 34	C 29	N 3	O 2	0	0
2	A	1	Total 34	C 29	N 3	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	34	29	3	2	0	0

- Molecule 3 is trimethylamine oxide (three-letter code: TMO) (formula: C₃H₉NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	C	1	5	3	1	1	0	0
3	A	1	5	3	1	1	0	0
3	B	1	5	3	1	1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
4	C	1	1	1	0	0
4	A	1	1	1	0	0
4	B	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	7	Total 7	O 7	0	0
5	A	11	Total 11	O 11	0	0
5	B	10	Total 10	O 10	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	164.38Å 252.96Å 260.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.54 – 3.00 49.54 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.54-3.00) 100.0 (49.54-3.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.197 , 0.232 0.200 , 0.231	Depositor DCC
R_{free} test set	5422 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	94.7	Xtriage
Anisotropy	0.417	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20097	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TMO, QXK, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/6905	0.59	0/9370
1	B	0.36	0/6762	0.59	0/9174
1	C	0.37	0/6838	0.59	0/9280
All	All	0.37	0/20505	0.59	0/27824

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6716	0	6529	6	0
1	B	6580	0	6399	11	0
1	C	6653	0	6469	10	0
2	A	34	0	0	0	0
2	B	34	0	0	0	0
2	C	34	0	0	0	0
3	A	5	0	9	0	0
3	B	5	0	9	0	0
3	C	5	0	9	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	11	0	0	0	0
5	B	10	0	0	0	0
5	C	7	0	0	0	0
All	All	20097	0	19424	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:689:LEU:HD12	1:C:699:VAL:HG11	1.80	0.64
1:B:238:ILE:HD13	1:B:348:ILE:HD13	1.81	0.62
1:C:686:TYR:HB3	1:C:689:LEU:HD23	1.83	0.59
1:B:61:LEU:HD22	1:B:881:LEU:HD23	1.89	0.55
1:C:118:ILE:HD12	1:C:599:LEU:CD2	2.41	0.51
1:C:439:TYR:CE1	1:C:482:ILE:HD13	2.46	0.50
1:A:238:ILE:HG23	1:A:249:LEU:HD11	1.92	0.50
1:C:118:ILE:HD12	1:C:599:LEU:HD22	1.94	0.50
1:C:651:TYR:HB2	1:C:699:VAL:HB	1.96	0.48
1:B:651:TYR:HB2	1:B:699:VAL:HB	1.96	0.48
1:B:118:ILE:HD12	1:B:599:LEU:CD2	2.44	0.47
1:B:189:VAL:HG13	1:B:204:PRO:HA	1.96	0.47
1:B:376:GLU:HG3	1:B:397:ARG:HB2	1.96	0.47
1:C:61:LEU:HD22	1:C:881:LEU:HD23	1.97	0.46
1:B:118:ILE:HD12	1:B:599:LEU:HD22	1.98	0.46
1:B:627:LEU:HD13	1:B:628:PRO:HD2	1.98	0.46
1:B:65:LEU:HD21	1:B:881:LEU:HD22	1.98	0.45
1:A:118:ILE:HD12	1:A:599:LEU:CD2	2.47	0.45
1:A:762:LEU:HD13	1:A:811:VAL:HG21	2.00	0.44
1:C:334:THR:CG2	1:C:785:ILE:HA	2.47	0.44
1:B:302:LEU:HD22	1:B:390:ALA:HB1	1.99	0.43
1:A:752:HIS:CD2	1:A:880:LEU:HD21	2.53	0.43
1:B:273:LEU:HD13	1:B:279:ARG:NH2	2.34	0.43
1:C:689:LEU:HD12	1:C:699:VAL:CG1	2.46	0.43
1:C:65:LEU:HD21	1:C:881:LEU:HD22	2.00	0.43
1:A:596:CYS:SG	1:A:623:SER:HB2	2.60	0.41
1:A:118:ILE:HD12	1:A:599:LEU:HD22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	823/898 (92%)	786 (96%)	37 (4%)	0	100	100
1	B	804/898 (90%)	773 (96%)	30 (4%)	1 (0%)	51	85
1	C	814/898 (91%)	775 (95%)	36 (4%)	3 (0%)	34	72
All	All	2441/2694 (91%)	2334 (96%)	103 (4%)	4 (0%)	47	82

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	317	THR
1	C	626	PRO
1	B	445	ILE
1	C	445	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	733/795 (92%)	716 (98%)	17 (2%)	50	80
1	B	719/795 (90%)	701 (98%)	18 (2%)	47	79
1	C	727/795 (91%)	710 (98%)	17 (2%)	50	80
All	All	2179/2385 (91%)	2127 (98%)	52 (2%)	49	79

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	97	ASP
1	C	105	SER
1	C	129	MET
1	C	170	ASP
1	C	173	GLN
1	C	260	GLU
1	C	324	ARG
1	C	376	GLU
1	C	443	THR
1	C	473	LYS
1	C	545	ASP
1	C	550	HIS
1	C	592	LYS
1	C	614	LYS
1	C	629	ASP
1	C	718	PHE
1	C	881	LEU
1	A	138	LEU
1	A	164	VAL
1	A	170	ASP
1	A	173	GLN
1	A	214	ILE
1	A	346	SER
1	A	376	GLU
1	A	473	LYS
1	A	545	ASP
1	A	614	LYS
1	A	627	LEU
1	A	657	GLN
1	A	704	ASN
1	A	708	CYS
1	A	718	PHE
1	A	834	GLU
1	A	876	TYR
1	B	64	LEU
1	B	129	MET
1	B	173	GLN
1	B	189	VAL
1	B	324	ARG
1	B	340	LYS
1	B	376	GLU
1	B	431	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	473	LYS
1	B	545	ASP
1	B	614	LYS
1	B	622	ASP
1	B	627	LEU
1	B	688	ARG
1	B	690	ASN
1	B	705	ARG
1	B	718	PHE
1	B	829	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	111	ASN
1	C	173	GLN
1	C	550	HIS
1	C	837	HIS
1	A	111	ASN
1	A	199	GLN
1	A	200	GLN
1	A	882	HIS
1	B	108	ASN
1	B	111	ASN
1	B	657	GLN
1	B	862	GLN
1	B	882	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TMO	A	902	-	4,4,4	6.53	1 (25%)	6,6,6	0.25	0
2	QXK	B	901	1	35,37,37	0.76	0	47,51,51	1.47	8 (17%)
2	QXK	C	901	1	35,37,37	0.75	0	47,51,51	1.38	6 (12%)
3	TMO	C	902	-	4,4,4	6.29	1 (25%)	6,6,6	0.25	0
3	TMO	B	902	-	4,4,4	6.46	1 (25%)	6,6,6	0.23	0
2	QXK	A	901	1	35,37,37	0.75	0	47,51,51	1.34	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QXK	C	901	1	-	5/24/37/37	0/4/4/4
2	QXK	B	901	1	-	5/24/37/37	0/4/4/4
2	QXK	A	901	1	-	7/24/37/37	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	TMO	OAE-NAC	-12.90	1.25	1.42
3	B	902	TMO	OAE-NAC	-12.80	1.25	1.42
3	C	902	TMO	OAE-NAC	-12.51	1.25	1.42

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	901	QXK	C1-C8-N10	4.38	121.92	115.43
2	B	901	QXK	C1-C8-N10	4.16	121.61	115.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	QXK	O9-C8-C1	-3.89	114.95	121.21
2	A	901	QXK	O9-C8-C1	-3.88	114.96	121.21
2	C	901	QXK	O9-C8-C1	-3.80	115.09	121.21
2	A	901	QXK	C1-C8-N10	3.73	120.96	115.43
2	B	901	QXK	C22-C21-N18	-2.77	110.19	114.14
2	B	901	QXK	C21-C22-C27	2.53	124.35	119.93
2	B	901	QXK	C29-C26-C25	-2.51	117.33	123.19
2	C	901	QXK	C29-C26-C25	-2.43	117.50	123.19
2	C	901	QXK	C21-C22-C27	2.36	124.05	119.93
2	A	901	QXK	C29-C26-C25	-2.21	118.03	123.19
2	B	901	QXK	C16-C11-C12	-2.20	117.05	119.65
2	B	901	QXK	C19-N18-C17	2.17	113.72	108.83
2	A	901	QXK	C7-C13-C12	-2.15	116.09	120.25
2	C	901	QXK	C16-C11-C12	-2.14	117.11	119.65
2	B	901	QXK	C7-C13-C12	-2.13	116.14	120.25
2	C	901	QXK	C3-C2-C1	2.12	116.86	114.38
2	A	901	QXK	C19-N18-C17	2.12	113.59	108.83
2	A	901	QXK	C21-C22-C27	2.02	123.47	119.93

There are no chirality outliers.


All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	901	QXK	C2-C1-C8-O9
2	C	901	QXK	C2-C1-C8-N10
2	A	901	QXK	C2-C1-C8-O9
2	A	901	QXK	C2-C1-C8-N10
2	C	901	QXK	N18-C21-C22-C27
2	A	901	QXK	N18-C21-C22-C27
2	B	901	QXK	N18-C21-C22-C27
2	A	901	QXK	C4-C1-C2-C3
2	C	901	QXK	N18-C21-C22-C23
2	B	901	QXK	N18-C21-C22-C23
2	A	901	QXK	N18-C21-C22-C23
2	A	901	QXK	C6-C1-C2-C3
2	A	901	QXK	C4-C1-C8-O9
2	B	901	QXK	C2-C1-C8-O9
2	B	901	QXK	C2-C1-C8-N10
2	B	901	QXK	C4-C1-C8-N10
2	C	901	QXK	C8-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	827/898 (92%)	-0.02	22 (2%) 54 26	73, 95, 152, 225	0
1	B	812/898 (90%)	0.00	21 (2%) 56 27	71, 92, 145, 203	0
1	C	820/898 (91%)	-0.00	13 (1%) 72 44	73, 98, 155, 202	0
All	All	2459/2694 (91%)	-0.01	56 (2%) 60 31	71, 95, 152, 225	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	75	MET	6.3
1	C	319	PRO	5.7
1	B	323	THR	4.3
1	B	320	MET	4.3
1	C	326	ALA	4.2
1	B	318	SER	3.9
1	B	325	ARG	3.8
1	A	317	THR	3.8
1	C	318	SER	3.7
1	C	862	GLN	3.7
1	A	865	HIS	3.5
1	C	864	ARG	3.5
1	B	317	THR	3.5
1	A	260	GLU	3.4
1	A	73	GLY	3.3
1	B	326	ALA	3.3
1	C	320	MET	3.3
1	A	318	SER	3.1
1	B	106	GLY	3.1
1	A	850	ARG	3.0
1	A	322	GLU	3.0
1	B	108	ASN	3.0
1	B	327	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	256	LEU	2.9
1	B	526	GLY	2.9
1	A	106	GLY	2.8
1	B	365	ILE	2.8
1	C	871	GLU	2.8
1	B	319	PRO	2.7
1	A	108	ASN	2.7
1	A	261	GLU	2.6
1	A	323	THR	2.6
1	B	451	ASP	2.5
1	C	278	ASP	2.5
1	A	326	ALA	2.5
1	B	525	HIS	2.5
1	B	278	ASP	2.5
1	A	258	ASN	2.4
1	A	525	HIS	2.3
1	B	422	ARG	2.3
1	A	105	SER	2.2
1	A	278	ASP	2.2
1	B	259	MET	2.2
1	A	138	LEU	2.2
1	C	260	GLU	2.1
1	A	164	VAL	2.1
1	C	422	ARG	2.1
1	B	527	SER	2.1
1	C	425	LEU	2.1
1	B	490	LYS	2.1
1	C	865	HIS	2.1
1	A	862	GLN	2.0
1	B	524	ARG	2.0
1	C	108	ASN	2.0
1	A	140	GLN	2.0
1	A	526	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	QXK	B	901	34/34	0.80	0.38	102,124,136,137	0
4	CL	C	903	1/1	0.80	0.36	87,87,87,87	0
4	CL	B	903	1/1	0.81	0.43	97,97,97,97	0
2	QXK	A	901	34/34	0.84	0.36	106,126,138,139	0
2	QXK	C	901	34/34	0.85	0.33	112,125,139,142	0
3	TMO	C	902	5/5	0.91	0.44	90,92,92,95	0
3	TMO	A	902	5/5	0.91	0.42	89,94,97,102	0
4	CL	A	903	1/1	0.92	0.43	96,96,96,96	0
3	TMO	B	902	5/5	0.93	0.44	94,98,100,101	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around QXK A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around QXK C 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.