



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 14, 2023 – 08:01 PM JST

PDB ID : 6A5S  
Title : Structure of 14-3-3 gamma in complex with TFEB 14-3-3 binding motif  
Authors : Xu, Y.; Ren, J.Q.; Feng, W.  
Deposited on : 2018-06-25  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	
1	B	248	
1	D	248	
1	G	248	
2	C	15	
2	E	15	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	15	 <p>13% 53% 20% 27%</p>
2	H	15	 <p>7% 53% 20% 27%</p>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	Total 1887	C 1183	N 319	O 375	S 10	0	5	0
1	B	229	Total 1863	C 1165	N 317	O 372	S 9	0	1	0
1	D	233	Total 1893	C 1181	N 323	O 380	S 9	0	2	0
1	G	228	Total 1856	C 1164	N 316	O 367	S 9	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	SER	-	expression tag	UNP P61981
B	248	SER	-	expression tag	UNP P61981
D	248	SER	-	expression tag	UNP P61981
G	248	SER	-	expression tag	UNP P61981

- Molecule 2 is a protein called TFEB pS211-peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
2	E	11	Total 77	C 43	N 11	O 21	P 1	S 1	0	0	0
2	C	11	Total 77	C 43	N 11	O 21	P 1	S 1	0	0	0
2	F	11	Total 77	C 43	N 11	O 21	P 1	S 1	0	0	0
2	H	11	Total 77	C 43	N 11	O 21	P 1	S 1	0	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		



- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	207	Total	O	0	0
			207	207		
5	E	11	Total	O	0	0
			11	11		
5	B	198	Total	O	0	0
			198	198		
5	C	12	Total	O	0	0
			12	12		
5	D	173	Total	O	0	0
			173	173		
5	F	11	Total	O	0	0
			11	11		
5	G	181	Total	O	0	0
			181	181		
5	H	10	Total	O	0	0
			10	10		







- Molecule 2: TFEB pS211-peptide



- Molecule 2: TFEB pS211-peptide



- Molecule 2: TFEB pS211-peptide



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.09Å 82.23Å 88.09Å 90.00° 90.11° 90.00°	Depositor
Resolution (Å)	44.04 – 2.10 44.04 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.04-2.10) 92.2 (44.04-2.10)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.89 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.180 , 0.227 0.182 , 0.227	Depositor DCC
$R_{free}$ test set	2028 reflections (2.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtrriage
Anisotropy	0.790	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.031 for l,k,-h 0.000 for h,-k,-l 0.000 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8612	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0938e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1929	0.50	0/2601
1	B	0.38	0/1893	0.48	0/2553
1	D	0.37	0/1927	0.48	0/2600
1	G	0.38	0/1889	0.46	0/2548
2	C	0.45	0/66	0.50	0/89
2	E	0.38	0/66	0.48	0/89
2	F	0.41	0/66	0.51	0/89
2	H	0.44	0/66	0.51	0/89
All	All	0.39	0/7902	0.48	0/10658

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1887	0	1874	11	0
1	B	1863	0	1839	13	0
1	D	1893	0	1864	14	0
1	G	1856	0	1836	15	0
2	C	77	0	67	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	77	0	67	1	0
2	F	77	0	67	2	0
2	H	77	0	67	1	0
3	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	207	0	0	4	0
5	B	198	0	0	0	0
5	C	12	0	0	0	0
5	D	173	0	0	1	0
5	E	11	0	0	0	0
5	F	11	0	0	1	0
5	G	181	0	0	2	0
5	H	10	0	0	0	0
All	All	8612	0	7681	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ASN:HD21	2:C:11:LEU:H	1.08	0.92
1:B:139:THR:H	1:B:142:LYS:HE2	1.50	0.75
2:F:2:VAL:N	5:F:101:HOH:O	2.21	0.72
1:D:16[B]:GLN:NE2	5:D:304:HOH:O	2.30	0.64
1:A:142:LYS:NZ	5:A:402:HOH:O	2.26	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/248 (93%)	228 (99%)	3 (1%)	0	100	100
1	B	226/248 (91%)	224 (99%)	2 (1%)	0	100	100
1	D	233/248 (94%)	226 (97%)	7 (3%)	0	100	100
1	G	226/248 (91%)	223 (99%)	3 (1%)	0	100	100
2	C	8/15 (53%)	8 (100%)	0	0	100	100
2	E	8/15 (53%)	8 (100%)	0	0	100	100
2	F	8/15 (53%)	8 (100%)	0	0	100	100
2	H	8/15 (53%)	8 (100%)	0	0	100	100
All	All	948/1052 (90%)	933 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/217 (96%)	207 (100%)	1 (0%)	88	92
1	B	204/217 (94%)	202 (99%)	2 (1%)	76	82
1	D	207/217 (95%)	205 (99%)	2 (1%)	76	82
1	G	202/217 (93%)	200 (99%)	2 (1%)	76	82
2	C	9/12 (75%)	9 (100%)	0	100	100
2	E	9/12 (75%)	9 (100%)	0	100	100
2	F	9/12 (75%)	9 (100%)	0	100	100
2	H	9/12 (75%)	8 (89%)	1 (11%)	6	3
All	All	857/916 (94%)	849 (99%)	8 (1%)	78	84

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	2	VAL
1	G	212	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	179	TYR
1	D	77	LYS
1	G	179	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	43	ASN
1	G	6	GLN
1	G	98	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SEP	H	6	2	8,9,10	1.49	1 (12%)	8,12,14	0.98	0
2	SEP	C	6	2	8,9,10	1.49	1 (12%)	8,12,14	1.41	2 (25%)
2	SEP	F	6	2	8,9,10	1.49	1 (12%)	8,12,14	1.07	0
2	SEP	E	6	2	8,9,10	1.49	1 (12%)	8,12,14	1.37	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	H	6	2	-	0/5/8/10	-
2	SEP	C	6	2	-	0/5/8/10	-
2	SEP	F	6	2	-	0/5/8/10	-
2	SEP	E	6	2	-	0/5/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	6	SEP	P-O1P	3.26	1.61	1.50
2	H	6	SEP	P-O1P	3.26	1.61	1.50
2	C	6	SEP	P-O1P	3.08	1.60	1.50
2	F	6	SEP	P-O1P	2.98	1.60	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	SEP	OG-CB-CA	2.95	111.01	108.14
2	E	6	SEP	P-OG-CB	-2.50	111.42	118.30
2	E	6	SEP	OG-CB-CA	2.44	110.52	108.14
2	C	6	SEP	P-OG-CB	-2.15	112.37	118.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/248 (92%)	-0.23	2 (0%) 84 86	12, 20, 42, 60	0
1	B	229/248 (92%)	-0.23	3 (1%) 77 80	13, 21, 41, 60	0
1	D	233/248 (93%)	-0.05	11 (4%) 31 37	12, 22, 51, 82	0
1	G	228/248 (91%)	-0.24	3 (1%) 77 80	13, 21, 41, 53	0
2	C	10/15 (66%)	0.57	1 (10%) 7 9	20, 30, 45, 65	0
2	E	10/15 (66%)	0.38	1 (10%) 7 9	19, 27, 45, 54	0
2	F	10/15 (66%)	0.87	2 (20%) 1 1	22, 32, 47, 58	0
2	H	10/15 (66%)	0.47	1 (10%) 7 9	21, 30, 45, 59	0
All	All	960/1052 (91%)	-0.16	24 (2%) 57 62	12, 22, 45, 82	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	74	GLY	6.2
1	D	73	ASP	5.8
2	C	2	VAL	5.1
2	H	2	VAL	4.8
2	F	2	VAL	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SEP	E	6	10/11	0.98	0.09	14,17,21,22	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SEP	C	6	10/11	0.99	0.07	16,17,21,23	0
2	SEP	F	6	10/11	0.99	0.10	16,21,23,24	0
2	SEP	H	6	10/11	0.99	0.09	15,20,21,22	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	A	301	1/1	0.87	0.20	41,41,41,41	0
4	MG	B	301	1/1	0.92	0.11	37,37,37,37	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.