



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 11:15 pm BST

PDB ID : 6A5Z  
Title : Crystal structure of human FXR/RXR-LBD heterodimer bound to HNC180 and 9cRA and SRC1  
Authors : Wang, N.; Liu, J.  
Deposited on : 2018-06-25  
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	 86% 10%
1	H	229	 86% 11%
2	D	238	 79% 9% 11%
2	L	238	 78% 9% 13%
3	E	16	 69% 25% 6%
3	F	16	 63% 6% 31%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	16	 <p>6% 63% 6% 31%</p>
3	I	16	 <p>44% 13% 44%</p>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bile acid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	220	Total 1808	C 1158	N 300	O 341	S 9	0	0	0
1	H	224	Total 1838	C 1174	N 307	O 348	S 9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	432	GLU	CYS	engineered mutation	UNP Q96RI1
A	466	GLU	CYS	engineered mutation	UNP Q96RI1
H	432	GLU	CYS	engineered mutation	UNP Q96RI1
H	466	GLU	CYS	engineered mutation	UNP Q96RI1


- Molecule 2 is a protein called Retinoic acid receptor RXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	212	Total 1684	C 1081	N 289	O 304	S 10	0	1	0
2	L	208	Total 1645	C 1058	N 282	O 295	S 10	0	0	0

- Molecule 3 is a protein called Nuclear receptor coactivator 1.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	15	Total 127	C 79	N 27	O 21	0	0	0
3	F	11	Total 94	C 60	N 20	O 14	0	0	0
3	G	11	Total 101	C 64	N 23	O 14	0	0	0
3	I	9	Total 81	C 53	N 18	O 10	0	0	0

- Molecule 4 is 2-[(1R,5S)-9-[[3-[2,6-bis(chloranyl)phenyl]-5-cyclopropyl-1,2-oxazol-4-yl]methoxy]-3-azabicyclo[3.3.1]nonan-3-yl]-1,3-benzothiazole-6-carboxylic acid (three-letter code: 9R3) (formula: C<sub>29</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	N	O			S
4	A	1	Total	C	Cl	N	O	S	0	0
			39	29	2	3	4	1		
4	H	1	Total	C	Cl	N	O	S	0	0
			39	29	2	3	4	1		

- Molecule 5 is (9cis)-retinoic acid (three-letter code: 9CR) (formula: C<sub>20</sub>H<sub>28</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			22	20	2		
5	L	1	Total	C	O	0	0
			22	20	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	O	0	0
			2	2		
6	D	1	Total	O	0	0
			1	1		
6	H	8	Total	O	0	0
			8	8		
6	L	5	Total	O	0	0
			5	5		
6	I	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bile acid receptor




- Molecule 1: Bile acid receptor



- Molecule 2: Retinoic acid receptor RXR-alpha



- Molecule 2: Retinoic acid receptor RXR-alpha







- Molecule 3: Nuclear receptor coactivator 1




- Molecule 3: Nuclear receptor coactivator 1



- Molecule 3: Nuclear receptor coactivator 1



- Molecule 3: Nuclear receptor coactivator 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.53Å 96.13Å 114.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.55 – 2.95 73.55 – 2.95	Depositor EDS
% Data completeness (in resolution range)	97.3 (73.55-2.95) 97.3 (73.55-2.95)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 2.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.221 , 0.282 0.228 , 0.285	Depositor DCC
$R_{free}$ test set	1069 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.1	Xtrriage
Anisotropy	0.373	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	7517	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9CR, 9R3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/1843	0.70	0/2488
1	H	0.47	0/1874	0.68	0/2528
2	D	0.49	0/1721	0.73	2/2327 (0.1%)
2	L	0.48	0/1677	0.74	3/2266 (0.1%)
3	E	0.46	0/129	0.71	0/171
3	F	0.49	0/95	0.76	0/125
3	G	0.41	0/102	0.66	0/134
3	I	0.52	0/82	0.75	0/108
All	All	0.48	0/7523	0.71	5/10147 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	302	ARG	NE-CZ-NH1	5.62	123.11	120.30
2	L	316	ARG	NE-CZ-NH1	5.58	123.09	120.30
2	D	316	ARG	NE-CZ-NH1	5.24	122.92	120.30
2	L	347	ASP	CB-CG-OD2	5.13	122.91	118.30
2	D	347	ASP	CB-CG-OD2	-5.06	113.75	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1808	0	1796	9	0
1	H	1838	0	1819	8	0
2	D	1684	0	1716	5	0
2	L	1645	0	1686	3	0
3	E	127	0	131	1	0
3	F	94	0	100	0	0
3	G	101	0	110	0	0
3	I	81	0	91	0	0
4	A	39	0	0	2	0
4	H	39	0	0	2	0
5	D	22	0	23	3	0
5	L	22	0	23	2	0
6	A	2	0	0	0	0
6	D	1	0	0	0	0
6	H	8	0	0	0	0
6	I	1	0	0	0	0
6	L	5	0	0	0	0
All	All	7517	0	7495	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:328:MET:HE2	4:H:501:9R3:CL1	2.35	0.63
1:A:309:GLN:HA	1:A:316:GLN:NE2	2.18	0.59
1:A:309:GLN:HA	1:A:316:GLN:HE22	1.70	0.57
1:H:328:MET:CE	4:H:501:9R3:CL1	2.90	0.56
1:A:328:MET:HE2	4:A:501:9R3:CL2	2.45	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/229 (93%)	201 (94%)	13 (6%)	0	100	100
1	H	218/229 (95%)	202 (93%)	16 (7%)	0	100	100
2	D	209/238 (88%)	202 (97%)	6 (3%)	1 (0%)	29	64
2	L	204/238 (86%)	198 (97%)	5 (2%)	1 (0%)	29	64
3	E	13/16 (81%)	9 (69%)	4 (31%)	0	100	100
3	F	9/16 (56%)	8 (89%)	1 (11%)	0	100	100
3	G	9/16 (56%)	7 (78%)	2 (22%)	0	100	100
3	I	7/16 (44%)	6 (86%)	1 (14%)	0	100	100
All	All	883/998 (88%)	833 (94%)	48 (5%)	2 (0%)	47	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	445	THR
2	L	445	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/212 (95%)	188 (93%)	14 (7%)	15	44
1	H	205/212 (97%)	185 (90%)	20 (10%)	8	27
2	D	184/205 (90%)	168 (91%)	16 (9%)	10	33
2	L	179/205 (87%)	163 (91%)	16 (9%)	9	32
3	E	14/15 (93%)	11 (79%)	3 (21%)	1	4
3	F	10/15 (67%)	9 (90%)	1 (10%)	7	26
3	G	11/15 (73%)	10 (91%)	1 (9%)	9	31
3	I	9/15 (60%)	7 (78%)	2 (22%)	1	3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	814/894 (91%)	741 (91%)	73 (9%)	9 32

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	285	LEU
1	H	350	GLU
3	E	747	LYS
1	H	328	MET
1	H	379	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	306	ASN
1	H	263	GLN
2	L	306	ASN
2	D	270	GLN
2	L	385	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	9CR	D	501	-	19,22,22	4.83	8 (42%)	26,30,30	9.97	13 (50%)
4	9R3	H	501	-	36,45,45	1.29	3 (8%)	42,67,67	1.76	11 (26%)
4	9R3	A	501	-	36,45,45	1.35	5 (13%)	42,67,67	1.77	10 (23%)
5	9CR	L	501	-	19,22,22	4.32	7 (36%)	26,30,30	9.48	12 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	9CR	D	501	-	-	9/13/32/32	0/1/1/1
4	9R3	H	501	-	-	0/7/45/45	0/7/7/7
4	9R3	A	501	-	-	0/7/45/45	0/7/7/7
5	9CR	L	501	-	-	8/13/32/32	0/1/1/1

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	L	501	9CR	C5-C6	12.24	1.55	1.34
5	D	501	9CR	C5-C6	11.71	1.54	1.34
5	D	501	9CR	C8-C9	-10.30	1.23	1.45
5	D	501	9CR	C12-C13	-8.26	1.28	1.45
5	L	501	9CR	C8-C9	-8.02	1.28	1.45

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	501	9CR	C7-C8-C9	32.77	175.75	126.23
5	D	501	9CR	C11-C10-C9	31.19	171.82	127.31
5	D	501	9CR	C7-C8-C9	29.45	170.74	126.23
5	L	501	9CR	C11-C10-C9	25.15	163.21	127.31
5	D	501	9CR	C11-C12-C13	15.75	170.67	126.42

There are no chirality outliers.

5 of 17 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	D	501	9CR	C11-C10-C9-C8
5	D	501	9CR	C11-C10-C9-C19
5	D	501	9CR	C10-C11-C12-C13
5	L	501	9CR	C10-C11-C12-C13
5	D	501	9CR	C9-C10-C11-C12

There are no ring outliers.



4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	501	9CR	3	0
4	H	501	9R3	2	0
4	A	501	9R3	2	0
5	L	501	9CR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	220/229 (96%)	0.16	2 (0%) 84 71	24, 51, 102, 124	0
1	H	224/229 (97%)	0.53	9 (4%) 38 25	39, 71, 112, 133	0
2	D	212/238 (89%)	-0.07	2 (0%) 84 71	23, 37, 63, 97	0
2	L	208/238 (87%)	0.00	1 (0%) 91 81	29, 44, 66, 89	0
3	E	15/16 (93%)	0.16	0 100 100	46, 58, 96, 97	0
3	F	11/16 (68%)	0.70	1 (9%) 9 5	37, 47, 99, 109	0
3	G	11/16 (68%)	0.51	1 (9%) 9 5	79, 91, 113, 118	0
3	I	9/16 (56%)	-0.47	0 100 100	32, 40, 53, 57	0
All	All	910/998 (91%)	0.17	16 (1%) 68 51	23, 50, 102, 133	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	640	GLY	8.4
1	H	342	SER	6.5
1	H	333	ALA	4.7
1	A	346	ASP	4.2
2	D	228	GLU	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	9R3	H	501	39/39	0.91	0.28	46,54,76,83	0
5	9CR	D	501	22/22	0.94	0.23	30,34,36,37	0
4	9R3	A	501	39/39	0.94	0.21	33,37,51,52	0
5	9CR	L	501	22/22	0.94	0.22	28,31,38,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 9CR D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 9R3 A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.