

# Full wwPDB X-ray Structure Validation Report (i)

#### Aug 16, 2023 - 10:34 AM EDT

PDB ID	:	2A8H
Title	:	Crystal structure of catalytic domain of TACE with Thiomorpholine Sulfon-
		amide Hydroxamate inhibitor
Authors	:	Levin, J.I.; Chen, J.M.; Laakso, L.M.; Du, M.; Schmid, J.; Xu, W.; Cummons,
		T.; Xu, J.; Jin, G.; Barone, D.; Skotnicki, J.S.
Deposited on	:	2005-07-08
Resolution	:	2.30  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575(2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	271	3% 80%	14%	•••
1	В	271	8%	20%	• 5%



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	259	Total 2054	C 1290	N 347	O 404	S 13	0	0	0
1	В	257	Total 2038	C 1282	N 344	O 399	S 13	0	0	0

• Molecule 1 is a protein called ADAM 17.

Chain	Residue	Modelled	Actual	Comment	Reference
А	266	ALA	SER	engineered mutation	UNP P78536
А	452	GLN	ASN	engineered mutation	UNP P78536
А	478	GLY	-	expression tag	UNP P78536
А	479	SER	-	expression tag	UNP P78536
А	480	HIS	-	expression tag	UNP P78536
А	481	HIS	-	expression tag	UNP P78536
А	482	HIS	-	expression tag	UNP P78536
А	483	HIS	-	expression tag	UNP P78536
А	484	HIS	-	expression tag	UNP P78536
А	485	HIS	-	expression tag	UNP P78536
В	266	ALA	SER	engineered mutation	UNP P78536
В	452	GLN	ASN	engineered mutation	UNP P78536
В	478	GLY	-	expression tag	UNP P78536
В	479	SER	-	expression tag	UNP P78536
В	480	HIS	-	expression tag	UNP P78536
В	481	HIS	-	expression tag	UNP P78536
В	482	HIS	-	expression tag	UNP P78536
В	483	HIS	-	expression tag	UNP P78536
В	484	HIS	-	expression tag	UNP P78536
В	485	HIS	-	expression tag	UNP P78536

There are 20 discrepancies between the modelled and reference sequences:

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0

• Molecule 3 is 4-({4-[(4-AMINOBUT-2-YNYL)OXY]PHENYL}SULFONYL)-N-HYDRO XY-2,2-DIMETHYLTHIOMORPHOLINE-3-CARBOXAMIDE (three-letter code: 4NH) (formula:  $C_{17}H_{23}N_3O_5S_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
3	А	1	Total 27	C 17	N 3	O 5	${ m S} { m 2}$	0	0
3	В	1	Total 27	C 17	N 3	O 5	S 2	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	91	Total O 91 91	0	0
4	В	56	Total         O           56         56	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: ADAM 17



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	48.37Å 59.21Å 198.01Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution (Å)	20.00 - 2.30	Depositor
Resolution (A)	19.80 - 2.30	EDS
% Data completeness	99.1 (20.00-2.30)	Depositor
(in resolution range)	99.1 (19.80-2.30)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$11.88 (at 2.30 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
P. P.	0.215 , $0.281$	Depositor
$n, n_{free}$	0.216 , $0.279$	DCC
$R_{free}$ test set	1313 reflections $(5.07\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	29.2	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, $49.1$	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4295	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 4NH, ZN  $\,$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
MIOI	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.69	0/2101	0.67	0/2837	
1	В	0.54	0/2085	0.63	0/2814	
All	All	0.62	0/4186	0.65	0/5651	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	372	PRO	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2054	0	1960	30	0
1	В	2038	0	1951	52	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
3	А	27	0	22	1	0
3	В	27	0	22	3	0
4	А	91	0	0	2	0
4	В	56	0	0	3	0
All	All	4295	0	3955	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:370:TYR:HA	1:B:371:SER:CB	1.52	1.37
1:B:370:TYR:CA	1:B:371:SER:HB2	1.68	1.21
1:B:371:SER:N	1:B:372:PRO:HD2	1.40	1.17
1:B:371:SER:OG	1:B:376:LYS:O	1.70	1.08
1:B:373:VAL:HG22	1:B:374:GLY:H	1.10	1.06
1:A:369:TYR:CE2	1:A:378:ILE:HG12	1.95	1.02
1:B:371:SER:H	1:B:372:PRO:HD2	1.19	1.01
1:A:378:ILE:HG13	1:A:379:TYR:N	1.77	0.99
1:A:283:ARG:HB2	1:A:283:ARG:HH11	1.31	0.95
1:B:278:GLN:HE22	1:B:474:SER:N	1.64	0.94
1:B:373:VAL:CG2	1:B:374:GLY:N	2.30	0.94
1:B:371:SER:N	1:B:372:PRO:CD	2.30	0.92
1:B:278:GLN:HE22	1:B:474:SER:H	0.94	0.92
1:B:373:VAL:HG22	1:B:374:GLY:N	1.75	0.92
1:B:278:GLN:NE2	1:B:474:SER:H	1.74	0.83
1:B:370:TYR:CA	1:B:371:SER:CB	2.37	0.76
1:A:415:HIS:HE1	3:A:158:4NH:O4	1.66	0.76
1:B:307:GLU:HG2	4:B:66:HOH:O	1.85	0.75
1:B:370:TYR:HA	1:B:371:SER:HB2	0.76	0.74
1:B:269:ASN:HD21	1:B:452:GLN:HE22	1.37	0.72
1:A:297:HIS:HD2	1:A:299:ASN:H	1.39	0.71
1:A:269:ASN:HD21	1:A:452:GLN:HE22	1.43	0.65
1:B:312:TRP:H	1:B:341:GLN:NE2	1.97	0.63
1:A:279:ILE:HG21	1:A:282:ILE:HG13	1.79	0.63
1:A:306:ASN:O	1:A:309:LYS:HG2	1.98	0.62
1:A:290:GLU:HB3	1:B:429:GLN:HG2	1.80	0.62



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:283:ARG:HH11	1:A:283:ARG:CB	2.09	0.62
1:A:297:HIS:CD2	1:A:299:ASN:H	2.17	0.61
1:B:428:ASP:OD1	1:B:428:ASP:N	2.27	0.61
1:A:378:ILE:HD11	1:A:380:LEU:CD2	2.32	0.60
1:B:415:HIS:CE1	3:B:159:4NH:O4	2.54	0.60
1:A:378:ILE:HD11	1:A:380:LEU:HD23	1.84	0.59
1:B:306:ASN:HD21	1:B:308:GLU:HG3	1.67	0.59
1:B:361:HIS:HA	4:B:75:HOH:O	2.04	0.58
1:B:375:LYS:O	1:B:376:LYS:CB	2.51	0.58
1:B:361:HIS:HB3	1:B:380:LEU:HD12	1.85	0.58
1:B:304:TYR:CD1	1:B:305:PRO:HA	2.39	0.57
1:A:234:ARG:O	1:A:238:TYR:HD1	1.88	0.56
1:B:439:ALA:HB1	3:B:159:4NH:HN21	1.70	0.56
1:B:409:HIS:CD2	3:B:159:4NH:O5	2.59	0.56
1:B:297:HIS:CD2	1:B:299:ASN:H	2.24	0.56
1:A:369:TYR:HE2	1:A:378:ILE:HG12	1.63	0.55
1:A:260:ASP:OD1	1:A:263:ARG:NH2	2.38	0.54
1:B:226:LYS:HB3	1:B:280:GLU:HB2	1.89	0.54
1:B:375:LYS:O	1:B:376:LYS:HB3	2.09	0.53
1:A:369:TYR:CZ	1:A:378:ILE:HG21	2.44	0.53
1:B:357:ARG:CZ	1:B:360:SER:HB2	2.39	0.53
1:A:436:TYR:CD1	1:A:437:PRO:HD2	2.45	0.52
1:B:306:ASN:HD22	1:B:306:ASN:C	2.14	0.50
1:B:370:TYR:HA	1:B:371:SER:HB3	1.75	0.50
1:A:289:GLN:HG2	1:A:297:HIS:CG	2.47	0.50
1:A:292:LYS:HB3	1:A:293:PRO:HD2	1.94	0.49
1:B:219:ASP:C	1:B:220:PRO:O	2.51	0.49
1:B:269:ASN:HD21	1:B:452:GLN:NE2	2.09	0.49
1:B:230:VAL:HG22	1:B:283:ARG:HB2	1.94	0.48
1:B:370:TYR:C	1:B:372:PRO:HD2	2.25	0.48
1:B:224:THR:HG21	1:B:278:GLN:HE21	1.77	0.48
1:B:255:ILE:HG23	1:B:277:ILE:HD12	1.95	0.47
1:B:220:PRO:C	1:B:222:LYS:H	2.17	0.47
1:B:312:TRP:H	1:B:341:GLN:HE21	1.61	0.46
1:A:348:LEU:HD21	1:A:389:ASN:HB2	1.98	0.46
1:B:330:SER:HB3	1:B:379:TYR:CZ	2.51	0.46
1:B:356:PRO:HD3	4:B:146:HOH:O	2.16	0.45
1:A:217:ASP:HB3	4:A:9:HOH:O	2.15	0.45
1:A:397:LYS:O	1:A:401:LEU:HG	2.17	0.45
1:B:389:ASN:O	1:B:392:LYS:HG3	2.17	0.44
1:B:373:VAL:HG23	1:B:374:GLY:N	2.27	0.44

Continued from previous page...



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:416:ASP:HA	1:A:417:PRO:HD3	1.86	0.44
1:A:279:ILE:HG21	1:A:282:ILE:CG1	2.47	0.44
1:B:343:PHE:HB2	1:B:347:THR:HB	2.00	0.43
1:B:370:TYR:CG	1:B:371:SER:HB3	2.52	0.43
1:B:304:TYR:CG	1:B:305:PRO:HA	2.53	0.43
1:A:406:GLU:OE1	1:A:406:GLU:HA	2.18	0.43
1:B:371:SER:O	1:B:372:PRO:C	2.58	0.42
1:B:401:LEU:HD11	1:B:442:GLY:CA	2.49	0.42
1:A:306:ASN:HB2	1:A:309:LYS:HE2	2.02	0.42
1:B:352:TYR:HB3	1:B:362:GLY:HA2	2.02	0.42
1:A:369:TYR:O	1:A:378:ILE:HG23	2.20	0.41
1:A:308:GLU:HB2	4:A:21:HOH:O	2.21	0.41
1:B:458:ILE:HG22	1:B:462:ILE:HD12	2.01	0.41
1:A:378:ILE:HD11	1:A:380:LEU:HD21	2.02	0.40

Continued from previous page...

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	257/271 (95%)	247~(96%)	8(3%)	2(1%)	19 23
1	В	255/271 (94%)	240 (94%)	10 (4%)	5 (2%)	7 6
All	All	512/542 (94%)	487 (95%)	18 (4%)	7 (1%)	11 11

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	371	SER
1	В	372	PRO
1	В	376	LYS
1	А	312	TRP



 $Continued \ from \ previous \ page...$ 

Mol	Chain	Res	Type
1	А	365	CYS
1	В	365	CYS
1	В	220	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	$\mathbf{ntiles}$
1	А	222/232~(96%)	211~(95%)	11 (5%)	24	34
1	В	220/232~(95%)	207~(94%)	13~(6%)	19	27
All	All	442/464~(95%)	418 (95%)	24 (5%)	22	30

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	223	ASN
1	А	277	ILE
1	А	283	ARG
1	А	302	LYS
1	А	306	ASN
1	А	308	GLU
1	А	327	GLU
1	А	341	GLN
1	А	378	ILE
1	А	407	LEU
1	А	448	LYS
1	В	223	ASN
1	В	306	ASN
1	В	308	GLU
1	В	309	LYS
1	В	322	SER
1	В	392	LYS
1	В	407	LEU
1	В	414	GLU
1	В	420	LEU



Continued from previous page...

Mol	Chain	Res	Type
1	В	422	GLU
1	В	426	ASN
1	В	428	ASP
1	В	460	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	223	ASN
1	А	297	HIS
1	А	341	GLN
1	А	452	GLN
1	В	223	ASN
1	В	278	GLN
1	В	297	HIS
1	В	306	ASN
1	В	320	GLN
1	В	341	GLN
1	В	426	ASN
1	В	429	GLN
1	В	452	GLN
1	В	471	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Type Chain		in Res	Dec	Dag	Dec	Tiple	Bo	ond leng	ths	B	ond ang	gles
	Unain			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2				
3	4NH	В	159	2	26,28,28	1.99	3 (11%)	31,40,40	1.08	3 (9%)			
3	4NH	А	158	2	26,28,28	1.98	3 (11%)	31,40,40	1.08	3 (9%)			

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4NH	В	159	2	-	3/22/41/41	0/2/2/2
3	4NH	А	158	2	-	6/22/41/41	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	В	159	4NH	S2-N1	8.75	1.76	1.63
3	А	158	4NH	S2-N1	8.72	1.76	1.63
3	В	159	4NH	C16-N3	3.21	1.38	1.33
3	А	158	4NH	C16-N3	3.21	1.38	1.33
3	А	158	4NH	C4-N1	2.12	1.52	1.48
3	В	159	4NH	C4-N1	2.09	1.52	1.48

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	158	4NH	C6-S2-N1	-2.87	102.24	107.36
3	В	159	4NH	C6-S2-N1	-2.86	102.27	107.36
3	В	159	4NH	O1-S2-C6	-2.59	104.77	108.05
3	А	158	4NH	O1-S2-C6	-2.58	104.79	108.05
3	А	158	4NH	01-S2-N1	2.39	111.33	106.97
3	В	159	4NH	01-S2-N1	2.39	111.32	106.97

There are no chirality outliers.



Mol	Chain	Res	Type	Atoms
3	А	158	4NH	C12-C13-C14-C15
3	В	159	4NH	C12-C13-C14-C15
3	А	158	4NH	N3-C16-C3-N1
3	А	158	4NH	C10-C9-O3-C12
3	В	159	4NH	N3-C16-C3-N1
3	А	158	4NH	C8-C9-O3-C12
3	А	158	4NH	O4-C16-C3-N1
3	В	159	4NH	O4-C16-C3-N1
3	А	158	4NH	O3-C12-C13-C14

All (9) torsion outliers are listed below:

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	159	4NH	3	0
3	А	158	4NH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	259/271~(95%)	0.04	8 (3%) 49 56	12, 25, 43, 56	1 (0%)
1	В	257/271~(94%)	0.50	21 (8%) 11 15	16, 36, 59, 72	0
All	All	516/542~(95%)	0.27	29 (5%) 24 30	12, 30, 53, 72	1 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	373	VAL	10.3
1	В	372	PRO	5.2
1	В	369	TYR	4.2
1	В	370	TYR	4.2
1	В	359	ASN	4.1
1	А	293	PRO	4.1
1	В	361	HIS	4.0
1	В	375	LYS	3.7
1	В	294	GLY	3.2
1	А	291	VAL	3.2
1	В	360	SER	3.1
1	В	292	LYS	2.8
1	В	391	GLY	2.7
1	В	371	SER	2.7
1	В	295	GLU	2.7
1	В	270	ALA	2.7
1	А	428	ASP	2.7
1	А	308	GLU	2.6
1	В	293	PRO	2.5
1	В	390	TYR	2.4
1	В	221	MET	2.4
1	В	357	ARG	2.3
1	A	241	ARG	2.2
1	А	302	LYS	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	А	238	TYR	2.1
1	В	327	GLU	2.1
1	В	364	VAL	2.1
1	В	374	GLY	2.1
1	А	292	LYS	2.0

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
3	4NH	В	159	27/27	0.89	0.17	29,38,39,40	0
3	4NH	А	158	27/27	0.90	0.16	23,30,33,34	0
2	ZN	В	486	1/1	0.97	0.06	27,27,27,27	0
2	ZN	А	486	1/1	0.99	0.06	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers (i)

There are no such residues in this entry.

