



# wwPDB X-ray Structure Validation Summary Report

Oct 31, 2021 – 12:50 AM EDT

PDB ID : 2ABZ  
Title : Crystal structure of C19A/C43A mutant of leech carboxypeptidase inhibitor in complex with bovine carboxypeptidase A  
Authors : Arolas, J.L.; Popowicz, G.M.; Bronsoms, S.; Aviles, F.X.; Huber, R.; Holak, T.A.; Ventura, S.  
Deposited on : 2005-07-18  
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| $R_{free}$            | 130704                      | 1479 (2.16-2.16)                                      |
| Clashscore            | 141614                      | 1585 (2.16-2.16)                                      |
| Ramachandran outliers | 138981                      | 1560 (2.16-2.16)                                      |
| Sidechain outliers    | 138945                      | 1559 (2.16-2.16)                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 309    |                  |
| 1   | B     | 309    |                  |
| 2   | C     | 67     |                  |
| 2   | D     | 67     |                  |
| 2   | E     | 67     |                  |
| 2   | F     | 67     |                  |

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carboxypeptidase A1.

| Mol | Chain | Residues | Atoms |      |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
|     |       |          | Total | C    | N   | O   | S |         |         |       |
| 1   | A     | 299      | 2379  | 1531 | 390 | 453 | 5 | 0       | 0       | 0     |
| 1   | B     | 301      | 2392  | 1538 | 392 | 457 | 5 | 0       | 0       | 0     |

There are 4 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment        | Reference  |
|-------|---------|----------|--------|----------------|------------|
| A     | 228     | ALA      | GLU    | SEE REMARK 999 | UNP P00730 |
| A     | 305     | VAL      | LEU    | SEE REMARK 999 | UNP P00730 |
| B     | 228     | ALA      | GLU    | SEE REMARK 999 | UNP P00730 |
| B     | 305     | VAL      | LEU    | SEE REMARK 999 | UNP P00730 |

- Molecule 2 is a protein called Metallocoarboxypeptidase inhibitor.

| Mol | Chain | Residues | Atoms |     |    |    |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|---------|-------|
|     |       |          | Total | C   | N  | O  | S |         |         |       |
| 2   | C     | 62       | 476   | 301 | 79 | 90 | 6 | 0       | 0       | 0     |
| 2   | D     | 46       | 345   | 214 | 58 | 67 | 6 | 0       | 0       | 0     |
| 2   | E     | 60       | 463   | 293 | 77 | 87 | 6 | 0       | 0       | 0     |
| 2   | F     | 58       | 447   | 283 | 75 | 83 | 6 | 0       | 0       | 0     |

There are 12 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment             | Reference  |
|-------|---------|----------|--------|---------------------|------------|
| C     | 1       | GLY      | -      | cloning artifact    | UNP P81511 |
| C     | 19      | ALA      | CYS    | engineered mutation | UNP P81511 |
| C     | 43      | ALA      | CYS    | engineered mutation | UNP P81511 |
| D     | 1       | GLY      | -      | cloning artifact    | UNP P81511 |

*Continued on next page...*

*Continued from previous page...*

| Chain | Residue | Modelled | Actual | Comment             | Reference  |
|-------|---------|----------|--------|---------------------|------------|
| D     | 19      | ALA      | CYS    | engineered mutation | UNP P81511 |
| D     | 43      | ALA      | CYS    | engineered mutation | UNP P81511 |
| E     | 1       | GLY      | -      | cloning artifact    | UNP P81511 |
| E     | 19      | ALA      | CYS    | engineered mutation | UNP P81511 |
| E     | 43      | ALA      | CYS    | engineered mutation | UNP P81511 |
| F     | 1       | GLY      | -      | cloning artifact    | UNP P81511 |
| F     | 19      | ALA      | CYS    | engineered mutation | UNP P81511 |
| F     | 43      | ALA      | CYS    | engineered mutation | UNP P81511 |


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms           | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 3   | A     | 1        | Total Zn<br>1 1 | 0       | 0       |
| 3   | B     | 1        | Total Zn<br>1 1 | 0       | 0       |


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carboxypeptidase A1




- Molecule 1: Carboxypeptidase A1



- Molecule 2: Metallocoarboxypeptidase inhibitor



- Molecule 2: Metallocoarboxypeptidase inhibitor




- Molecule 2: Metallocoarboxypeptidase inhibitor





- Molecule 2: Metallocooxypeptidase inhibitor

Chain F:   
70% 12% 13%



## 4 Data and refinement statistics

| Property  | Value   | Source           |
|---|---|------------------|
| Space group   | P 43 21 2   | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 124.93Å 124.93Å 154.98Å<br>90.00° 90.00° 90.00°             | Depositor        |
| Resolution (Å)  | 30.00 – 2.16<br>97.26 – 2.16                                | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | (Not available) (30.00-2.16)<br>99.8 (97.26-2.16)           | Depositor<br>EDS |
| $R_{merge}$   | (Not available)   | Depositor        |
| $R_{sym}$   | (Not available)   | Depositor        |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>                              | 2.71 (at 2.16Å)   | Xtrriage         |
| Refinement program  | REFMAC 5.1.2  | Depositor        |
| R, $R_{free}$   | 0.189 , 0.234<br>0.241 , 0.257                              | Depositor<br>DCC |
| $R_{free}$ test set   | 3360 reflections (5.07%)                                    | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 28.9  | Xtrriage         |
| Anisotropy  | 0.335   | Xtrriage         |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.39 , 26.1   | EDS              |
| L-test for twinning <sup>2</sup>  | $\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$ | Xtrriage         |
| Estimated twinning fraction   | No twinning to report.                                      | Xtrriage         |
| $F_o, F_c$ correlation  | 0.92  | EDS              |
| Total number of atoms   | 6504  | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 32.0  | wwPDB-VP         |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |               | Bond angles |         |
|-----|-------|--------------|---------------|-------------|---------|
|     |       | RMSZ         | # Z  >5       | RMSZ        | # Z  >5 |
| 1   | A     | 0.47         | 0/2444        | 0.62        | 0/3322  |
| 1   | B     | 0.50         | 1/2457 (0.0%) | 0.63        | 0/3340  |
| 2   | C     | 0.46         | 0/494         | 0.70        | 0/680   |
| 2   | D     | 0.52         | 0/352         | 0.64        | 0/477   |
| 2   | E     | 0.56         | 0/479         | 0.68        | 0/658   |
| 2   | F     | 0.52         | 0/462         | 0.75        | 0/635   |
| All | All   | 0.49         | 1/6688 (0.0%) | 0.64        | 0/9112  |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z    | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 1   | B     | 136 | SER  | N-CA  | 6.76 | 1.59        | 1.46     |

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 2379  | 0        | 2290     | 60      | 1            |
| 1   | B     | 2392  | 0        | 2302     | 81      | 0            |
| 2   | C     | 476   | 0        | 431      | 34      | 0            |
| 2   | D     | 345   | 0        | 306      | 14      | 0            |

*Continued on next page...*



Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 2   | E     | 463   | 0        | 417      | 22      | 1            |
| 2   | F     | 447   | 0        | 407      | 12      | 0            |
| 3   | A     | 1     | 0        | 0        | 0       | 0            |
| 3   | B     | 1     | 0        | 0        | 0       | 0            |
| All | All   | 6504  | 0        | 6153     | 187     | 1            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 187 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1           | Atom-2          | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|-----------------|--------------------------|-------------------|
| 1:A:276:ARG:HH12 | 1:B:295:LEU:CD2 | 1.36                     | 1.38              |
| 1:B:133:THR:HG22 | 1:B:168:LYS:NZ  | 1.38                     | 1.34              |
| 1:B:276:ARG:NH2  | 1:B:277:TYR:HE2 | 1.35                     | 1.24              |
| 1:A:15:LEU:O     | 1:A:15:LEU:HD23 | 1.13                     | 1.22              |
| 1:B:15:LEU:C     | 1:B:15:LEU:HD23 | 1.60                     | 1.17              |

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1          | Atom-2               | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|----------------------|--------------------------|-------------------|
| 1:A:137:LEU:CD2 | 2:E:7:GLU:OE2[6_555] | 1.63                     | 0.57              |

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Favoured  | Allowed | Outliers | Percentiles |
|-----|-------|---------------|-----------|---------|----------|-------------|
| 1   | A     | 295/309 (96%) | 282 (96%) | 12 (4%) | 1 (0%)   | 41 37       |
| 1   | B     | 297/309 (96%) | 285 (96%) | 11 (4%) | 1 (0%)   | 41 37       |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed      | Favoured  | Allowed | Outliers | Percentiles |     |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 2   | C     | 60/67 (90%)   | 59 (98%)  | 1 (2%)  | 0        | 100         | 100 |
| 2   | D     | 36/67 (54%)   | 34 (94%)  | 2 (6%)  | 0        | 100         | 100 |
| 2   | E     | 56/67 (84%)   | 56 (100%) | 0       | 0        | 100         | 100 |
| 2   | F     | 52/67 (78%)   | 51 (98%)  | 0       | 1 (2%)   | 8           | 2   |
| All | All   | 796/886 (90%) | 767 (96%) | 26 (3%) | 3 (0%)   | 34          | 29  |

All (3) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | A     | 199 | SER  |
| 1   | B     | 199 | SER  |
| 2   | F     | 51  | VAL  |

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Rotameric | Outliers | Percentiles |    |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1   | A     | 256/265 (97%) | 248 (97%) | 8 (3%)   | 40          | 39 |
| 1   | B     | 258/265 (97%) | 249 (96%) | 9 (4%)   | 36          | 34 |
| 2   | C     | 52/56 (93%)   | 48 (92%)  | 4 (8%)   | 13          | 8  |
| 2   | D     | 39/56 (70%)   | 33 (85%)  | 6 (15%)  | 2           | 1  |
| 2   | E     | 50/56 (89%)   | 47 (94%)  | 3 (6%)   | 19          | 14 |
| 2   | F     | 50/56 (89%)   | 46 (92%)  | 4 (8%)   | 12          | 7  |
| All | All   | 705/754 (94%) | 671 (95%) | 34 (5%)  | 25          | 22 |

5 of 34 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2   | E     | 39  | THR  |
| 2   | E     | 49  | GLU  |
| 2   | F     | 49  | GLU  |
| 1   | B     | 199 | SER  |

Continued on next page...

*Continued from previous page...*

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | B     | 95  | SER  |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | B     | 285 | GLN  |
| 2   | E     | 35  | ASN  |
| 2   | D     | 57  | GLN  |
| 1   | B     | 37  | GLN  |
| 1   | B     | 249 | GLN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.