



# wwPDB NMR Structure Validation Summary Report ⓘ

Aug 7, 2020 – 11:51 AM BST

PDB ID : 1AC0  
Title : GLUCOAMYLASE, GRANULAR STARCH-BINDING DOMAIN COMPLEX WITH CYCLODEXTRIN, NMR, MINIMIZED AVERAGE STRUCTURE  
Authors : Sorimachi, K.; Le Gal-Coeffet, M.-F.; Williamson, G.; Archer, D.B.; Williamson, M.P.  
Deposited on : 1997-02-10

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.13.1  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1



## 2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.


### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1907 atoms, of which 915 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called GLUCOAMYLASE.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	108	1613	527	775	127	182	2	0

- Molecule 2 is an oligosaccharide called Cycloheptakis-(1-4)-(alpha-D-glucopyranose).




Mol	Chain	Residues	Atoms				Trace
			Total	C	H	O	
2	B	7	147	42	70	35	0
2	C	7	147	42	70	35	0


## 4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: GLUCOAMYLASE



- Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



- Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 1 were deposited, based on the following criterion: *RANDOM FROM 81 GOOD STRUCTURES*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	
X-PLOR	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	0	0	0	0
2	B	77	70	0	0
2	C	77	70	0	0
All	All	154	140	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	0	-	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	0	-	-	-	-

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	GLC	B	1	2	11,11,12	0.54	0 (0%)
2	GLC	B	2	2	11,11,12	0.39	0 (0%)



Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	GLC	B	3	2	11,11,12	0.43	0 (0%)
2	GLC	B	4	2	11,11,12	0.63	0 (0%)
2	GLC	B	5	2	11,11,12	0.52	0 (0%)
2	GLC	B	6	2	11,11,12	0.54	0 (0%)
2	GLC	B	7	2	11,11,12	0.43	0 (0%)
2	GLC	C	1	2	11,11,12	0.42	0 (0%)
2	GLC	C	2	2	11,11,12	0.67	0 (0%)
2	GLC	C	3	2	11,11,12	0.41	0 (0%)
2	GLC	C	4	2	11,11,12	0.65	0 (0%)
2	GLC	C	5	2	11,11,12	0.46	0 (0%)
2	GLC	C	6	2	11,11,12	0.53	0 (0%)
2	GLC	C	7	2	11,11,12	0.41	0 (0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	GLC	B	1	2	15,15,17	1.29	0 (0%)
2	GLC	B	2	2	15,15,17	0.82	0 (0%)
2	GLC	B	3	2	15,15,17	0.88	0 (0%)
2	GLC	B	4	2	15,15,17	0.79	0 (0%)
2	GLC	B	5	2	15,15,17	1.03	0 (0%)
2	GLC	B	6	2	15,15,17	1.22	0 (0%)
2	GLC	B	7	2	15,15,17	0.65	0 (0%)
2	GLC	C	1	2	15,15,17	1.31	0 (0%)
2	GLC	C	2	2	15,15,17	1.48	0 (0%)
2	GLC	C	3	2	15,15,17	0.45	0 (0%)
2	GLC	C	4	2	15,15,17	1.19	0 (0%)
2	GLC	C	5	2	15,15,17	1.02	0 (0%)
2	GLC	C	6	2	15,15,17	0.85	0 (0%)
2	GLC	C	7	2	15,15,17	0.96	0 (0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0,2,19,22	0,1,1,1
2	GLC	B	2	2	1,1,5,5	0,2,19,22	0,1,1,1
2	GLC	B	3	2	-	0,2,19,22	0,1,1,1
2	GLC	B	4	2	-	0,2,19,22	0,1,1,1
2	GLC	B	5	2	1,1,5,5	0,2,19,22	0,1,1,1
2	GLC	B	6	2	-	0,2,19,22	0,1,1,1
2	GLC	B	7	2	-	0,2,19,22	0,1,1,1
2	GLC	C	1	2	-	0,2,19,22	0,1,1,1
2	GLC	C	2	2	-	0,2,19,22	0,1,1,1
2	GLC	C	3	2	-	0,2,19,22	0,1,1,1
2	GLC	C	4	2	-	0,2,19,22	0,1,1,1
2	GLC	C	5	2	-	0,2,19,22	0,1,1,1
2	GLC	C	6	2	1,1,5,5	0,2,19,22	0,1,1,1
2	GLC	C	7	2	1,1,5,5	0,2,19,22	0,1,1,1

There are no bond-length outliers.

There are no bond-angle outliers.


All chiral outliers are listed below.


Mol	Chain	Res	Type	Atoms
2	C	7	GLC	C1
2	B	5	GLC	C1
2	B	2	GLC	C1
2	C	6	GLC	C1

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided