



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 12:33 pm BST

PDB ID : 4ACO
Title : Structure of the budding yeast Ndc10 N-terminal domain
Authors : Perriches, T.; Singleton, M.R.
Deposited on : 2011-12-16
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	956	 % 39% 7% 53%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CENTROMERE DNA-BINDING PROTEIN COMPLEX CBF3 SUBUNIT A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	452	3793	2473	619	691	10	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	86	Total	O	0	0
			86	86		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.74Å 87.76Å 104.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.22 – 1.89 47.05 – 1.89	Depositor EDS
% Data completeness (in resolution range)	98.4 (67.22-1.89) 98.5 (47.05-1.89)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.193 , 0.232 0.192 , 0.232	Depositor DCC
R_{free} test set	2054 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtrriage
Anisotropy	0.194	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3879	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.05	3/3908 (0.1%)	1.04	13/5297 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	369	TRP	CD2-CE2	6.78	1.49	1.41
1	A	193	TRP	CD2-CE2	5.53	1.48	1.41
1	A	401	PHE	CB-CG	-5.23	1.42	1.51

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	407	MET	CG-SD-CE	-15.01	76.19	100.20
1	A	231	ARG	NE-CZ-NH1	10.19	125.39	120.30
1	A	221	ARG	NE-CZ-NH1	8.55	124.58	120.30
1	A	347	ARG	NE-CZ-NH1	8.55	124.58	120.30
1	A	231	ARG	NE-CZ-NH2	-7.65	116.47	120.30
1	A	235	ARG	NE-CZ-NH1	6.59	123.60	120.30
1	A	122	LEU	CB-CG-CD1	-6.32	100.26	111.00
1	A	235	ARG	NE-CZ-NH2	-6.29	117.16	120.30
1	A	241	LEU	CB-CG-CD1	-6.08	100.66	111.00
1	A	123	LYS	CD-CE-NZ	-5.16	99.83	111.70
1	A	294	PHE	CB-CG-CD2	-5.06	117.26	120.80
1	A	239	LEU	CB-CG-CD2	5.05	119.58	111.00
1	A	325	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3793	0	3693	44	0
2	A	86	0	0	0	0
All	All	3879	0	3693	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:537:ARG:NH1	1:A:537:ARG:HB3	1.79	0.98
1:A:482:HIS:HD2	1:A:484:ILE:H	1.33	0.77
1:A:509:SER:O	1:A:527:LYS:HE2	1.85	0.75
1:A:537:ARG:HB3	1:A:537:ARG:HH11	1.53	0.71
1:A:379:ASN:ND2	1:A:381:GLU:HG3	2.09	0.68
1:A:379:ASN:HD21	1:A:381:GLU:HG3	1.57	0.68
1:A:537:ARG:CZ	1:A:537:ARG:HB3	2.25	0.66
1:A:202:GLU:O	1:A:206:LYS:HG3	1.97	0.65
1:A:482:HIS:CD2	1:A:484:ILE:HG12	2.32	0.64
1:A:493:SER:O	1:A:497:SER:HB2	1.99	0.63
1:A:166:LYS:HA	1:A:166:LYS:HE3	1.81	0.62
1:A:168:ILE:HG21	1:A:519:GLY:O	1.99	0.62
1:A:414:THR:HG22	1:A:414:THR:O	2.01	0.61
1:A:264:PRO:O	1:A:265:GLY:O	2.19	0.61
1:A:298:PRO:HG3	1:A:487:ASP:HA	1.82	0.59
1:A:487:ASP:OD1	1:A:489:ILE:HB	2.03	0.58
1:A:202:GLU:O	1:A:206:LYS:CG	2.52	0.57
1:A:473:PRO:HG2	1:A:502:LYS:HD3	1.87	0.55
1:A:482:HIS:CD2	1:A:484:ILE:H	2.22	0.49
1:A:537:ARG:CB	1:A:537:ARG:HH11	2.25	0.48
1:A:203:LYS:HE3	1:A:204:PHE:CZ	2.49	0.48
1:A:509:SER:OG	1:A:511:ILE:HG22	2.14	0.48
1:A:270:LEU:N	1:A:270:LEU:HD12	2.29	0.48
1:A:431:LEU:HD21	1:A:494:ASP:HB2	1.96	0.47
1:A:168:ILE:HD11	1:A:520:PHE:CE2	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:TYR:CZ	1:A:253:LYS:HG2	2.51	0.45
1:A:489:ILE:HG23	1:A:495:PHE:CE2	2.51	0.45
1:A:511:ILE:HG21	1:A:511:ILE:HD13	1.70	0.45
1:A:107:GLU:HG2	1:A:111:GLU:HB2	1.97	0.45
1:A:509:SER:O	1:A:527:LYS:CE	2.61	0.45
1:A:302:LYS:HB2	1:A:302:LYS:HE3	1.51	0.44
1:A:162:ILE:HD11	1:A:413:TYR:CE2	2.53	0.44
1:A:183:PRO:HA	1:A:184:PRO:HD3	1.83	0.43
1:A:447:TYR:CE1	1:A:455:ARG:HG2	2.54	0.43
1:A:131:PHE:CE2	1:A:135:LEU:HD22	2.53	0.43
1:A:224:LEU:HD22	1:A:270:LEU:HB3	2.01	0.43
1:A:427:PRO:HB3	1:A:469:LEU:HG	2.01	0.42
1:A:391:GLU:OE2	1:A:397:ASP:N	2.48	0.42
1:A:427:PRO:HA	1:A:428:PRO:HD3	1.92	0.42
1:A:441:GLU:HG3	1:A:444:ARG:HH21	1.85	0.42
1:A:129:LEU:HA	1:A:129:LEU:HD23	1.84	0.41
1:A:441:GLU:HG3	1:A:444:ARG:NH2	2.36	0.40
1:A:387:LEU:HA	1:A:387:LEU:HD12	1.88	0.40
1:A:414:THR:O	1:A:414:THR:CG2	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	440/956 (46%)	421 (96%)	16 (4%)	3 (1%)	22 12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	GLY
1	A	201	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	489	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	427/897 (48%)	404 (95%)	23 (5%)	22 13

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	47	THR
1	A	49	TYR
1	A	72	SER
1	A	108	ASP
1	A	166	LYS
1	A	239	LEU
1	A	253	LYS
1	A	299	SER
1	A	302	LYS
1	A	330	THR
1	A	338	PHE
1	A	339	SER
1	A	347	ARG
1	A	375	ARG
1	A	391	GLU
1	A	470	SER
1	A	489	ILE
1	A	493	SER
1	A	497	SER
1	A	507	LYS
1	A	522	LYS
1	A	537	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	379	ASN
1	A	482	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/956 (47%)	0.12	13 (2%) 51 54	12, 25, 53, 91	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	264	PRO	4.9
1	A	257	SER	4.7
1	A	107	GLU	4.0
1	A	414	THR	3.6
1	A	425	PHE	3.5
1	A	298	PRO	3.0
1	A	446	ASP	2.7
1	A	445	HIS	2.7
1	A	299	SER	2.6
1	A	97	ASP	2.6
1	A	96	ASP	2.4
1	A	522	LYS	2.3
1	A	168	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.