

# wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 08:19 pm BST

odec
odec

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment is 26%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f NMR} \ {f archive} \ (\#{f Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	А	12	75%	25%			
1	В	12	58%	42%			
1	С	12	58%	42%			
1	D	12	75%	25%			



# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues							
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model							
1	/	B:111-B:122, D:311-D:322	1.18	1			

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 9, 10, 11, 13, 14, 16, 18, 20
2	3, 17, 19
3	8, 12
Single-model clusters	7; 15



#### 7ACB

# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 872 atoms, of which 460 are hydrogens and 0 are deuteriums.

Mol	Chain	Residues		Atoms					Trace
1 1	1 A	12	Total	С	Η	Ν	0	S	0
	A	12	218	66	115	22	13	2	0
1	В	12	Total	С	Η	Ν	Ο	S	0
	D	12	218	66	115	22	13	2	0
1	C	12	Total	С	Н	Ν	Ο	S	0
	U	12	218	66	115	22	13	2	0
1	Л	12	Total	С	Η	Ν	Ο	S	0
	D	12	218	66	115	22	13	2	0

• Molecule 1 is a protein called Capra hircus Cathelicidin-1 (dodecylphosphocholine).



# 4 Residue-property plots (i)

# 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Capra hircus Cathelicidin-1 (dodecylphosphocholine)

Chain A:	75%	25%
R11 L17 L17 R19 R20 R21 R22 R22		
• Molecule 1: Cap	ra hircus Cathelicidin-1 (dodeo	cylphosphocholine)
Chain B:	58%	42%
R111 112 C113 C113 F116 V116 L117 R122 R122		
• Molecule 1: Cap	ra hircus Cathelicidin-1 (dodeo	cylphosphocholine)
Chain C:	58%	42%
R211 1212 C213 7215 7215 1217 1217 1217 1217 1217 1217		
• Molecule 1: Cap	ra hircus Cathelicidin-1 (dodeo	cylphosphocholine)
Chain D:	75%	25%
R311 1317 1318 R319 C321 R322		

# 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: Capra hircus Cathelicidin-1 (dodecylphosphocholine)



Chain A:	58%	42%	
R11 14 14 118 118 118 118 118 118			
• Molecule 1:	Capra hircus Cathelicidin-1	(dodecylphosphocholine)	
Chain B:	33%	67%	
R111 112 1112 0113 7115 F115 1118 1118 1118 R119	11.73 11.12		
• Molecule 1:	Capra hircus Cathelicidin-1	(dodecylphosphocholine)	
Chain C:	33%	67%	
R211 1212 7213 7214 7215 1217 1218 1218 1218			
• Molecule 1:	Capra hircus Cathelicidin-1	(dodecylphosphocholine)	
Chain D:	67%	25%	8%
R311 (314 (314 (314 (314 (314) (318) (318) (318) (318) (318) (318) (321) (321) (321)			



# 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: torsion angle dynamics.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	3.98.13
CYANA	structure calculation	3.98.13

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	2
Total number of shifts	207
Number of shifts mapped to atoms	207
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	26%



# 6 Model quality (i)

# 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

# 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	103	115	114	$5\pm 2$
1	В	103	115	114	3±1
1	С	103	115	114	$3\pm 2$
1	D	103	115	114	$5\pm 2$
All	All	8240	9200	9120	180

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

5 of 86 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:17:LEU:HD23	1:D:317:LEU:HD23	0.85	1.48	1	4
1:A:17:LEU:HD23	1:D:317:LEU:HD12	0.72	1.60	19	1
1:C:217:LEU:HD22	1:D:317:LEU:HD13	0.67	1.66	12	1
1:C:217:LEU:HD22	1:D:317:LEU:HD12	0.66	1.68	4	2
1:A:17:LEU:HD22	1:D:317:LEU:HD23	0.66	1.66	20	3



# 6.3 Torsion angles (i)

## 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perc	entiles
1	А	10/12~(83%)	$9{\pm}1$ ( $93{\pm}6\%$ )	$1 \pm 1 \ (7 \pm 6\%)$	0±0 (0±0%)	100	100
1	В	10/12~(83%)	$8\pm1$ ( $84\pm7\%$ )	$1{\pm}1$ (8 ${\pm}10\%$ )	1±0 (8±4%)	2	14
1	С	10/12~(83%)	$8\pm1$ (78 $\pm9\%$ )	$2\pm1$ (16 $\pm12\%$ )	$1\pm0~(6\pm5\%)$	3	20
1	D	10/12~(83%)	$9{\pm}1$ ( $94{\pm}7\%$ )	$1 \pm 1 \ (6 \pm 7\%)$	0±0 (0±0%)	100	100
All	All	800/960~(83%)	695~(87%)	77 (10%)	28 (4%)	6	35

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	В	113	CYS	16
1	С	213	CYS	12

## 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	entiles
1	А	12/12~(100%)	$11 \pm 1 (91 \pm 8\%)$	$1{\pm}1$ (9 ${\pm}8\%$ )	13	59
1	В	12/12~(100%)	$10{\pm}1$ (84 ${\pm}9\%$ )	$2\pm1 (16\pm9\%)$	5	41
1	С	12/12~(100%)	$10{\pm}1$ (82 ${\pm}9\%$ )	$2\pm1 (18\pm9\%)$	4	37
1	D	12/12~(100%)	$10\pm1~(87\pm9\%)$	$2\pm1~(13\pm9\%)$	7	49
All	All	960/960~(100%)	824 (86%)	136 (14%)	6	46

5 of 25 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.



Mol	Chain	Res	Type	Models (Total)
1	С	222	ARG	15
1	D	318	ILE	14
1	С	215	PHE	13
1	В	115	PHE	12
1	В	122	ARG	10

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry (i)

There are no ligands in this entry.

## 6.7 Other polymers (i)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 26% for the well-defined parts and 26% for the entire structure.

# 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_1

# 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	109
Number of shifts mapped to atoms	109
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

## 7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

## 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 13%, i.e. 109 atoms were assigned a chemical shift out of a possible 808. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}\mathbf{N}$
Backbone	34/240~(14%)	23/96~(24%)	0/96~(0%)	11/48~(23%)
Sidechain	73/528~(14%)	69/344~(20%)	0/144~(0%)	4/40~(10%)
Aromatic	2/40~(5%)	2/20~(10%)	0/20~(0%)	0/0 (%)
Overall	109/808~(13%)	94/460~(20%)	0/260~(0%)	15/88~(17%)

# 7.1.4 Statistically unusual chemical shifts (i)


There are no statistically unusual chemical shifts.



# 7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



# 7.2 Chemical shift list 2

File name: working\_cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_2

# 7.2.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	98
Number of shifts mapped to atoms	98
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

# 7.2.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).



#### 7.2.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 12%, i.e. 98 atoms were assigned a chemical shift out of a possible 808. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	33/240~(14%)	22/96~(23%)	0/96~(0%)	11/48~(23%)
Sidechain	63/528~(12%)	60/344~(17%)	0/144~(0%)	3/40~(8%)
Aromatic	2/40~(5%)	2/20~(10%)	0/20~(0%)	0/0 (%)
Overall	98/808~(12%)	84/460~(18%)	0/260~(0%)	14/88~(16%)


#### 7.2.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

# 7.2.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:





# 8 NMR restraints analysis (i)

# 8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	705
Intra-residue ( i-j =0)	298
Sequential ( i-j =1)	152
Medium range ( $ i-j >1$ and $ i-j <5$ )	12
Long range $( i-j  \ge 5)$	0
Inter-chain	107
Hydrogen bond restraints	112
Disulfide bond restraints	24
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	14.7
Number of long range restraints per residue <sup>1</sup>	0.0

<sup>1</sup>Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

# 8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

## 8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	20.6	0.2
0.2-0.5 (Medium)	19.4	0.5
>0.5 (Large)	25.1	9.92



## 8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than  $1^\circ$  are not included in the calculation. There are no dihedral-angle violations



# 9 Distance violation analysis (i)


# 9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destusiate toma	Count	$\%^1$	Vi	iolated	3	Consistently Violated		
Restraints type	Count	%0 <sup>-1</sup>	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue ( i-j =0)	298	42.3	75	25.2	10.6	18	6.0	2.6
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	202	28.7	51	25.2	7.2	17	8.4	2.4
Sidechain-Sidechain	96	13.6	24	25.0	3.4	1	1.0	0.1
Sequential ( i-j =1)	152	21.6	32	21.1	4.5	13	8.6	1.8
Backbone-Backbone	52	7.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	98	13.9	30	30.6	4.3	13	13.3	1.8
Sidechain-Sidechain	2	0.3	2	100.0	0.3	0	0.0	0.0
Medium range ( $ i-j  > 1 \&  i-j  < 5$ )	12	1.7	2	16.7	0.3	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	8	1.1	2	25.0	0.3	0	0.0	0.0
Sidechain-Sidechain	4	0.6	0	0.0	0.0	0	0.0	0.0
Long range $( i-j  \ge 5)$	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	107	15.2	19	17.8	2.7	2	1.9	0.3
Backbone-Backbone	70	9.9	10	14.3	1.4	2	2.9	0.3
Backbone-Sidechain	37	5.2	9	24.3	1.3	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	112	15.9	3	2.7	0.4	0	0.0	0.0
Disulfide bond	24	3.4	1	4.2	0.1	0	0.0	0.0
Total	705	100.0	132	18.7	18.7	33	4.7	4.7
Backbone-Backbone	234	33.2	13	5.6	1.8	2	0.9	0.3
Backbone-Sidechain	345	48.9	92	26.7	13.0	30	8.7	4.3
Sidechain-Sidechain	126	17.9	27	21.4	3.8	1	0.8	0.1

 $^1$  percentage calculated with respect to the total number of distance restraints,  $^2$  percentage calculated with respect to the number of restraints in a particular restraint category,  $^3$  violated in at least one model,  $^4$  violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

# 9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID	Number of violations						Moon (Å)	Mov (Å)	$SD^6$ (Å)	Median (Å)
Model ID	$\mathrm{IR}^{1}$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (Å)	Max (Å)	$SD^{*}(A)$	Median (A)
1	41	20	0	0	4	65	1.17	8.99	1.84	0.35
2	37	23	0	0	3	63	1.19	9.52	1.9	0.36
3	37	18	0	0	6	61	1.24	9.45	1.9	0.42
4	41	20	0	0	4	65	1.19	9.0	1.83	0.33
5	37	21	0	0	7	65	1.19	9.02	1.84	0.31
6	40	22	1	0	9	72	1.12	9.27	1.82	0.31
7	40	20	0	0	4	64	1.2	9.92	1.93	0.34
8	41	22	0	0	3	66	1.18	9.28	1.81	0.4
9	36	23	0	0	5	64	1.18	9.05	1.85	0.39
10	39	18	0	0	8	65	1.16	8.82	1.79	0.39
11	38	24	0	0	6	68	1.14	9.56	1.78	0.32


Continued on next page...




Madal ID	Number of violations						Mean (Å)	N/ (Å)	$SD^6$ (Å)	Madian (Å)
Model ID	$\mathrm{IR}^{1}$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (A)	Max (Å)	$SD^{\circ}(A)$	Median (Å)
12	40	19	1	0	9	69	1.11	9.1	1.76	0.35
13	38	20	0	0	11	69	1.14	9.49	1.79	0.26
14	41	17	0	0	8	66	1.18	9.2	1.8	0.37
15	35	23	0	0	6	64	1.17	8.72	1.8	0.32
16	39	21	0	0	8	68	1.18	9.31	1.88	0.32
17	35	20	0	0	6	61	1.25	8.82	1.88	0.29
18	36	19	0	0	7	62	1.21	9.85	1.85	0.42
19	39	18	0	0	8	65	1.16	8.67	1.8	0.34
20	38	18	0	0	5	61	1.23	9.27	1.84	0.46

Continued from previous page...

<sup>1</sup>Intra-residue restraints, <sup>2</sup>Sequential restraints, <sup>3</sup>Medium range restraints, <sup>4</sup>Long range restraints, <sup>5</sup>Inter-chain restraints, <sup>6</sup>Standard deviation





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right




# 9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 441(IR:223, SQ:120, MR:10, LR:0, IC:88) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restra	aints	Fractio	n of the ensemble
$IR^1$	$SQ^2$	$MR^3$	LR <sup>4</sup>	IC <sup>5</sup>	Total	$\operatorname{Count}^6$	%
18	2	2	0	6	28	1	5.0
12	1	0	0	1	14	2	10.0
2	2	0	0	1	5	3	15.0
1	3	0	0	2	6	4	20.0
2	0	0	0	3	5	5	25.0
3	0	0	0	0	3	6	30.0
0	2	0	0	1	3	7	35.0
0	1	0	0	1	2	8	40.0
0	2	0	0	0	2	9	45.0
1	1	0	0	1	3	10	50.0
0	1	0	0	0	1	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
1	2	0	0	0	3	14	70.0
2	0	0	0	0	2	15	75.0
2	1	0	0	1	4	16	80.0
0	0	0	0	0	0	17	85.0
5	0	0	0	0	5	18	90.0
8	1	0	0	0	9	19	95.0
18	13	0	0	2	33	20	100.0

 $^{1}$ Intra-residue restraints,  $^{2}$ Sequential restraints,  $^{3}$ Medium range restraints,  $^{4}$ Long range restraints,  $^{5}$ Inter-chain restraints,  $^{6}$  Number of models with violations






## 9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

# 9.4 Most violated distance restraints in the ensemble (i)

## 9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





#### 9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	$\mathbf{Models}^1$	Mean (Å)	$SD^1$ (Å)	Median (Å)
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	20	8.84	0.57	8.96
(1,541)	1:A:21:CYS:HA	1:D:313:CYS:H	20	8.82	0.58	8.83
(1,561)	1:C:221:CYS:HB2	1:C:222:ARG:HA	20	5.28	0.05	5.29
(1,559)	1:B:121:CYS:HB2	1:B:122:ARG:HA	20	5.16	0.1	5.12
(1,547)	1:B:117:LEU:HD11	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,547)	1:B:117:LEU:HD12	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,547)	1:B:117:LEU:HD13	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,547)	1:B:117:LEU:HD21	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,547)	1:B:117:LEU:HD22	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,547)	1:B:117:LEU:HD23	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,551)	1:B:117:LEU:HD11	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,551)	1:B:117:LEU:HD12	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,551)	1:B:117:LEU:HD13	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,551)	1:B:117:LEU:HD21	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,551)	1:B:117:LEU:HD22	1:B:118:ILE:H	20	3.27	0.17	3.2
(1,551)	1:B:117:LEU:HD23	1:B:118:ILE:H	20	3.27	0.17	3.2

Continued on next page...



Key	Atom-1	Atom-2	$\mathbf{Models}^1$	Mean (Å)	$SD^1$ (Å)	Median (Å)
(1,555)	1:C:216:VAL:HG11	1:C:217:LEU:HA	20	3.26	0.07	3.26
(1,555)	1:C:216:VAL:HG12	1:C:217:LEU:HA	20	3.26	0.07	3.26
(1,555)	1:C:216:VAL:HG13	1:C:217:LEU:HA	20	3.26	0.07	3.26
(1,555)	1:C:216:VAL:HG21	1:C:217:LEU:HA	20	3.26	0.07	3.26
(1,555)	1:C:216:VAL:HG22	1:C:217:LEU:HA	20	3.26	0.07	3.26
(1,555)	1:C:216:VAL:HG23	1:C:217:LEU:HA	20	3.26	0.07	3.26
(1,557)	1:B:116:VAL:HG11	1:B:117:LEU:HA	20	3.23	0.06	3.24
(1,557)	1:B:116:VAL:HG12	1:B:117:LEU:HA	20	3.23	0.06	3.24
(1,557)	1:B:116:VAL:HG13	1:B:117:LEU:HA	20	3.23	0.06	3.24
(1,557)	1:B:116:VAL:HG21	1:B:117:LEU:HA	20	3.23	0.06	3.24
(1,557)	1:B:116:VAL:HG22	1:B:117:LEU:HA	20	3.23	0.06	3.24
(1,557)	1:B:116:VAL:HG23	1:B:117:LEU:HA	20	3.23	0.06	3.24
(1,549)	1:C:217:LEU:HD11	1:C:218:ILE:H	20	3.19	0.2	3.18
(1,549)	1:C:217:LEU:HD12	1:C:218:ILE:H	20	3.19	0.2	3.18
(1,549)	1:C:217:LEU:HD13	1:C:218:ILE:H	20	3.19	0.2	3.18
(1,549)	1:C:217:LEU:HD21	1:C:218:ILE:H	20	3.19	0.2	3.18
(1,549)	1:C:217:LEU:HD22	1:C:218:ILE:H	20	3.19	0.2	3.18
(1,549)	1:C:217:LEU:HD23	1:C:218:ILE:H	20	3.19	0.2	3.18
(1,553)	1:C:217:LEU:HD11	1:C:218:ILE:H	20	3.19	0.2	3.18

Continued from previous page...


<sup>1</sup>Number of violated models, <sup>2</sup>Standard deviation

# 9.5 All violated distance restraints (i)

## 9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





## 9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,541)	1:A:21:CYS:HA	1:D:313:CYS:H	7	9.92
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	18	9.85
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	11	9.56
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	2	9.52
(1,541)	1:A:21:CYS:HA	1:D:313:CYS:H	13	9.49
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	3	9.45
(1,541)	1:A:21:CYS:HA	1:D:313:CYS:H	16	9.31
(1,541)	1:A:21:CYS:HA	1:D:313:CYS:H	8	9.28
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	6	9.27
(1,539)	1:A:13:CYS:H	1:D:321:CYS:HA	20	9.27



# 10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

