



## wwPDB EM Validation Summary Report ⓘ

Apr 5, 2023 – 02:03 pm BST

PDB ID : 8ACQ  
EMDB ID : EMD-14715  
Title : S-layer Deinoxanthin-Binding Complex (SDBC), subunit DR\_2577 assembled with its SOD DR\_0644  
Authors : Farci, D.; Piano, D.  
Deposited on : 2022-07-06  
Resolution : 2.54 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev50  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.2

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1167	
1	B	1167	
1	C	1167	
2	H	206	
2	I	206	
2	L	206	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	JPI	A	1201	X	-	-	-
3	JPI	B	1302	X	-	-	-
3	JPI	C	1201	X	-	-	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 25170 atoms, of which 714 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called S-layer protein SlpA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	949	Total	C	N	O	S	0	0
			7117	4415	1265	1431	6		
1	B	949	Total	C	N	O	S	0	0
			7117	4415	1265	1431	6		
1	C	949	Total	C	N	O	S	0	0
			7117	4415	1265	1431	6		

- Molecule 2 is a protein called DR\_0644, only-Cu Superoxide Dismutase.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	H	120	Total	C	N	O	S	0	0
			864	547	147	165	5		
2	I	120	Total	C	N	O	S	0	0
			864	547	147	165	5		
2	L	120	Total	C	N	O	S	0	0
			864	547	147	165	5		

- Molecule 3 is (3 {S},5 {R},6 {R})-5-[(3 {S},7 {R},12 {S},16 {S},20 {S})-3,7,12,16,20,24-hexamethyl-24-oxidanyl-pentacosyl]-4,4,6-trimethyl-cyclohexane-1,3-diol (three-letter code: JPI) (formula: C<sub>40</sub>H<sub>80</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf	
			Total	C	H		O
3	A	1	119	40	76	3	0
3	B	1	119	40	76	3	0
3	C	1	119	40	76	3	0

- Molecule 4 is [(2 {S})-2-acetyloxy-3-[[[(2 {S})-3-[(2 {R},3 {S},4 {S},5 {S},6 {S})-6-(hydroxymethyl)-3-(octadecanoylamino)-4,5-bis(oxidanyl)oxan-2-yl]oxy-1-oxidanylidene-1-(pentylamino)propan-2-yl]oxy-oxidanyl-phosphoryl]oxy-propyl] ethanoate (three-letter code: JQ6) (formula: C<sub>39</sub>H<sub>73</sub>N<sub>2</sub>O<sub>15</sub>P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf	
4	A	1	Total	C	H	N	O	P	0
			129	39	72	2	15	1	
4	B	1	Total	C	H	N	O	P	0
			129	39	72	2	15	1	
4	C	1	Total	C	H	N	O	P	0
			129	39	72	2	15	1	

- Molecule 5 is [(2 {S})-3-[[[(2 {S})-3-[(2 {S},3 {S},4 {S},5 {S},6 {S})-6-(hydroxymethyl)-4,5-bis(oxidanyl)-3-(propanoylamino)oxan-2-yl]oxy-1-oxidanylidene-1-(pentadecylamino)propan-2-yl]oxy-oxidanyl-phosphoryl]oxy-2-octanoyloxy-propyl] decanoate (three-letter code: JPX) (formula: C<sub>48</sub>H<sub>91</sub>N<sub>2</sub>O<sub>15</sub>P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf	
5	A	1	Total	C	H	N	O	P	0
			156	48	90	2	15	1	
5	B	1	Total	C	H	N	O	P	0
			156	48	90	2	15	1	
5	B	1	Total	C	H	N	O	P	0
			156	48	90	2	15	1	

- Molecule 6 is COPPER (II) ION (three-letter code: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
6	A	3	Total	Cu	0
			3	3	
6	B	3	Total	Cu	0
			3	3	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		AltConf
6	C	3	Total 3	Cu 3	0
6	H	1	Total 1	Cu 1	0
6	I	1	Total 1	Cu 1	0
6	L	1	Total 1	Cu 1	0


- Molecule 7 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
7	A	1	Total 1	Fe 1	0
7	B	1	Total 1	Fe 1	0
7	C	1	Total 1	Fe 1	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: S-layer protein SlpA










• Molecule 1: S-layer protein SlpA






• Molecule 1: S-layer protein SlpA







• Molecule 2: DR\_0644, only-Cu Superoxide Dismutase



• Molecule 2: DR\_0644, only-Cu Superoxide Dismutase



• Molecule 2: DR\_0644, only-Cu Superoxide Dismutase



THR	THR
ALA	ALA
ALA	ALA
PRO	PRO
MET	MET
GLY	GLY
SER	SER
ALA	ALA
PRO	PRO
ALA	ALA
THR	THR
ASN	ASN
THR	THR
ASN	ASN
MET	MET
ALA	ALA
MET	MET
SER	SER
SER	SER
THR	THR
ASP	ASP
ALA	ALA
THR	THR
ALA	ALA
THR	THR
THR	THR
THR	THR
ALA	ALA
SER	SER
THR	THR
THR	THR
SER	SER
THR	THR
THR	THR
THR	THR
SER	SER
THR	THR
THR	THR
ASP	ASP
MET	MET
THR	THR
THR	THR
THR	THR
VAL	VAL
THR	THR
THR	THR
VAL	VAL
ALA	ALA
ALA	ALA
SER	SER
THR	THR
ASP	ASP
MET	MET
THR	THR
THR	THR
THR	THR
VAL	VAL
THR	THR
MET	MET
ALA	ALA
PRO	PRO
VAL	VAL
THR	THR
ALA	ALA
ALA	ALA
PRO	PRO
ASN	ASN
PRO	PRO

CYS	THR
ASN	ALA
S142	S142
D143	D143
G144	G144
P145	P145
A146	A146
I147	I147
M148	M148
E149	E149
S150	S150
R151	R151
Q155	Q155
A156	A156
D159	D159
V162	V162
T163	T163
P169	P169
L172	L172
I173	I173
R174	R174
D175	D175
I179	I179
H183	H183
G184	G184
R185	R185
D186	D186
F187	F187
S188	S188
G189	G189
A190	A190
D193	D193
S194	S194
G195	G195
V196	V196
I197	I197
C198	C198
T199	T199
P200	P200
I201	I201
THR	THR
MET	MET
THR	THR
MET	MET
ARG	ARG

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C3	Depositor
Number of particles used	252122	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.3	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1000	Depositor
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	2.993	Depositor
Minimum map value	-1.662	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.060	Depositor
Recommended contour level	0.46	Depositor
Map size (Å)	431.90402, 431.90402, 431.90402	wwPDB
Map dimensions	528, 528, 528	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.818, 0.818, 0.818	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JPI, JPX, CU, FE, JQ6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/7258	0.53	0/9865
1	B	0.29	0/7258	0.53	0/9865
1	C	0.29	0/7258	0.53	0/9865
2	H	0.72	0/882	0.84	0/1205
2	I	0.73	0/882	0.80	0/1205
2	L	0.67	0/882	0.76	0/1205
All	All	0.36	0/24420	0.57	0/33210

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	696	TYR	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7117	0	6745	350	0
1	B	7117	0	6745	352	0
1	C	7117	0	6745	362	0
2	H	864	0	871	22	0
2	I	864	0	871	9	0
2	L	864	0	871	28	0
3	A	43	76	0	7	0
3	B	43	76	0	5	0
3	C	43	76	0	1	0
4	A	57	72	0	3	0
4	B	57	72	0	3	0
4	C	57	72	0	7	0
5	A	66	90	0	8	0
5	B	132	180	0	11	0
6	A	3	0	0	0	0
6	B	3	0	0	0	0
6	C	3	0	0	0	0
6	H	1	0	0	0	0
6	I	1	0	0	0	0
6	L	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
All	All	24456	714	22848	1046	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 1046 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1301:JPX:C30	5:B:1301:JPX:O15	1.82	1.28
4:A:1202:JQ6:C20	5:A:1203:JPX:C50	2.23	1.15
1:B:247:ALA:HB1	2:L:187:PHE:CE2	1.88	1.08
5:A:1203:JPX:O15	5:A:1203:JPX:C31	2.00	1.08
4:A:1202:JQ6:C21	5:A:1203:JPX:C16	2.33	1.05

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	947/1167 (81%)	848 (90%)	97 (10%)	2 (0%)	47	60
1	B	947/1167 (81%)	846 (89%)	99 (10%)	2 (0%)	47	60
1	C	947/1167 (81%)	850 (90%)	95 (10%)	2 (0%)	47	60
2	H	116/206 (56%)	97 (84%)	19 (16%)	0	100	100
2	I	116/206 (56%)	89 (77%)	27 (23%)	0	100	100
2	L	116/206 (56%)	96 (83%)	20 (17%)	0	100	100
All	All	3189/4119 (77%)	2826 (89%)	357 (11%)	6 (0%)	50	60

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	818	GLN
1	B	818	GLN
1	C	818	GLN
1	B	1087	ASN
1	A	1087	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	726/891 (82%)	724 (100%)	2 (0%)	92	96
1	B	726/891 (82%)	724 (100%)	2 (0%)	92	96
1	C	726/891 (82%)	724 (100%)	2 (0%)	92	96

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	H	89/157 (57%)	88 (99%)	1 (1%)	73	83
2	I	89/157 (57%)	88 (99%)	1 (1%)	73	83
2	L	89/157 (57%)	89 (100%)	0	100	100
All	All	2445/3144 (78%)	2437 (100%)	8 (0%)	92	96

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	74	HIS
2	H	52	THR
1	C	360	LYS
1	B	368	ARG
1	C	368	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1054	ASN
1	C	493	ASN
1	B	596	ASN
1	B	493	ASN
1	C	240	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 24 ligands modelled in this entry, 15 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	JQ6	C	1202	-	56,57,57	0.25	0	66,71,71	0.67	2 (3%)
4	JQ6	B	1304	5	56,57,57	0.28	0	66,71,71	0.53	0
3	JPI	C	1201	-	43,43,43	2.29	3 (6%)	55,58,58	2.10	8 (14%)
5	JPX	B	1303	-	65,66,66	0.37	0	75,80,80	1.27	7 (9%)
3	JPI	A	1201	-	43,43,43	2.29	3 (6%)	55,58,58	2.03	8 (14%)
5	JPX	B	1301	4	65,66,66	0.47	1 (1%)	75,80,80	0.76	3 (4%)
3	JPI	B	1302	-	43,43,43	2.25	3 (6%)	55,58,58	2.04	8 (14%)
5	JPX	A	1203	-	65,66,66	0.46	1 (1%)	75,80,80	0.72	2 (2%)
4	JQ6	A	1202	-	56,57,57	0.30	0	66,71,71	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	JQ6	C	1202	-	-	27/58/78/78	0/1/1/1
4	JQ6	B	1304	5	-	24/58/78/78	0/1/1/1
3	JPI	C	1201	-	4/4/11/11	16/37/61/61	0/1/1/1
5	JPX	B	1303	-	-	31/69/89/89	0/1/1/1
3	JPI	A	1201	-	4/4/11/11	10/37/61/61	0/1/1/1
5	JPX	B	1301	4	-	38/69/89/89	0/1/1/1
3	JPI	B	1302	-	4/4/11/11	18/37/61/61	0/1/1/1
5	JPX	A	1203	-	-	34/69/89/89	0/1/1/1
4	JQ6	A	1202	-	-	31/58/78/78	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1201	JPI	C5-C6	-11.82	1.34	1.54
3	A	1201	JPI	C5-C6	-11.70	1.34	1.54
3	B	1302	JPI	C5-C6	-11.39	1.35	1.54
3	A	1201	JPI	O1-C4	-8.79	1.24	1.43
3	B	1302	JPI	O1-C4	-8.66	1.25	1.43

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1201	JPI	C7-C5-C6	9.28	126.14	113.04
3	C	1201	JPI	C7-C5-C6	9.22	126.06	113.04
3	B	1302	JPI	C7-C5-C6	8.58	125.15	113.04
5	B	1303	JPX	O6-C4-O5	-5.99	95.79	109.97
3	B	1302	JPI	O1-C4-C5	5.60	122.18	110.66

5 of 12 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	1201	JPI	C5
3	A	1201	JPI	C4
3	A	1201	JPI	C2
3	A	1201	JPI	C6
3	B	1302	JPI	C5

5 of 229 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1302	JPI	C6-C10-C11-C12
3	B	1302	JPI	C11-C10-C6-C1
3	C	1201	JPI	C11-C10-C6-C5
4	A	1202	JQ6	O1-C1-C2-O5
4	A	1202	JQ6	C2-C1-C3-O7

There are no ring outliers.

9 monomers are involved in 40 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1202	JQ6	7	0
4	B	1304	JQ6	3	0
3	C	1201	JPI	1	0
5	B	1303	JPX	3	0
3	A	1201	JPI	7	0


*Continued on next page...*



Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1301	JPX	8	0
3	B	1302	JPI	5	0
5	A	1203	JPX	8	0
4	A	1202	JQ6	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14715. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 264




Y Index: 264




Z Index: 264


### 6.2.2 Raw map



X Index: 264



Y Index: 264




Z Index: 264


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 268




Y Index: 272




Z Index: 246


### 6.3.2 Raw map



X Index: 268



Y Index: 271




Z Index: 247


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.46. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [\(i\)](#)




The volume at the recommended contour level is 131 nm<sup>3</sup>; this corresponds to an approximate mass of 118 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.394 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.394 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.54	-	-
Author-provided FSC curve	2.54	2.93	2.61
Unmasked-calculated*	6.88	12.11	7.72

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.88 differs from the reported value 2.54 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14715 and PDB model 8ACQ. Per-residue inclusion information can be found in section 3 on page 8.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.46 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.46).















## 9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.46) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8270	 0.5790
A	 0.8730	 0.5900
B	 0.8740	 0.5910
C	 0.8700	 0.5890
H	 0.4570	 0.5020
I	 0.4560	 0.4930
L	 0.4430	 0.4710

