

Full wwPDB X-ray Structure Validation Report (i)

Jun 23, 2024 – 01:39 PM EDT

PDB ID : 5ADJ

Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex

with 7-((3-(2-(Methylamino)ethyl)phenoxy)methyl)quinolin-2- amine

Authors : Li, H.; Poulos, T.L.

Deposited on : 2015-08-20

Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	443	82%	9%	9%			
1	В	443	81%	9%	9%			



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 6815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total 3220	As 1	C 2047	N 565	O 591	S 16	0	1	0
1	В	403	Total 3212	As 1	C 2042	N 566	O 587	S 16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	conflict	UNP P29473
В	100	ARG	CYS	conflict	UNP P29473

• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0



Continued from previous page...

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	D	1	Total	С	Fe	N	О	0	0
2	Ъ	1	43	34	1	4	4	0	U


• Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 17 9 5 3	0	0
3	В	1	Total C N O 17 9 5 3	0	0


• Molecule 4 is 7-[[3-[2-(methylamino)ethyl]phenoxy]methyl]quinolin-2-amine (three-letter code: M94) (formula: $C_{19}H_{21}N_3O$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 23 19 3 1	0	0
4	В	1	Total C N O 23 19 3 1	0	0

 \bullet Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 4 2 2	0	0
5	В	1	Total C O 4 2 2	0	0

• Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	В	1	Total C O 6 3 3	0	0

• Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Zn 1 1	0	0

• Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	105	Total O 105 105	0	0
8	В	83	Total O 83 83	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	57.61Å 105.81Å 156.31Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.14 - 2.22	Depositor
rtesolution (A)	48.14 - 2.22	EDS
% Data completeness	99.5 (48.14-2.22)	Depositor
(in resolution range)	99.5 (48.14-2.22)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.16 (at 2.22Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
P. P.	0.164 , 0.210	Depositor
R, R_{free}	0.164 , 0.210	DCC
R_{free} test set	2398 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	50.7	Xtriage
Anisotropy	0.536	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 46.3	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6815	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: H4B, GOL, M94, ACT, CAS, HEM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.43	0/3300	0.55	0/4494
1	В	0.41	0/3291	0.54	0/4480
All	All	0.42	0/6591	0.54	0/8974

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3220	0	3117	21	0
1	В	3212	0	3117	22	0
2	A	43	0	30	2	0
2	В	43	0	30	2	0
3	A	17	0	15	0	0
3	В	17	0	15	1	0
4	A	23	0	21	2	0
4	В	23	0	21	4	0
5	A	8	0	6	1	0
5	В	8	0	6	0	0
6	A	6	0	8	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
6	В	6	0	8	1	0
7	A	1	0	0	0	0
8	A	105	0	0	1	0
8	В	83	0	0	1	0
All	All	6815	0	6394	44	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	$-$ overlap (\mathring{A})
1:B:249:GLN:HE21	4:B:800:M94:H22	1.45	0.81
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.71	0.72
1:B:338:VAL:HG21	4:B:800:M94:C07	2.30	0.62
2:B:500:HEM:O2D	8:B:2081:HOH:O	2.16	0.61
1:A:338:VAL:HG21	4:A:800:M94:C07	2.31	0.60
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	1.86	0.57
1:B:370:CYS:HB3	1:B:378:LEU:HD22	1.86	0.57
1:A:236:ARG:NH1	1:A:349:GLU:OE1	2.38	0.57
4:B:800:M94:H303	6:B:880:GOL:H32	1.86	0.56
1:A:355:PHE:HE1	4:A:800:M94:H06	1.69	0.56
1:A:68:PRO:HG2	1:A:83:TYR:CE2	2.41	0.56
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.87	0.56
1:A:252:ARG:NH2	5:A:861:ACT:O	2.39	0.53
1:A:126:LEU:HD11	1:A:156:GLU:HA	1.91	0.53
1:B:167:LEU:HG	1:B:348:LEU:HD12	1.93	0.51
1:A:281:TRP:HB2	1:A:304:LEU:HD21	1.94	0.49
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.94	0.49
1:A:273:THR:O	1:A:277:ILE:HG13	2.16	0.46
1:B:130:ARG:NH2	1:B:156:GLU:OE2	2.48	0.46
1:A:392:THR:HB	1:B:423:HIS:HB2	1.97	0.46
1:A:186:CYS:HB2	2:A:500:HEM:ND	2.32	0.45
8:A:2076:HOH:O	1:B:463:HIS:ND1	2.36	0.45
1:B:366:THR:O	1:B:370:CYS:HB2	2.17	0.45
1:A:154:GLU:OE1	1:A:168:ARG:NH2	2.50	0.45
1:B:236:ARG:HD2	1:B:242:ASP:OD1	2.17	0.45
1:B:236:ARG:HD3	1:B:351:SER:HB3	2.00	0.44
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.57	0.43
1:A:126:LEU:HA	1:A:126:LEU:HD12	1.79	0.43
1:B:242:ASP:HB3	1:B:351:SER:OG	2.17	0.43



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:A:423:HIS:HB2	1:B:392:THR:HB	2.01	0.43
1:B:319:HIS:CG	1:B:320:PRO:HD2	2.53	0.42
1:A:237:ALA:HB3	1:A:240:ARG:HB3	2.00	0.42
1:A:240:ARG:HD2	1:A:241:GLY:O	2.20	0.42
1:B:355:PHE:HE1	4:B:800:M94:H06	1.84	0.42
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	2.00	0.42
1:B:126:LEU:HD21	1:B:156:GLU:HG3	2.02	0.42
1:A:480:ASP:HA	1:A:481:PRO:HD3	1.79	0.41
1:B:367:ARG:O	1:B:371:ASP:HB2	2.20	0.41
1:B:180:TRP:CE3	1:B:192:TRP:HA	2.55	0.41
1:B:449:TRP:HA	3:B:600:H4B:N1	2.34	0.41
1:A:231:THR:O	1:A:353:ALA:HA	2.21	0.41
1:B:183:ALA:HA	1:B:184:PRO:HD3	1.84	0.41
1:A:180:TRP:CE3	1:A:192:TRP:HA	2.57	0.40
1:B:70:PHE:HA	1:B:71:PRO:HD2	1.81	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	400/443 (90%)	392 (98%)	8 (2%)	0	100	100
1	В	398/443 (90%)	385 (97%)	13 (3%)	0	100	100
All	All	798/886 (90%)	777 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	343/375 (92%)	331 (96%)	12 (4%)	36 44		
1	В	342/375 (91%)	327 (96%)	15 (4%)	28 34		
All	All	685/750 (91%)	658 (96%)	27 (4%)	32 40		

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	125	LEU
1	A	169	GLU
1	A	259	GLN
1	A	260	ASP
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	458	LEU
1	В	69	LYS
1	В	91	GLN
1	В	100	ARG
1	В	102	LEU
1	В	124	GLN
1	В	141	LYS
1	В	143	SER
1	В	149	GLU
1	В	156	GLU
1	В	168	ARG
1	В	223	ARG
1	В	240	ARG
1	В	330	LEU
1	В	378	LEU
1	В	458	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	413	GLN
1	В	249	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Chain	Dag	Timle	B	ond leng	${ m gths}$	В	ond ang	gles
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
1	CAS	В	384	1	5,8,9	1.02	0	1,9,11	0.45	0
1	CAS	A	384	1	5,8,9	1.05	0	1,9,11	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	В	384	1	-	0/0/7/9	-
1	CAS	A	384	1	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Dag	Link	Во	ond leng	gths	В	ond ang	gles
Mol	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	H4B	A	600	-	16,18,18	0.91	0	11,26,26	2.75	6 (54%)
4	M94	В	800	-	25,25,25	0.89	1 (4%)	32,33,33	1.06	4 (12%)
5	ACT	В	860	-	3,3,3	0.77	0	3,3,3	0.66	0
2	HEM	A	500	1	41,50,50	1.84	7 (17%)	45,82,82	1.61	11 (24%)
5	ACT	A	861	-	3,3,3	0.83	0	3,3,3	0.75	0
3	H4B	В	600	-	16,18,18	0.97	1 (6%)	11,26,26	2.63	4 (36%)
5	ACT	В	861	-	3,3,3	0.83	0	3,3,3	0.61	0
5	ACT	A	860	_	3,3,3	0.76	0	3,3,3	0.74	0
6	GOL	В	880	_	5,5,5	0.31	0	5,5,5	0.41	0
2	HEM	В	500	1	41,50,50	2.00	7 (17%)	45,82,82	2.19	10 (22%)
4	M94	A	800	-	25,25,25	0.94	1 (4%)	32,33,33	1.00	1 (3%)
6	GOL	A	880	_	5,5,5	0.32	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
4	M94	В	800	-	-	2/9/9/9	0/3/3/3
2	HEM	A	500	1	-	1/12/54/54	-
3	H4B	В	600	-	-	0/8/17/17	0/2/2/2
6	GOL	В	880	-	-	2/4/4/4	-
2	HEM	В	500	1	-	0/12/54/54	_



$Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	M94	A	800	-	-	2/9/9/9	0/3/3/3
6	GOL	A	880	-	-	4/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\mathring{A}})$	$\operatorname{Ideal}(ext{\AA})$
2	В	500	HEM	C3D-C2D	7.96	1.53	1.36
2	A	500	HEM	C3D-C2D	7.11	1.51	1.36
2	В	500	HEM	C3C-C2C	-4.13	1.34	1.40
2	A	500	HEM	C3C-C2C	-4.05	1.34	1.40
2	В	500	HEM	C3C-CAC	3.92	1.55	1.47
2	A	500	HEM	C3C-CAC	3.50	1.55	1.47
2	A	500	HEM	CAB-C3B	2.94	1.55	1.47
2	В	500	HEM	CAB-C3B	2.89	1.55	1.47
2	В	500	HEM	FE-ND	2.82	2.10	1.96
2	В	500	HEM	FE-NB	2.76	2.10	1.96
2	A	500	HEM	CMD-C2D	2.20	1.55	1.50
2	A	500	HEM	CAA-C2A	2.13	1.55	1.52
4	A	800	M94	C02-N01	2.12	1.36	1.33
3	В	600	H4B	C4A-C4	-2.09	1.38	1.41
2	В	500	HEM	CMD-C2D	2.03	1.55	1.50
2	A	500	HEM	CMB-C2B	2.03	1.55	1.50
4	В	800	M94	C09-C10	-2.01	1.38	1.41

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	500	HEM	CBA-CAA-C2A	-7.03	100.62	112.62
2	В	500	HEM	C4D-ND-C1D	6.78	112.07	105.07
3	В	600	H4B	C8A-C4A-C4	5.72	119.65	114.57
3	A	600	H4B	C4-C4A-N5	4.47	122.88	119.12
2	A	500	HEM	C4D-ND-C1D	4.03	109.24	105.07
2	В	500	HEM	C4C-CHD-C1D	3.96	127.78	122.56
3	A	600	H4B	C8A-C4A-C4	3.94	118.07	114.57
3	A	600	H4B	N1-C2-N3	-3.84	119.39	125.42
3	В	600	H4B	C2-N3-C4	3.63	121.69	115.93
2	В	500	HEM	CBD-CAD-C3D	-3.47	103.00	112.63
3	A	600	H4B	C2-N3-C4	3.45	121.41	115.93
2	A	500	HEM	CMA-C3A-C4A	-3.45	123.17	128.46
2	В	500	HEM	C1B-NB-C4B	3.32	108.50	105.07
3	В	600	H4B	N1-C2-N3	-3.25	120.33	125.42
2	В	500	HEM	C4B-CHC-C1C	3.04	126.57	122.56



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	600	H4B	C2-N1-C8A	3.03	121.32	114.54
2	В	500	HEM	C3B-C2B-C1B	2.99	108.70	106.49
2	В	500	HEM	CHC-C4B-C3B	2.76	128.79	124.57
2	A	500	HEM	CHB-C1B-NB	2.65	127.66	124.38
2	A	500	HEM	CBA-CAA-C2A	-2.61	108.17	112.62
2	A	500	HEM	CAD-CBD-CGD	-2.58	108.05	113.60
2	A	500	HEM	CAD-C3D-C4D	2.49	129.00	124.66
2	A	500	HEM	C1D-C2D-C3D	-2.48	104.35	106.96
2	A	500	HEM	CMA-C3A-C2A	2.46	129.58	124.94
2	A	500	HEM	CBD-CAD-C3D	-2.37	106.04	112.63
2	В	500	HEM	CHA-C4D-ND	2.36	127.30	124.38
3	В	600	H4B	C2-N1-C8A	2.33	119.75	114.54
4	В	800	M94	C04-C03-C02	2.32	121.29	119.48
3	A	600	H4B	N2-C2-N3	2.24	120.73	117.25
4	В	800	M94	C03-C02-N01	-2.24	119.39	122.08
2	A	500	HEM	CMD-C2D-C1D	2.15	128.31	125.04
2	В	500	HEM	CMD-C2D-C1D	2.12	128.27	125.04
4	В	800	M94	C27-C28-N29	2.12	116.53	111.83
2	A	500	HEM	CMC-C2C-C3C	2.12	128.65	124.68
4	В	800	M94	C03-C04-C05	-2.06	117.64	120.82
4	A	800	M94	C28-C27-C25	-2.04	108.14	112.87

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	800	M94	C25-C27-C28-N29
6	A	880	GOL	O1-C1-C2-C3
6	A	880	GOL	C1-C2-C3-O3
6	В	880	GOL	C1-C2-C3-O3
6	В	880	GOL	O2-C2-C3-O3
4	A	800	M94	C27-C28-N29-C30
6	A	880	GOL	O2-C2-C3-O3
6	A	880	GOL	O1-C1-C2-O2
4	A	800	M94	C25-C27-C28-N29
2	A	500	HEM	CAD-CBD-CGD-O2D
4	В	800	M94	C08-C11-O12-C21


There are no ring outliers.

7 monomers are involved in 12 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	800	M94	4	0
2	A	500	HEM	2	0
5	A	861	ACT	1	0
3	В	600	H4B	1	0
6	В	880	GOL	1	0
2	В	500	HEM	2	0
4	A	800	M94	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	403/443 (90%)	0.23	21 (5%) 27 2	5	33, 51, 89, 112	0
1	В	402/443 (90%)	0.09	12 (2%) 50 4	8	35, 56, 88, 122	0
All	All	805/886 (90%)	0.16	33 (4%) 37 3	5	33, 53, 89, 122	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	259	GLN	4.4
1	В	261	GLY	3.8
1	A	259	GLN	3.7
1	A	126	LEU	3.7
1	В	123	GLU	3.5
1	A	91	GLN	3.3
1	A	448	ALA	3.3
1	A	311	LEU	2.9
1	A	69	LYS	2.9
1	В	270	VAL	2.8
1	A	122	ALA	2.7
1	A	156	GLU	2.7
1	В	448	ALA	2.7
1	A	103	GLY	2.6
1	В	447	TRP	2.6
1	В	263	VAL	2.5
1	A	121	PRO	2.5
1	В	323	GLU	2.5
1	A	153	GLN	2.5
1	В	122	ALA	2.5
1	A	160	ALA	2.4
1	В	144	GLY	2.2
1	A	260	ASP	2.2
1	В	274	GLU	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	257	ARG	2.2
1	A	155	VAL	2.2
1	A	452	PRO	2.2
1	A	186	CYS	2.2
1	A	274	GLU	2.2
1	A	157	ALA	2.1
1	В	260	ASP	2.1
1	A	187	VAL	2.1
1	A	449	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	CAS	A	384	9/10	0.96	0.14	47,53,109,109	0
1	CAS	В	384	9/10	0.96	0.12	67,71,138,197	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
5	ACT	В	861	4/4	0.87	0.41	55,59,62,63	0
6	GOL	A	880	6/6	0.87	0.28	78,87,90,93	0
6	GOL	В	880	6/6	0.89	0.15	69,72,73,74	0
4	M94	A	800	23/23	0.91	0.19	33,59,102,102	0
4	M94	В	800	23/23	0.91	0.20	37,64,97,98	0
5	ACT	A	861	4/4	0.91	0.14	53,56,58,59	0
5	ACT	A	860	4/4	0.94	0.14	54,60,62,64	0




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
5	ACT	В	860	4/4	0.95	0.41	66,69,69,78	0
3	H4B	A	600	17/17	0.96	0.17	43,47,52,55	0
3	H4B	В	600	17/17	0.98	0.14	39,46,51,51	0
2	HEM	В	500	43/43	0.98	0.17	32,39,58,59	0
2	HEM	A	500	43/43	0.98	0.23	20,31,64,69	0
7	ZN	A	900	1/1	1.00	0.14	54,54,54,54	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

