wwPDB X-ray Structure Validation Summary Report (i)

Jun 24, 2024 – 02:26 PM EDT

PDB ID	:	6ADH
Title	:	STRUCTURE OF TRICLINIC TERNARY COMPLEX OF HORSE LIVER ALCOHOL DEHYDROGENASE AT 2.9 ANGSTROMS RESOLUTION
Authors	:	Eklund, H.
Deposited on	:	1984-01-16
Resolution	:	2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	374	24% 44% 27% 5%
1	B	374	24% 44% 26% 6%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5669 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HOLO-LIVER ALCOHOL DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	374	2784	1769	472	520	23	154	0	0
1	B	374	2785	1769	472	521	23	151	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	2	2	2	0	0
2	B	2	2	2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	44	21	7	14	2	0	0
			Total	C	N	O	P		
3	B	1	44	21	7	14	2	0	0

- Molecule 4 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
4	A	1	4	2	1	1	0	0
			Total	C	O	S		
4	B	1	4	2	1	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: HOLO-LIVER ALCOHOL DEHYDROGENASE

Chain A: 24% 44% 27% 5%



• Molecule 1: HOLO-LIVER ALCOHOL DEHYDROGENASE

Chain B: 24% 44% 26% 6%




S258
N259
V262
D263
F264
S265
F266
E267
V268
I269
G270
R271
L272
D273
T274
M275
A278
L279
S280
C281
C282
Q283
E284
A285
Y286
G287
V288
S289
V290
I291
V292
G293
V294
P295
P296
D297
L301
S302
M303
N304
P305
M306
L307
L308
L309
S310
G311
R312
T313
W314
K315
I318
F319
F322
K323
S324
K325
D326
S327
V328
P329
K330
L331
V332
A333
D334
F335
M336
K339
F340
A341
L342
D343
P344
L345
I346
T347
H348
V349
L350
P351
F352
E353
K354
I355
N356
E357
L361
L362
R363
S364
G365
E366
S367
I368
R369
T370
I371
L372
T373
F374

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	52.00Å 44.60Å 94.40Å 104.40° 101.90° 70.70°	Depositor
Resolution (Å)	(Not available) – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5669	wwPDB-VP
Average B, all atoms (Å^2)	16.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, DMS, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.03	1/2836 (0.0%)	1.51	43/3834 (1.1%)
1	B	1.15	6/2837 (0.2%)	1.65	54/3834 (1.4%)
All	All	1.09	7/5673 (0.1%)	1.58	97/7668 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	0

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	9	CYS	CA-CB	8.44	1.72	1.53
1	B	282	CYS	CA-CB	7.60	1.70	1.53
1	A	240	CYS	CB-SG	7.06	1.94	1.82
1	B	170	CYS	CB-SG	7.06	1.94	1.82
1	B	9	CYS	CB-SG	6.70	1.93	1.82

The worst 5 of 97 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	46	CYS	N-CA-CB	23.70	153.26	110.60
1	B	122	THR	O-C-N	11.07	140.41	122.70
1	A	195	CYS	CA-CB-SG	9.37	130.87	114.00
1	A	1	SER	O-C-N	9.19	137.40	122.70
1	B	73	VAL	O-C-N	9.15	137.34	122.70

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	46	CYS	CA

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2784	0	2848	324	55
1	B	2785	0	2849	358	59
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	44	0	26	0	0
3	B	44	0	25	4	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
All	All	5669	0	5760	655	63

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 60.

The worst 5 of 655 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:CYS:HB2	1:A:137:ILE:CD1	1.61	1.29
1:A:255:THR:O	1:A:259:ASN:HA	1.42	1.18
1:B:24:GLU:HG2	1:B:132:CYS:SG	1.86	1.15
1:A:23:ILE:CD1	1:A:353:GLU:HA	1.77	1.13
1:A:132:CYS:HB2	1:A:137:ILE:HD11	1.22	1.12

The worst 5 of 63 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:LYS:NZ	1:B:245:ASP:OD1[1_655]	0.31	1.89
1:B:16:GLU:OE2	1:B:81:THR:CG2[1_545]	0.68	1.52
1:A:245:ASP:O	1:B:248:LYS:CB[1_655]	0.71	1.49
1:A:247:LYS:N	1:B:247:LYS:N[1_655]	0.72	1.48
1:A:248:LYS:CD	1:B:245:ASP:CB[1_655]	0.75	1.45

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	372/374 (100%)	281 (76%)	53 (14%)	38 (10%)	0 1
1	B	372/374 (100%)	282 (76%)	58 (16%)	32 (9%)	1 2
All	All	744/748 (100%)	563 (76%)	111 (15%)	70 (9%)	0 1

5 of 70 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	ALA
1	A	33	ALA
1	A	65	ALA
1	A	99	LYS
1	A	109	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/308 (100%)	196 (64%)	112 (36%)	0	0
1	B	308/308 (100%)	195 (63%)	113 (37%)	0	0
All	All	616/616 (100%)	391 (64%)	225 (36%)	0	0

5 of 225 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	19	LYS
1	B	369	ARG
1	B	112	LEU
1	B	357	GLU
1	B	306	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	109	ASN
1	B	124	GLN
1	B	300	ASN
1	B	225	ASN
1	B	242	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	DMS	B	378	2	3,3,3	0.60	0	3,3,3	0.17	0
4	DMS	A	378	2	3,3,3	0.56	0	3,3,3	0.15	0
3	NAD	B	377	-	42,48,48	1.65	4 (9%)	50,73,73	1.92	7 (14%)
3	NAD	A	377	-	42,48,48	1.18	3 (7%)	50,73,73	1.71	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	B	377	-	-	9/26/62/62	0/5/5/5
3	NAD	A	377	-	-	9/26/62/62	0/5/5/5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	377	NAD	C5B-C4B	-7.72	1.27	1.51
3	A	377	NAD	C3N-C7N	4.51	1.57	1.50
3	B	377	NAD	C3N-C7N	3.87	1.56	1.50
3	A	377	NAD	C2B-C1B	-2.37	1.50	1.53
3	B	377	NAD	C6N-N1N	2.29	1.41	1.35

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	377	NAD	C5N-C4N-C3N	-6.61	112.52	120.34
3	A	377	NAD	C5N-C4N-C3N	-6.18	113.03	120.34
3	B	377	NAD	C6N-C5N-C4N	5.84	127.92	119.44
3	A	377	NAD	C6N-C5N-C4N	5.71	127.74	119.44
3	B	377	NAD	O4B-C4B-C5B	5.04	125.96	109.37

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	377	NAD	C5B-O5B-PA-O1A
3	A	377	NAD	O4D-C1D-N1N-C2N
3	A	377	NAD	O4D-C1D-N1N-C6N
3	A	377	NAD	C2D-C1D-N1N-C2N
3	A	377	NAD	C2D-C1D-N1N-C6N

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	377	NAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NAD B 377



Bond lengths



Bond angles



Torsions



Rings

Ligand NAD A 377



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.