

wwPDB NMR Structure Validation Summary Report (i)

May 28, 2020 – 07:13 pm BST

PDB ID	:	1AFP
Title	:	SOLUTION STRUCTURE OF THE ANTIFUNGAL PROTEIN FROM AS-
		PERGILLUS GIGANTEUS. EVIDENCE FOR DISULPHIDE CONFIGURA-
		TIONAL ISOMERISM
Authors	:	Campos-Olivas, R.; Bruix, M.; Santoro, J.; Lacadena, J.; Del Pozo, A.M.;
		Gavilanes, J.G.; Rico, M.
Deposited on	:	1994-11-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


Cyrange	:	Kirchner and Güntert (2011)
$\operatorname{NmrClust}$:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	$v_1n_11_5_13_A$ (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
${ m ShiftChecker}$:	2.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	${f NMR} { m archive} \ (\#{ m Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length		Quality of chain		
1	А	51	8%	71%	12%	10%



2 Ensemble composition and analysis (i)

This entry contains 40 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

	Well-defined (core) protein residues									
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model							
1	A:1-A:18, A:24-A:51 (46)	0.38	1							

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 787 atoms, of which 387 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called ANTIFUNGAL PROTEIN FROM ASPERGILLUS GIGAN-TEUS.

Mol	Chain	Residues		Atoms								
1	Λ	51	Total	С	Η	Ν	Ο	S	0			
	A	51	787	251	387	69	72	8	0			



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: ANTIFUNGAL PROTEIN FROM ASPERGILLUS GIGANTEUS

Chain A:	8%		71%		12%	10%
A1 Y3 N4 G5 C7	Y8 K49 D111 113 C14	K15 Y16 K17 K17 A18 Q19 G21 K22 K22 K22 K22 K22 K22 K22 K22 K22 K	C26 C26 C28 C28 V30 K31 K31	C33 P34 P34 P34 D36 D36 C33 C33 C33 C33 C40 C40 C40 C40 C40 C40 C40 C40 C40 C40	140 K46 K48 K48 C49 C49 C51	

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: ANTIFUNGAL PROTEIN FROM ASPERGILLUS GIGANTEUS

Chain	А	: •			204	%			•							4	13%	6												25	%		·	10	%
A1 T2 N 3 N4	Y8	K9 K10	D11	N12 113	C14	K15 Y16	K17	A18	S20	G21	K22 T23	A24	125	C26	K27 r.28	020	U30	K31	K32	C33	P34	R35	D36	C40	E41	F 42	D43	# 2	K48	C49 VEO	C51				



5 Refinement protocol and experimental data overview (i)

Of the ? calculated structures, 40 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DIANA	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	365	352	367	41±7
All	All	14600	14080	14759	1636

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 56.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Moo	dels
Atom-1	Atom-2	1-2 Clash(A) Distance(A)		Worst	Total
1:A:40:CYS:SG	1:A:51:CYS:HA	1.24	1.70	9	2
1:A:28:CYS:SG	1:A:51:CYS:SG	1.24	2.35	24	20
1:A:14:CYS:HB2	1:A:51:CYS:SG	1.23	1.72	36	3
1:A:40:CYS:SG	1:A:51:CYS:SG	1.22	2.36	36	1
1:A:49:CYS:SG	1:A:51:CYS:SG	1.09	2.49	36	3

5 of 420 unique clashes are listed below, sorted by their clash magnitude.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	44/51~(86%)	$30\pm3~(69\pm7\%)$	$9\pm2~(21\pm6\%)$	$5\pm2~(10\pm4\%)$	1 9
All	All	1760/2040~(86%)	1214~(69%)	366~(21%)	180 (10%)	1 9

5 of 17 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	30	VAL	29
1	А	32	LYS	29
1	А	24	ALA	23
1	А	27	LYS	15
1	А	25	ILE	14

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	39/43~(91%)	$26\pm2(67\pm6\%)$	13 ± 2 (33±6%)	1 12
All	All	1560/1720~(91%)	1044~(67%)	516 (33%)	1 12

5 of 33 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	49	CYS	32
1	А	36	ASP	30
1	А	45	TYR	30
1	А	2	THR	27
1	А	10	LYS	23

6.3.3 RNA (i)

There are no RNA molecules in this entry.



6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

No chemical shift data were provided

