



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2023 – 02:28 AM EDT

PDB ID : 3AF1  
Title : Pantothenate kinase from Mycobacterium tuberculosis (MtPanK) in complex with GDP  
Authors : Chetnani, B.; Kumar, P.; Surolia, A.; Vijayan, M.  
Deposited on : 2010-02-19  
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div> <p style="margin-top: 5px;">5%                      81%                      15%                      ..</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	A	314	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
5	GOL	A	328	-	-	-	X






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 13 6 7	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total Cl 3 3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	208	Total O 208 208	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pantothenate kinase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.90Å 103.90Å 90.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.06 – 2.50 34.06 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (34.06-2.50) 100.0 (34.06-2.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.193 , 0.243 0.188 , 0.236	Depositor DCC
$R_{free}$ test set	958 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.2	Xtrriage
Anisotropy	0.446	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2798	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, GOL, GDP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/2528	0.60	0/3441

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2468	0	2456	39	0
2	A	28	0	12	1	0
3	A	13	0	5	4	0
4	A	3	0	0	1	0
5	A	78	0	104	6	0
6	A	208	0	0	5	0
All	All	2798	0	2577	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:SER:HB2	2:A:313:GDP:O2B	1.56	1.04
1:A:208:THR:HG22	5:A:318:GOL:H32	1.39	1.03
1:A:179:HIS:HB2	3:A:314:FLC:HG2	1.61	0.82
1:A:208:THR:HG22	5:A:318:GOL:C3	2.11	0.81
1:A:208:THR:CG2	5:A:318:GOL:H32	2.17	0.73
1:A:81:PRO:O	1:A:82:GLN:HB3	1.88	0.72
1:A:146:ARG:HH22	5:A:328:GOL:H32	1.56	0.71
3:A:314:FLC:CBC	3:A:314:FLC:OG2	2.38	0.71
1:A:240:LEU:O	1:A:243:ARG:HD3	1.92	0.69
1:A:146:ARG:NH2	5:A:328:GOL:H32	2.15	0.61
1:A:179:HIS:ND1	3:A:314:FLC:HA1	2.16	0.61
1:A:240:LEU:O	1:A:243:ARG:CD	2.47	0.61
1:A:80:GLU:HA	6:A:356:HOH:O	2.02	0.60
1:A:104:SER:HB3	6:A:334:HOH:O	2.02	0.59
1:A:180:LEU:N	3:A:314:FLC:OG1	2.31	0.59
1:A:222:LEU:HD23	1:A:294:LEU:HB3	1.83	0.59
1:A:204:ASN:HB2	4:A:316:CL:CL	2.40	0.58
1:A:67:ARG:HD3	1:A:219:ASP:HB3	1.87	0.57
1:A:81:PRO:HD2	6:A:356:HOH:O	2.04	0.57
1:A:261:SER:H	1:A:264:GLN:NE2	2.03	0.56
1:A:240:LEU:HA	1:A:243:ARG:HD2	1.87	0.56
1:A:302:HIS:HE1	6:A:374:HOH:O	1.91	0.53
1:A:35:VAL:HG22	1:A:38:ARG:NH2	2.24	0.53
1:A:81:PRO:CD	6:A:356:HOH:O	2.57	0.53
1:A:15:ARG:NH2	1:A:48:GLU:HA	2.24	0.52
1:A:81:PRO:O	1:A:82:GLN:CB	2.56	0.51
1:A:27:LEU:HD22	1:A:50:GLU:HG3	1.92	0.51
1:A:150:PRO:HB3	1:A:207:GLN:HE22	1.78	0.49
1:A:223:TYR:CD2	1:A:289:ARG:HG3	2.48	0.48
1:A:129:ASP:HB3	1:A:177:TYR:HB3	1.98	0.46
1:A:278:ARG:HB3	1:A:279:PRO:HD3	1.99	0.43
1:A:122:ARG:HD2	1:A:195:PRO:HA	2.01	0.43
1:A:301:ASP:OD1	1:A:303:SER:HB3	2.18	0.43
1:A:212:LEU:HD12	1:A:212:LEU:HA	1.92	0.42
1:A:43:GLN:HB3	5:A:319:GOL:H31	2.02	0.42
1:A:84:ASN:HA	1:A:85:PRO:HD3	1.79	0.42
1:A:192:VAL:HG13	1:A:195:PRO:HG3	2.02	0.41
1:A:119:HIS:ND1	1:A:119:HIS:N	2.66	0.41
1:A:54:LEU:H	1:A:55:PRO:HD2	1.87	0.40
1:A:296:LEU:HD23	1:A:307:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	305/312 (98%)	285 (93%)	18 (6%)	2 (1%)	22 39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	81	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	263/270 (97%)	253 (96%)	10 (4%)	33 58

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLU
1	A	24	SER
1	A	43	GLN
1	A	68	GLN
1	A	70	LEU
1	A	104	SER
1	A	114	LEU
1	A	119	HIS
1	A	139	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	237	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	GLN
1	A	207	GLN
1	A	253	HIS
1	A	264	GLN
1	A	305	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 3 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	318	-	5,5,5	0.32	0	5,5,5	0.47	0
2	GDP	A	313	-	24,30,30	1.47	4 (16%)	30,47,47	1.60	7 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	329	-	5,5,5	0.35	0	5,5,5	0.48	0
5	GOL	A	320	-	5,5,5	0.35	0	5,5,5	0.39	0
5	GOL	A	328	-	5,5,5	0.36	0	5,5,5	0.36	0
5	GOL	A	326	-	5,5,5	0.36	0	5,5,5	0.20	0
5	GOL	A	319	-	5,5,5	0.41	0	5,5,5	0.25	0
5	GOL	A	327	-	5,5,5	0.35	0	5,5,5	0.23	0
5	GOL	A	322	-	5,5,5	0.31	0	5,5,5	0.45	0
5	GOL	A	323	-	5,5,5	0.38	0	5,5,5	0.25	0
3	FLC	A	314	-	12,12,12	1.05	0	17,17,17	1.52	4 (23%)
5	GOL	A	321	-	5,5,5	0.42	0	5,5,5	0.23	0
5	GOL	A	330	-	5,5,5	0.36	0	5,5,5	0.42	0
5	GOL	A	324	-	5,5,5	0.41	0	5,5,5	0.44	0
5	GOL	A	325	-	5,5,5	0.37	0	5,5,5	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	318	-	-	4/4/4/4	-
2	GDP	A	313	-	-	3/12/32/32	0/3/3/3
5	GOL	A	329	-	-	1/4/4/4	-
5	GOL	A	320	-	-	2/4/4/4	-
5	GOL	A	328	-	-	4/4/4/4	-
5	GOL	A	326	-	-	4/4/4/4	-
5	GOL	A	319	-	-	2/4/4/4	-
5	GOL	A	327	-	-	2/4/4/4	-
5	GOL	A	322	-	-	0/4/4/4	-
5	GOL	A	323	-	-	0/4/4/4	-
3	FLC	A	314	-	-	9/16/16/16	-
5	GOL	A	321	-	-	0/4/4/4	-
5	GOL	A	330	-	-	2/4/4/4	-
5	GOL	A	324	-	-	1/4/4/4	-
5	GOL	A	325	-	-	4/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	313	GDP	C5-C6	-4.02	1.39	1.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	313	GDP	PB-O1B	3.22	1.60	1.50
2	A	313	GDP	O4'-C1'	2.32	1.44	1.41
2	A	313	GDP	C6-N1	2.04	1.40	1.37

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	314	FLC	OB1-CBC-CB	-3.16	117.77	122.25
2	A	313	GDP	O3B-PB-O3A	3.12	115.11	104.64
2	A	313	GDP	C2-N1-C6	-3.12	119.36	125.10
2	A	313	GDP	C3'-C2'-C1'	2.77	105.15	100.98
2	A	313	GDP	C5-C6-N1	2.64	118.62	113.95
2	A	313	GDP	C8-N7-C5	2.64	108.02	102.99
3	A	314	FLC	OG1-CGC-CG	-2.64	115.23	122.94
2	A	313	GDP	O6-C6-C5	-2.38	119.73	124.37
3	A	314	FLC	OB2-CBC-CB	2.15	116.79	113.05
3	A	314	FLC	OA1-CAC-CA	-2.07	116.90	122.94
2	A	313	GDP	C5'-C4'-C3'	-2.03	107.58	115.18

There are no chirality outliers.

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	314	FLC	CA-CB-CBC-OB1
3	A	314	FLC	CA-CB-CBC-OB2
3	A	314	FLC	OHB-CB-CBC-OB1
3	A	314	FLC	OHB-CB-CBC-OB2
5	A	320	GOL	O1-C1-C2-O2
5	A	320	GOL	O1-C1-C2-C3
5	A	325	GOL	O1-C1-C2-C3
5	A	325	GOL	C1-C2-C3-O3
5	A	327	GOL	O1-C1-C2-O2
5	A	327	GOL	O1-C1-C2-C3
5	A	328	GOL	O1-C1-C2-C3
3	A	314	FLC	CA-CB-CG-CGC
3	A	314	FLC	OHB-CB-CG-CGC
5	A	325	GOL	O1-C1-C2-O2
5	A	318	GOL	O1-C1-C2-C3
5	A	318	GOL	C1-C2-C3-O3
5	A	319	GOL	C1-C2-C3-O3
5	A	326	GOL	O1-C1-C2-C3
5	A	326	GOL	C1-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
5	A	328	GOL	C1-C2-C3-O3
5	A	330	GOL	C1-C2-C3-O3
5	A	318	GOL	O1-C1-C2-O2
5	A	319	GOL	O2-C2-C3-O3
5	A	325	GOL	O2-C2-C3-O3
5	A	328	GOL	O1-C1-C2-O2
5	A	330	GOL	O2-C2-C3-O3
5	A	318	GOL	O2-C2-C3-O3
5	A	326	GOL	O2-C2-C3-O3
3	A	314	FLC	CBC-CB-CG-CGC
2	A	313	GDP	PA-O3A-PB-O1B
5	A	326	GOL	O1-C1-C2-O2
5	A	328	GOL	O2-C2-C3-O3
5	A	324	GOL	C1-C2-C3-O3
2	A	313	GDP	PA-O3A-PB-O3B
3	A	314	FLC	CB-CG-CGC-OG1
3	A	314	FLC	CB-CG-CGC-OG2
5	A	329	GOL	C1-C2-C3-O3
2	A	313	GDP	C3'-C4'-C5'-O5'

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	318	GOL	3	0
2	A	313	GDP	1	0
5	A	328	GOL	2	0
5	A	319	GOL	1	0
3	A	314	FLC	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/312 (98%)	0.15	17 (5%) <span style="border: 1px solid red; padding: 2px;">25</span> <span style="border: 1px solid red; padding: 2px;">26</span>	27, 42, 74, 103	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	86	ASP	4.9
1	A	85	PRO	4.8
1	A	82	GLN	3.9
1	A	81	PRO	3.7
1	A	171	TYR	3.4
1	A	117	TRP	3.1
1	A	24	SER	3.1
1	A	120	HIS	3.0
1	A	84	ASN	3.0
1	A	83	GLN	2.9
1	A	119	HIS	2.8
1	A	94	GLY	2.7
1	A	222	LEU	2.6
1	A	95	VAL	2.3
1	A	224	VAL	2.2
1	A	101	VAL	2.1
1	A	118	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	320	6/6	0.69	0.33	82,85,89,90	0
5	GOL	A	324	6/6	0.73	0.29	70,79,84,85	0
5	GOL	A	327	6/6	0.77	0.26	85,89,90,90	0
5	GOL	A	328	6/6	0.78	0.44	76,84,87,87	0
5	GOL	A	321	6/6	0.83	0.20	71,74,76,77	0
4	CL	A	317	1/1	0.83	0.30	96,96,96,96	0
5	GOL	A	326	6/6	0.84	0.28	81,89,91,91	0
5	GOL	A	329	6/6	0.84	0.33	84,88,89,90	0
3	FLC	A	314	13/13	0.85	0.20	67,85,87,88	0
5	GOL	A	325	6/6	0.85	0.20	94,96,97,98	0
5	GOL	A	323	6/6	0.85	0.49	66,81,83,85	0
5	GOL	A	330	6/6	0.85	0.53	65,75,82,84	0
5	GOL	A	319	6/6	0.88	0.23	97,98,98,99	0
5	GOL	A	318	6/6	0.91	0.33	71,73,76,78	0
5	GOL	A	322	6/6	0.93	0.57	52,66,72,79	0
4	CL	A	315	1/1	0.95	0.08	82,82,82,82	0
2	GDP	A	313	28/28	0.97	0.17	30,36,48,62	0
4	CL	A	316	1/1	0.98	0.31	58,58,58,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.