



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 09:47 am BST

PDB ID : 5AFO  
Title : Long Polar Fimbriae adhesin LpfD from the adherent invasive E. coli strain LF82  
Authors : Coppens, F.; Iyyathurai, J.; Remaut, H.  
Deposited on : 2015-01-23  
Resolution : 1.82 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	 86% 7% • 5%
1	B	362	 87% 7% 5%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FIMBRIAE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	343	2665	1682	440	535	8	0	10	0
1	B	343	2673	1686	444	535	8	0	9	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	327	GLY	-	expression tag	UNP E2QGGQ5
A	328	GLY	-	expression tag	UNP E2QGGQ5
A	329	GLY	-	expression tag	UNP E2QGGQ5
A	330	GLY	-	expression tag	UNP E2QGGQ5
A	331	GLY	-	expression tag	UNP E2QGGQ5
A	332	GLY	-	expression tag	UNP E2QGGQ5
A	333	GLY	-	expression tag	UNP E2QGGQ5
A	334	GLY	-	expression tag	UNP E2QGGQ5
A	335	GLY	-	expression tag	UNP E2QGGQ5
A	336	GLY	-	expression tag	UNP E2QGGQ5
A	337	ALA	-	expression tag	UNP E2QGGQ5
A	338	THR	-	expression tag	UNP E2QGGQ5
A	339	THR	-	expression tag	UNP E2QGGQ5
A	340	ASP	-	expression tag	UNP E2QGGQ5
A	341	LEU	-	expression tag	UNP E2QGGQ5
A	342	GLY	-	expression tag	UNP E2QGGQ5
A	343	ALA	-	expression tag	UNP E2QGGQ5
A	344	LYS	-	expression tag	UNP E2QGGQ5
A	345	GLY	-	expression tag	UNP E2QGGQ5
A	346	THR	-	expression tag	UNP E2QGGQ5
A	347	LEU	-	expression tag	UNP E2QGGQ5
A	348	LYS	-	expression tag	UNP E2QGGQ5
A	349	PHE	-	expression tag	UNP E2QGGQ5
A	350	SER	-	expression tag	UNP E2QGGQ5
A	351	LEU	-	expression tag	UNP E2QGGQ5

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	352	LYS	-	expression tag	UNP E2QGGQ5
A	353	ILE	-	expression tag	UNP E2QGGQ5
A	354	SER	-	expression tag	UNP E2QGGQ5
A	355	GLN	-	expression tag	UNP E2QGGQ5
A	356	GLY	-	expression tag	UNP E2QGGQ5
A	357	HIS	-	expression tag	UNP E2QGGQ5
A	358	HIS	-	expression tag	UNP E2QGGQ5
A	359	HIS	-	expression tag	UNP E2QGGQ5
A	360	HIS	-	expression tag	UNP E2QGGQ5
A	361	HIS	-	expression tag	UNP E2QGGQ5
A	362	HIS	-	expression tag	UNP E2QGGQ5
B	327	GLY	-	expression tag	UNP E2QGGQ5
B	328	GLY	-	expression tag	UNP E2QGGQ5
B	329	GLY	-	expression tag	UNP E2QGGQ5
B	330	GLY	-	expression tag	UNP E2QGGQ5
B	331	GLY	-	expression tag	UNP E2QGGQ5
B	332	GLY	-	expression tag	UNP E2QGGQ5
B	333	GLY	-	expression tag	UNP E2QGGQ5
B	334	GLY	-	expression tag	UNP E2QGGQ5
B	335	GLY	-	expression tag	UNP E2QGGQ5
B	336	GLY	-	expression tag	UNP E2QGGQ5
B	337	ALA	-	expression tag	UNP E2QGGQ5
B	338	THR	-	expression tag	UNP E2QGGQ5
B	339	THR	-	expression tag	UNP E2QGGQ5
B	340	ASP	-	expression tag	UNP E2QGGQ5
B	341	LEU	-	expression tag	UNP E2QGGQ5
B	342	GLY	-	expression tag	UNP E2QGGQ5
B	343	ALA	-	expression tag	UNP E2QGGQ5
B	344	LYS	-	expression tag	UNP E2QGGQ5
B	345	GLY	-	expression tag	UNP E2QGGQ5
B	346	THR	-	expression tag	UNP E2QGGQ5
B	347	LEU	-	expression tag	UNP E2QGGQ5
B	348	LYS	-	expression tag	UNP E2QGGQ5
B	349	PHE	-	expression tag	UNP E2QGGQ5
B	350	SER	-	expression tag	UNP E2QGGQ5
B	351	LEU	-	expression tag	UNP E2QGGQ5
B	352	LYS	-	expression tag	UNP E2QGGQ5
B	353	ILE	-	expression tag	UNP E2QGGQ5
B	354	SER	-	expression tag	UNP E2QGGQ5
B	355	GLN	-	expression tag	UNP E2QGGQ5
B	356	GLY	-	expression tag	UNP E2QGGQ5
B	357	HIS	-	expression tag	UNP E2QGGQ5

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	358	HIS	-	expression tag	UNP E2QGG5
B	359	HIS	-	expression tag	UNP E2QGG5
B	360	HIS	-	expression tag	UNP E2QGG5
B	361	HIS	-	expression tag	UNP E2QGG5
B	362	HIS	-	expression tag	UNP E2QGG5

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	399	Total	O	0	0
			399	399		
3	B	259	Total	O	0	0
			259	259		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FIMBRIAE



- Molecule 1: FIMBRIAE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.01Å 372.13Å 99.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.59 – 1.82 48.59 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.59-1.82) 99.9 (48.59-1.82)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 1.82Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.167 , 0.196 0.176 , 0.203	Depositor DCC
$R_{free}$ test set	4108 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtrriage
Anisotropy	0.557	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6008	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.00	4/2732 (0.1%)	0.97	5/3726 (0.1%)
1	B	0.95	0/2731	0.95	4/3722 (0.1%)
All	All	0.98	4/5463 (0.1%)	0.96	9/7448 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	132	GLN	CD-OE1	-5.98	1.10	1.24
1	A	41	GLN	CB-CG	-5.65	1.37	1.52
1	A	112	SER	CB-OG	-5.10	1.35	1.42
1	A	191	SER	CB-OG	5.05	1.48	1.42

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	292	ARG	NE-CZ-NH2	-7.54	116.53	120.30
1	A	63	ARG	NE-CZ-NH1	7.03	123.82	120.30
1	A	340	ASP	CB-CG-OD1	6.82	124.43	118.30
1	B	292	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	B	195	ASP	CB-CG-OD1	6.32	123.99	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2665	0	2609	16	0
1	B	2673	0	2611	18	0
2	A	6	0	8	0	0
2	B	6	0	8	2	0
3	A	399	0	0	6	0
3	B	259	0	0	4	1
All	All	6008	0	5236	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:GLN:OE1	3:B:2181:HOH:O	2.01	0.79
1:A:92[B]:THR:HG23	3:A:2184:HOH:O	1.90	0.71
1:A:130:LYS:HD3	3:A:2112:HOH:O	1.91	0.70
1:A:213:GLN:O	1:A:214:ASN:ND2	2.28	0.66
1:A:326:ASP:OD1	1:A:326:ASP:C	2.34	0.65

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:2016:HOH:O	3:B:2016:HOH:O[3_556]	1.72	0.48

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/362 (96%)	343 (98%)	6 (2%)	0	100	100
1	B	348/362 (96%)	343 (99%)	5 (1%)	0	100	100
All	All	697/724 (96%)	686 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/298 (101%)	297 (99%)	4 (1%)	69	61
1	B	300/298 (101%)	287 (96%)	13 (4%)	29	14
All	All	601/596 (101%)	584 (97%)	17 (3%)	46	29

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	84[B]	ILE
1	B	139	LYS
1	B	263	GLU
1	B	84[A]	ILE
1	B	275	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	219	GLN
1	B	276	ASN
1	B	41	GLN
1	A	202	HIS
1	A	276	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	1001	-	5,5,5	0.23	0	5,5,5	0.85	0
2	GOL	B	1002	-	5,5,5	0.54	0	5,5,5	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1001	-	-	4/4/4/4	-
2	GOL	B	1002	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1002	GOL	O1-C1-C2-O2
2	B	1002	GOL	O1-C1-C2-C3
2	A	1001	GOL	O1-C1-C2-O2
2	A	1001	GOL	O2-C2-C3-O3
2	A	1001	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1002	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	343/362 (94%)	-0.20	5 (1%) 73 70	15, 26, 44, 70	0
1	B	343/362 (94%)	0.16	20 (5%) 23 18	15, 31, 66, 88	0
All	All	686/724 (94%)	-0.02	25 (3%) 42 37	15, 28, 59, 88	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	309	LEU	5.3
1	B	208	THR	4.6
1	A	309	LEU	3.8
1	B	205	LYS	3.5
1	B	202	HIS	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	B	1002	6/6	0.85	0.18	38,45,47,53	0
2	GOL	A	1001	6/6	0.91	0.14	42,45,46,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.