



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 22, 2023 – 01:49 PM EDT

PDB ID : 3AGB
Title : F218V mutant of the substrate-free form of red chlorophyll catabolite reductase from *Arabidopsis thaliana*
Authors : Sugishima, M.; Fukuyama, K.
Deposited on : 2010-03-30
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 3% 76% 18% . .
1	B	276	 8% 74% 18% . 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Red chlorophyll catabolite reductase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	271	2143	1354	368	412	9	0	0	0
1	B	263	2056	1300	354	393	9	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLY	-	expression tag	UNP Q8LDU4
A	45	PRO	-	expression tag	UNP Q8LDU4
A	46	LEU	-	expression tag	UNP Q8LDU4
A	47	GLY	-	expression tag	UNP Q8LDU4
A	48	SER	-	expression tag	UNP Q8LDU4
A	218	VAL	PHE	engineered mutation	UNP Q8LDU4
B	44	GLY	-	expression tag	UNP Q8LDU4
B	45	PRO	-	expression tag	UNP Q8LDU4
B	46	LEU	-	expression tag	UNP Q8LDU4
B	47	GLY	-	expression tag	UNP Q8LDU4
B	48	SER	-	expression tag	UNP Q8LDU4
B	218	VAL	PHE	engineered mutation	UNP Q8LDU4

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	108	Total 108	O 108	0	0
3	B	47	Total 47	O 47	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Red chlorophyll catabolite reductase, chloroplastic



- Molecule 1: Red chlorophyll catabolite reductase, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.14Å 84.43Å 132.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.58 – 2.20 35.58 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.5 (35.58-2.20) 98.5 (35.58-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.52 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.221 , 0.292 0.237 , 0.309	Depositor DCC
R_{free} test set	1330 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtrriage
Anisotropy	0.103	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4356	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2185	0.65	0/2960
1	B	0.40	0/2101	0.58	2/2850 (0.1%)
All	All	0.44	0/4286	0.61	2/5810 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	155	LEU	CA-CB-CG	5.21	127.28	115.30
1	B	163	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2143	0	2131	33	0
1	B	2056	0	2038	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	108	0	0	1	0
3	B	47	0	0	0	0
All	All	4356	0	4169	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:157:GLN:HB2	1:B:163:LEU:HD23	1.66	0.78
1:A:294:LEU:O	1:A:298:ARG:NH2	2.26	0.69
1:A:51:LYS:HG2	1:A:52:PHE:N	2.11	0.66
1:A:97:ASN:HD22	1:A:97:ASN:C	2.05	0.60
1:B:268:VAL:HG12	1:B:273:ARG:HG2	1.85	0.58
1:A:51:LYS:HG2	1:A:52:PHE:H	1.69	0.58
1:B:234:LYS:O	1:B:238:ILE:HG12	2.05	0.57
1:A:183:LYS:HA	1:A:187:GLN:HB2	1.86	0.57
1:B:161:LYS:HB3	1:B:229:ALA:O	2.04	0.57
1:B:74:GLU:O	1:B:78:GLN:HG2	2.05	0.56
1:A:244:SER:HB2	1:A:245:PRO:HD3	1.87	0.56
1:B:69:LEU:HB3	1:B:138:ILE:HD12	1.86	0.56
1:B:240:ARG:HD3	1:B:241:ASP:OD1	2.04	0.56
1:B:110:ASP:OD2	1:B:111:LYS:O	2.24	0.55
1:B:78:GLN:HE22	1:B:107:ARG:HH22	1.55	0.55
1:B:183:LYS:HA	1:B:187[A]:GLN:HB2	1.89	0.55
1:A:171:HIS:H	1:A:171:HIS:CD2	2.26	0.53
1:B:175:LEU:HG	1:B:182:LEU:HD13	1.91	0.52
1:B:288:ILE:HD13	1:B:313:LYS:HE2	1.91	0.52
1:A:76:ARG:NH1	1:A:252:GLU:OE2	2.41	0.52
1:A:292:LEU:HD12	1:A:309:VAL:HG22	1.93	0.51
1:B:104:LEU:HD11	1:B:119:ILE:HD11	1.93	0.50
1:A:261:GLU:O	1:A:263:GLU:N	2.44	0.50
1:A:127:ILE:HD11	1:A:133:LEU:HB2	1.94	0.50
1:B:97:ASN:HD22	1:B:99:SER:H	1.61	0.49
1:A:289:GLU:O	1:A:294:LEU:HD13	2.13	0.48
1:B:111:LYS:C	1:B:113:SER:H	2.17	0.48
1:A:111:LYS:O	1:A:112:SER:OG	2.27	0.48
1:B:274:MET:HB3	1:B:278:ARG:HH21	1.78	0.47
1:A:208:VAL:HA	3:A:341:HOH:O	2.15	0.46
1:A:240:ARG:HG2	1:A:241:ASP:OD2	2.15	0.46
1:B:73:VAL:HG11	1:B:118:VAL:HG11	1.98	0.46
1:A:104:LEU:HD11	1:A:312:ILE:HD11	1.98	0.46
1:B:253:VAL:O	1:B:257:ARG:HB3	2.16	0.45
1:A:231:GLU:HB2	1:A:234:LYS:HB3	1.99	0.44
1:A:244:SER:CB	1:A:245:PRO:HD3	2.47	0.44
1:B:163:LEU:HD12	1:B:227:ILE:HD12	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:LEU:HD13	1:A:225:LEU:HD22	1.99	0.43
1:B:55:PHE:HB2	1:B:63:LYS:HG2	1.99	0.43
1:B:233:ASP:OD1	1:B:233:ASP:N	2.51	0.43
1:A:111:LYS:CE	1:A:111:LYS:HA	2.50	0.42
1:A:193:SER:O	1:A:197:SER:HB3	2.20	0.42
1:A:163:LEU:HD12	1:A:235:LEU:HD11	2.02	0.42
1:A:173:LYS:HD2	1:A:181:TYR:CD2	2.55	0.42
1:A:261:GLU:CD	1:A:262:GLU:H	2.23	0.42
1:A:56:PRO:O	1:A:57:TYR:HB2	2.20	0.42
1:A:169:LEU:HD11	1:A:254:TRP:HB2	2.02	0.42
1:A:319:LEU:HD13	1:A:319:LEU:OXT	2.19	0.42
1:A:175:LEU:HG	1:A:182:LEU:HD13	2.01	0.41
1:A:262:GLU:HG3	1:A:263:GLU:H	1.84	0.41
1:B:169:LEU:HD21	1:B:253:VAL:HG22	2.02	0.41
1:B:161:LYS:HB2	1:B:161:LYS:NZ	2.36	0.41
1:B:183:LYS:HE2	1:B:188:ASP:OD2	2.21	0.41
1:A:65:LEU:HD13	1:A:244:SER:HA	2.03	0.41
1:A:89:ASP:OD1	1:A:89:ASP:N	2.53	0.40
1:B:204:VAL:HG22	1:B:227:ILE:HG12	2.02	0.40
1:B:165:LEU:HD23	1:B:225:LEU:HD12	2.02	0.40
1:A:97:ASN:HD22	1:A:99:SER:H	1.70	0.40
1:A:147:LYS:NZ	1:A:185:TYR:OH	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/276 (98%)	255 (95%)	12 (4%)	2 (1%)	22	22
1	B	260/276 (94%)	245 (94%)	12 (5%)	3 (1%)	13	10
All	All	529/552 (96%)	500 (94%)	24 (4%)	5 (1%)	17	16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	262	GLU
1	A	260	LYS
1	B	52	PHE
1	B	231	GLU
1	B	128	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	245/255 (96%)	223 (91%)	22 (9%)	9 9
1	B	235/255 (92%)	213 (91%)	22 (9%)	8 8
All	All	480/510 (94%)	436 (91%)	44 (9%)	9 9

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ARG
1	A	69	LEU
1	A	81	LEU
1	A	82	LEU
1	A	89	ASP
1	A	97	ASN
1	A	107	ARG
1	A	111	LYS
1	A	162	SER
1	A	163	LEU
1	A	169	LEU
1	A	175	LEU
1	A	177	LEU
1	A	198	LEU
1	A	199	LEU
1	A	208	VAL
1	A	231	GLU
1	A	236	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	281	LYS
1	A	292	LEU
1	A	303	GLU
1	A	318	VAL
1	B	51	LYS
1	B	69	LEU
1	B	80	GLN
1	B	81	LEU
1	B	97	ASN
1	B	103	SER
1	B	113	SER
1	B	155	LEU
1	B	169	LEU
1	B	175	LEU
1	B	177	LEU
1	B	180	ASP
1	B	198	LEU
1	B	199	LEU
1	B	203	GLU
1	B	233	ASP
1	B	240	ARG
1	B	249	GLU
1	B	273	ARG
1	B	281	LYS
1	B	294	LEU
1	B	303	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	97	ASN
1	A	171	HIS
1	A	187	GLN
1	A	196	GLN
1	B	78	GLN
1	B	97	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/276 (98%)	0.10	8 (2%) 50 48	9, 17, 29, 36	0
1	B	263/276 (95%)	0.51	22 (8%) 11 9	14, 27, 38, 43	0
All	All	534/552 (96%)	0.30	30 (5%) 24 23	9, 22, 36, 43	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	160	SER	6.9
1	A	319	LEU	4.8
1	B	229	ALA	4.4
1	A	294	LEU	4.3
1	B	271	GLU	4.1
1	A	261	GLU	3.7
1	B	257	ARG	3.4
1	A	265	LYS	3.3
1	B	267	VAL	3.1
1	A	264	GLU	3.1
1	B	159	SER	3.0
1	B	129	THR	2.8
1	B	270	GLU	2.8
1	A	262	GLU	2.8
1	B	230	GLU	2.8
1	B	79	SER	2.7
1	B	231	GLU	2.5
1	B	53	MET	2.4
1	B	228	ASP	2.4
1	A	231	GLU	2.4
1	B	61	THR	2.4
1	B	152	VAL	2.3
1	B	131	VAL	2.3
1	B	233	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	130	GLY	2.2
1	B	128	PRO	2.2
1	B	161	LYS	2.2
1	A	190	ALA	2.2
1	B	269	GLY	2.1
1	B	235	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NA	B	2	1/1	0.67	0.10	41,41,41,41	0
2	NA	A	1	1/1	0.93	0.12	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.