



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 11:38 am BST

PDB ID : 3AGY
Title : Crystal structure of human Hsp40 Hdj1 peptide-binding domain complexed with a C-terminal peptide of Hsp70
Authors : Suzuki, H.; Noguchi, S.; Satow, Y.
Deposited on : 2010-04-12
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 5px;">5% 85% 12% ..</p>
1	B	181	<div style="display: flex; align-items: center;"> <div style="width: 23%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 5px;">23% 86% 11% ..</p>
2	C	8	<div style="display: flex; align-items: center;"> <div style="width: 75%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 5px;">75% 75% 13% 13%</p>
2	D	8	<div style="display: flex; align-items: center;"> <div style="width: 38%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 63%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 5px;">38% 100% 63%</p>
2	F	8	<div style="display: flex; align-items: center;"> <div style="width: 38%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 63%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 25%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 5px;">38% 63% 13% 25%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3220 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DnaJ homolog subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	176	1406	894	250	257	5	0	2	0
1	B	176	1425	905	258	257	5	0	7	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	MET	-	EXPRESSION TAG	UNP P25685
B	160	MET	-	EXPRESSION TAG	UNP P25685

- Molecule 2 is a protein called peptide of Heat shock cognate 71 kDa protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	7	51	32	7	12	0	0	0
2	D	8	60	36	8	16	0	0	0
2	F	6	47	30	6	11	0	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	123	Total 123	O 123	0	0
3	B	104	Total 104	O 104	0	0
3	F	4	Total 4	O 4	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: DnaJ homolog subfamily B member 1



- Molecule 1: DnaJ homolog subfamily B member 1



- Molecule 2: peptide of Heat shock cognate 71 kDa protein




- Molecule 2: peptide of Heat shock cognate 71 kDa protein



- Molecule 2: peptide of Heat shock cognate 71 kDa protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	104.75Å 41.13Å 62.87Å 90.00° 97.42° 90.00°	Depositor
Resolution (Å)	42.71 – 1.85 42.71 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.71-1.85) 98.8 (42.71-1.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.225 , 0.247 0.225 , 0.221	Depositor DCC
R_{free} test set	2282 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtrriage
Anisotropy	0.300	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3220	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	0/1443	0.68	0/1947
1	B	0.73	1/1491 (0.1%)	0.75	1/2006 (0.0%)
2	C	0.58	0/51	0.48	0/69
2	D	0.58	0/60	0.55	0/80
2	F	0.41	0/47	0.52	0/63
All	All	0.69	1/3092 (0.0%)	0.71	1/4165 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	269	CYS	CB-SG	-9.27	1.66	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	263	ARG	NE-CZ-NH1	-8.09	116.25	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1406	0	1470	22	0
1	B	1425	0	1507	11	0
2	C	51	0	48	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	60	0	52	3	0
2	F	47	0	46	1	0
3	A	123	0	0	1	0
3	B	104	0	0	0	0
3	F	4	0	0	0	0
All	All	3220	0	3123	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:ARG:H	1:A:293[A]:MET:HE3	1.00	1.10
1:A:290:ARG:N	1:A:293[A]:MET:HE3	1.78	0.97
1:A:290:ARG:H	1:A:293[A]:MET:CE	1.88	0.85
1:A:185:ILE:HD13	1:A:235:ILE:HD13	1.65	0.79
1:A:185:ILE:HD13	1:A:235:ILE:CD1	2.22	0.70
1:A:217:LYS:HE3	1:A:236:VAL:HG11	1.74	0.69
1:A:290:ARG:CZ	1:A:293[A]:MET:HE2	2.25	0.66
1:A:188:LYS:HB3	1:A:197:ILE:HG21	1.79	0.65
1:A:185:ILE:CD1	1:A:235:ILE:HD13	2.26	0.65
1:B:206:ILE:HG21	1:B:218:ILE:HD13	1.85	0.59
1:A:185:ILE:CD1	1:A:235:ILE:CD1	2.83	0.56
1:A:290:ARG:NH2	1:A:293[A]:MET:HE1	2.24	0.53
1:A:340:ILE:HG23	3:A:370:HOH:O	2.08	0.52
1:B:185:ILE:HG22	2:D:637:ILE:HG12	1.93	0.50
2:D:640:VAL:HG22	2:D:641:ASP:H	1.75	0.50
1:A:290:ARG:CZ	1:A:293[A]:MET:CE	2.88	0.50
1:A:185:ILE:HD12	2:C:637:ILE:HG12	1.93	0.50
1:A:172:LEU:HD21	1:A:249:ARG:HB2	1.94	0.49
1:B:217:LYS:HB2	2:F:638:GLU:HB3	1.95	0.48
1:B:294:ARG:HA	1:B:316:ILE:O	2.14	0.47
1:A:294:ARG:HA	1:A:316:ILE:O	2.14	0.47
1:A:323:PRO:HG2	1:A:326:ILE:HD11	1.96	0.46
1:A:188:LYS:HB3	1:A:197:ILE:CG2	2.45	0.46
1:A:240:LYS:HA	1:A:240:LYS:HD2	1.77	0.46
1:B:224:GLY:HA2	2:D:635:PRO:HG3	2.01	0.43
1:B:273:VAL:HG21	1:B:316:ILE:HD11	2.01	0.42
1:B:296[B]:LYS:HD2	1:B:313:ASP:OD2	2.19	0.42
1:B:314:LEU:HD23	1:B:315:ILE:N	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:LYS:HD2	1:A:231:ILE:HD11	2.01	0.41
1:B:197:ILE:HG12	1:B:227:THR:HG22	2.02	0.41
1:A:222:LYS:HB3	1:A:232:PRO:HG2	2.02	0.41
1:B:314:LEU:C	1:B:314:LEU:HD23	2.41	0.41
1:B:222:LYS:HB3	1:B:232:PRO:HG2	2.03	0.41
1:A:290:ARG:NH2	1:A:293[A]:MET:CE	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/181 (97%)	175 (99%)	1 (1%)	0	100	100
1	B	181/181 (100%)	177 (98%)	3 (2%)	1 (1%)	25	12
2	C	5/8 (62%)	5 (100%)	0	0	100	100
2	D	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
2	F	4/8 (50%)	4 (100%)	0	0	100	100
All	All	372/386 (96%)	366 (98%)	5 (1%)	1 (0%)	41	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	228	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/166 (98%)	159 (98%)	4 (2%)	47	31
1	B	168/166 (101%)	164 (98%)	4 (2%)	49	33
2	C	6/7 (86%)	6 (100%)	0	100	100
2	D	7/7 (100%)	6 (86%)	1 (14%)	3	0
2	F	6/7 (86%)	6 (100%)	0	100	100
All	All	350/353 (99%)	341 (97%)	9 (3%)	50	30

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	165	THR
1	A	187	HIS
1	A	227	THR
1	A	240	LYS
1	B	187	HIS
1	B	227	THR
1	B	245[A]	ASN
1	B	245[B]	ASN
2	D	636	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	176/181 (97%)	0.41	9 (5%) 28 26	13, 29, 52, 64	0
1	B	176/181 (97%)	1.21	41 (23%) 0 0	12, 32, 88, 90	0
2	C	7/8 (87%)	2.67	6 (85%) 0 0	71, 71, 74, 74	0
2	D	8/8 (100%)	4.34	8 (100%) 0 0	102, 102, 103, 103	0
2	F	6/8 (75%)	2.03	3 (50%) 0 0	65, 66, 67, 68	0
All	All	373/386 (96%)	0.94	67 (17%) 1 1	12, 32, 87, 103	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	231	ILE	12.7
1	B	227	THR	8.6
1	B	229	ASN	7.9
1	B	230	ASN	7.4
1	B	197	ILE	7.2
1	B	193	ASP	6.3
1	B	196	SER	6.1
1	B	228	SER	5.9
2	D	640	VAL	5.4
1	A	231	ILE	5.2
2	D	636	THR	5.2
1	B	232	PRO	4.9
1	B	190	LEU	4.9
2	D	635	PRO	4.9
1	A	166	HIS	4.6
2	D	638	GLU	4.6
2	D	641	ASP	4.5
1	B	200	GLU	4.5
1	A	197	ILE	4.5
2	C	634	GLY	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	189	ARG	4.3
1	B	233	ALA	4.2
1	B	203	ILE	4.1
2	D	637	ILE	4.0
1	B	195[A]	LYS	4.0
1	B	166	HIS	4.0
1	B	183	MET	3.8
1	B	204	LEU	3.8
1	A	229	ASN	3.7
1	B	199	ASN	3.6
1	B	236	VAL	3.6
1	B	222	LYS	3.5
1	B	185	ILE	3.3
1	B	194	GLY	3.3
1	B	202	LYS	3.3
1	B	226	GLN	3.2
1	B	184	LYS	3.2
1	B	192	PRO	3.1
1	B	201	ASP	3.1
2	D	634	GLY	3.1
2	F	637	ILE	3.1
2	D	639	GLU	3.0
2	F	636	THR	3.0
1	B	221	PRO	3.0
1	B	187	HIS	3.0
1	A	230	ASN	2.9
1	B	220	PHE	2.9
1	B	191	ASN	2.8
1	A	227	THR	2.8
2	C	636	THR	2.8
1	B	186	SER	2.7
1	B	235	ILE	2.7
2	C	635	PRO	2.7
1	B	206	ILE	2.7
2	C	638	GLU	2.6
1	B	224	GLY	2.5
1	B	188	LYS	2.4
1	B	218	ILE	2.4
2	C	640	VAL	2.3
2	C	637	ILE	2.3
1	B	219	THR	2.3
1	B	234	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	198	ARG	2.2
1	A	235	ILE	2.2
2	F	638	GLU	2.1
1	A	195	LYS	2.0
1	A	165	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.