



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 06:27 am BST

PDB ID : 5AIT
Title : A complex of of RNF4-RING domain, UbeV2, Ubc13-Ub (isopeptide crosslink)
Authors : Branigan, E.; Naismith, J.H.
Deposited on : 2015-02-17
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	 2% 74% 22% • •
2	B	154	 68% 27% • 5%
2	E	154	 66% 28% • 5%
3	C	76	 75% 24% •
3	F	76	 64% 34% •
4	D	147	 70% 24% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	G	147	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a green segment on the left labeled '71%', a yellow segment in the middle labeled '22%', and a red segment on the right labeled '6%'. A small red square is positioned at the start of the bar, and a small black dot is at the end of the red segment. A '%' symbol is located above the start of the bar.</p>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6742 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 UBIQUITIN-PROTEIN LIGASE RNF4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	999	614	185	184	16	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	GLY	-	expression tag	UNP O88846
A	128	ALA	-	expression tag	UNP O88846
A	129	MET	-	expression tag	UNP O88846
A	130	GLY	-	expression tag	UNP O88846
A	195	GLY	-	linker	UNP O88846

- Molecule 2 is a protein called UBIQUITIN-CONJUGATING ENZYME E2 N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	147	1173	755	202	213	3	0	0	0
2	E	147	1173	755	202	213	3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP P61088
B	0	ALA	-	expression tag	UNP P61088
B	87	LYS	CYS	engineered mutation	UNP P61088
B	92	ALA	LYS	engineered mutation	UNP P61088
E	-1	GLY	-	expression tag	UNP P61088
E	0	ALA	-	expression tag	UNP P61088
E	87	LYS	CYS	engineered mutation	UNP P61088
E	92	ALA	LYS	engineered mutation	UNP P61088

- Molecule 3 is a protein called POLYUBIQUITIN-C.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	75	Total	C	N	O	0	0	0
			593	373	104	116			
3	F	75	Total	C	N	O	0	0	0
			593	373	104	116			

- Molecule 4 is a protein called UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	140	Total	C	N	O	S	0	0	0
			1112	697	195	212	8			
4	G	138	Total	C	N	O	S	0	0	0
			1095	689	191	207	8			

There are 4 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	GLY	-	expression tag	UNP Q15819
D	0	ALA	-	expression tag	UNP Q15819
G	-1	GLY	-	expression tag	UNP Q15819
G	0	ALA	-	expression tag	UNP Q15819

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	Zn	0	0
			4	4		



- Molecule 4: UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 2



- Molecule 4: UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	77.58Å 77.58Å 328.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	67.19 – 3.40 65.83 – 3.40	Depositor EDS
% Data completeness (in resolution range)	89.0 (67.19-3.40) 89.0 (65.83-3.40)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 3.41Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.208 , 0.286 0.215 , 0.281	Depositor DCC
R_{free} test set	753 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	139.6	Xtrriage
Anisotropy	0.118	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 92.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6742	wwPDB-VP
Average B, all atoms (Å ²)	140.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.66	0/1018	0.82	0/1372
2	B	0.69	0/1203	0.79	0/1639
2	E	0.64	0/1203	0.76	0/1639
3	C	0.64	0/599	0.80	0/806
3	F	0.63	0/599	0.75	0/806
4	D	0.71	0/1135	0.81	0/1534
4	G	0.75	1/1118 (0.1%)	0.87	0/1512
All	All	0.68	1/6875 (0.0%)	0.80	0/9308

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	46	TRP	CB-CG	-6.30	1.39	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	999	0	980	24	0
2	B	1173	0	1185	51	0
2	E	1173	0	1186	35	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	593	0	617	18	2
3	F	593	0	617	26	0
4	D	1112	0	1111	39	0
4	G	1095	0	1099	30	0
5	A	4	0	0	0	0
All	All	6742	0	6795	206	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 206 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:87:LYS:HE3	2:E:121:LEU:HD21	1.13	1.12
4:D:30:THR:HG21	4:D:123:ARG:HH12	1.16	1.08
3:F:42:ARG:HD2	3:F:72:ARG:HE	1.25	1.00
3:C:39:ASP:O	3:C:72:ARG:HD3	1.62	0.99
4:G:35:LEU:HD12	4:G:45:ARG:O	1.63	0.99

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:57:SER:OG	3:C:57:SER:OG[5_675]	1.70	0.50
3:C:57:SER:CB	3:C:57:SER:OG[5_675]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	127/133 (96%)	110 (87%)	15 (12%)	2 (2%)	9 34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	145/154 (94%)	138 (95%)	6 (4%)	1 (1%)	22	55
2	E	145/154 (94%)	133 (92%)	10 (7%)	2 (1%)	11	37
3	C	73/76 (96%)	71 (97%)	2 (3%)	0	100	100
3	F	73/76 (96%)	70 (96%)	3 (4%)	0	100	100
4	D	138/147 (94%)	136 (99%)	2 (1%)	0	100	100
4	G	136/147 (92%)	132 (97%)	4 (3%)	0	100	100
All	All	837/887 (94%)	790 (94%)	42 (5%)	5 (1%)	25	57

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	120	PRO
1	A	246	ARG
1	A	186	HIS
2	E	100	GLN
2	E	19	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/119 (99%)	116 (98%)	2 (2%)	60	80
2	B	125/128 (98%)	125 (100%)	0	100	100
2	E	125/128 (98%)	123 (98%)	2 (2%)	62	81
3	C	67/68 (98%)	67 (100%)	0	100	100
3	F	67/68 (98%)	67 (100%)	0	100	100
4	D	125/129 (97%)	123 (98%)	2 (2%)	62	81
4	G	123/129 (95%)	123 (100%)	0	100	100
All	All	750/769 (98%)	744 (99%)	6 (1%)	81	91

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	120	GLN
2	E	44	ASP
4	D	123	ARG
1	A	228	CYS
2	E	28	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/133 (96%)	-0.08	2 (1%) 72 70	88, 134, 180, 231	0
2	B	147/154 (95%)	-0.10	0 100 100	89, 136, 184, 194	0
2	E	147/154 (95%)	-0.09	1 (0%) 87 87	96, 157, 204, 251	0
3	C	75/76 (98%)	-0.34	0 100 100	97, 145, 187, 213	0
3	F	75/76 (98%)	0.38	3 (4%) 38 37	95, 167, 203, 280	0
4	D	140/147 (95%)	-0.13	2 (1%) 75 74	89, 122, 161, 219	0
4	G	138/147 (93%)	-0.25	1 (0%) 87 87	83, 124, 158, 203	0
All	All	851/887 (95%)	-0.10	9 (1%) 80 79	83, 136, 190, 280	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	76	GLY	6.2
3	F	41	GLN	4.5
1	A	251	HIS	3.2
4	D	145	ASN	3.1
2	E	76	TYR	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ZN	A	1262	1/1	0.97	0.19	113,113,113,113	0
5	ZN	A	1260	1/1	0.98	0.21	115,115,115,115	0
5	ZN	A	1263	1/1	0.98	0.19	118,118,118,118	0
5	ZN	A	1261	1/1	0.99	0.21	102,102,102,102	0

6.5 Other polymers [i](#)

There are no such residues in this entry.