

wwPDB X-ray Structure Validation Summary Report (i)

Aug 30, 2023 – 04:20 pm BST

PDB ID	:	8AMC
Title	:	Crystal Structure of cat allergen Fel d 4
Authors	:	Schooltink, L.; Sagmeister, T.; Gottstein, N.; Pavkov-Keller, T.; Keller, W.
Deposited on	:	2022-08-03
Resolution	:	2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	193	^{2%} 76%	6%	18%
1	В	193	% 76%	6%	18%
1	С	193	.% 7 5%	7%	19%
1	D	193	% 70%	11%	19%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 10211 atoms, of which 4964 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	Δ	158	Total	С	Η	Ν	Ο	\mathbf{S}	0	0	0
	А	100	2533	827	1244	210	248	4	0	0	U
1	В	158	Total	С	Η	Ν	Ο	S	0	0	0
		100	2533	827	1244	210	248	4	0	0	0
1	С	157	Total	С	Η	Ν	0	S	0	0	0
	C	0 107	2518	822	1238	209	245	4	0	0	0
1	р	157	Total	С	Η	Ν	0	S	0	0	0
	I D	157	2518	822	1238	209	245	4	0	0	0

• Molecule 1 is a protein called Allergen Fel d 4.

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-21	MET	-	initiating methionine	UNP Q5VFH6
А	-20	GLY	-	expression tag	UNP Q5VFH6
А	-19	SER	-	expression tag	UNP Q5VFH6
A	-18	SER	-	expression tag	UNP Q5VFH6
А	-17	HIS	-	expression tag	UNP Q5VFH6
А	-16	HIS	-	expression tag	UNP Q5VFH6
А	-15	HIS	-	expression tag	UNP Q5VFH6
A	-14	HIS	-	expression tag	UNP Q5VFH6
А	-13	HIS	-	expression tag	UNP Q5VFH6
А	-12	HIS	-	expression tag	UNP Q5VFH6
А	-11	SER	-	expression tag	UNP Q5VFH6
А	-10	SER	-	expression tag	UNP Q5VFH6
А	-9	GLY	-	expression tag	UNP Q5VFH6
А	-8	GLU	-	expression tag	UNP Q5VFH6
A	-7	ASN	-	expression tag	UNP Q5VFH6
А	-6	LEU	-	expression tag	UNP Q5VFH6
A	-5	TYR	-	expression tag	UNP Q5VFH6
А	-4	PHE	-	expression tag	UNP Q5VFH6
А	-3	GLN	-	expression tag	UNP Q5VFH6
А	-2	GLY	-	expression tag	UNP Q5VFH6
А	-1	HIS	-	expression tag	UNP Q5VFH6



8A	١M	\mathbf{C}
<u> </u>		~


Continued from previous page							
Chain	Residue	Modelled	Actual	Comment	Reference		
А	0	MET	-	expression tag	UNP Q5VFH6		
В	-21	MET	-	initiating methionine	UNP Q5VFH6		
В	-20	GLY	-	expression tag	UNP Q5VFH6		
В	-19	SER	-	expression tag	UNP Q5VFH6		
В	-18	SER	-	expression tag	UNP Q5VFH6		
В	-17	HIS	-	expression tag	UNP Q5VFH6		
В	-16	HIS	-	expression tag	UNP Q5VFH6		
В	-15	HIS	-	expression tag	UNP Q5VFH6		
В	-14	HIS	-	expression tag	UNP Q5VFH6		
В	-13	HIS	-	expression tag	UNP Q5VFH6		
В	-12	HIS	-	expression tag	UNP Q5VFH6		
В	-11	SER	-	expression tag	UNP Q5VFH6		
В	-10	SER	-	expression tag	UNP Q5VFH6		
В	-9	GLY	-	expression tag	UNP Q5VFH6		
В	-8	GLU	-	expression tag	UNP Q5VFH6		
В	-7	ASN	-	expression tag	UNP Q5VFH6		
В	-6	LEU	-	expression tag	UNP Q5VFH6		
В	-5	TYR	-	expression tag	UNP Q5VFH6		
В	-4	PHE	-	expression tag	UNP Q5VFH6		
В	-3	GLN	-	expression tag	UNP Q5VFH6		
В	-2	GLY	-	expression tag	UNP Q5VFH6		
В	-1	HIS	-	expression tag	UNP Q5VFH6		
В	0	MET	-	expression tag	UNP Q5VFH6		
С	-21	MET	-	initiating methionine	UNP Q5VFH6		
С	-20	GLY	-	expression tag	UNP Q5VFH6		
С	-19	SER	-	expression tag	UNP Q5VFH6		
С	-18	SER	-	expression tag	UNP Q5VFH6		
С	-17	HIS	-	expression tag	UNP Q5VFH6		
С	-16	HIS	-	expression tag	UNP Q5VFH6		
С	-15	HIS	-	expression tag	UNP Q5VFH6		
С	-14	HIS	-	expression tag	UNP Q5VFH6		
С	-13	HIS	-	expression tag	UNP Q5VFH6		
С	-12	HIS	-	expression tag	UNP Q5VFH6		
С	-11	SER	-	expression tag	UNP Q5VFH6		
С	-10	SER	-	expression tag	UNP Q5VFH6		
С	-9	GLY	-	expression tag	UNP Q5VFH6		
С	-8	GLU	-	expression tag	UNP Q5VFH6		
С	-7	ASN	-	expression tag	UNP Q5VFH6		
С	-6	LEU	-	expression tag	UNP Q5VFH6		
С	-5	TYR	-	expression tag	UNP Q5VFH6		
С	-4	PHE	-	expression tag	UNP Q5VFH6		
С	-3	GLN	-	expression tag	UNP Q5VFH6		

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	GLY	-	expression tag	UNP Q5VFH6
С	-1	HIS	-	expression tag	UNP Q5VFH6
С	0	MET	-	expression tag	UNP Q5VFH6
D	-21	MET	-	initiating methionine	UNP Q5VFH6
D	-20	GLY	-	expression tag	UNP Q5VFH6
D	-19	SER	-	expression tag	UNP Q5VFH6
D	-18	SER	-	expression tag	UNP Q5VFH6
D	-17	HIS	-	expression tag	UNP Q5VFH6
D	-16	HIS	-	expression tag	UNP Q5VFH6
D	-15	HIS	-	expression tag	UNP Q5VFH6
D	-14	HIS	-	expression tag	UNP Q5VFH6
D	-13	HIS	-	expression tag	UNP Q5VFH6
D	-12	HIS	-	expression tag	UNP Q5VFH6
D	-11	SER	-	expression tag	UNP Q5VFH6
D	-10	SER	-	expression tag	UNP Q5VFH6
D	-9	GLY	-	expression tag	UNP Q5VFH6
D	-8	GLU	-	expression tag	UNP Q5VFH6
D	-7	ASN	-	expression tag	UNP Q5VFH6
D	-6	LEU	-	expression tag	UNP Q5VFH6
D	-5	TYR	-	expression tag	UNP Q5VFH6
D	-4	PHE	-	expression tag	UNP Q5VFH6
D	-3	GLN	-	expression tag	UNP Q5VFH6
D	-2	GLY	-	expression tag	UNP Q5VFH6
D	-1	HIS	-	expression tag	UNP Q5VFH6
D	0	MET	-	expression tag	UNP Q5VFH6

Continued from previous page...







Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	33	Total O 33 33	0	0
3	В	21	TotalO2121	0	0
3	С	26	TotalO2626	0	0
3	D	19	Total O 19 19	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Allergen Fel d 4



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	67.29Å 120.57Å 67.15Å	Denesiten
a, b, c, α , β , γ	90.00° 106.51° 90.00°	Depositor
Resolution (Å)	34.11 - 2.95	Depositor
Resolution (A)	34.11 - 2.95	EDS
% Data completeness	99.6 (34.11-2.95)	Depositor
(in resolution range)	99.7 (34.11 - 2.95)	EDS
R _{merge}	0.14	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.92 (at 2.95Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC 5.8.0267	Depositor
P. P.	0.179 , 0.230	Depositor
R, R_{free}	0.185 , 0.234	DCC
R_{free} test set	1063 reflections $(4.92%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	68.8	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36 , 44.4	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	0.157 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10211	wwPDB-VP
Average B, all atoms $(Å^2)$	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $\mathrm{SO4}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	Bond lengths		nd angles
	Mol Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.77	0/1313	0.89	0/1773
1	В	0.78	0/1313	0.91	1/1773~(0.1%)
1	С	0.78	0/1304	0.93	0/1761
1	D	0.77	0/1304	0.88	0/1761
All	All	0.77	0/5234	0.90	1/7068~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
1	С	0	1
1	D	0	1
All	All	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	160	ARG	CB-CA-C	5.98	122.37	110.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	4	GLU	Peptide
1	С	34	GLU	Peptide



Continued from previous page...

Mol	Chain	Res	Type	Group
1	D	34	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1289	1244	1274	4	1
1	В	1289	1244	1274	6	11
1	С	1280	1238	1268	8	13
1	D	1280	1238	1268	17	2
2	С	5	0	0	0	0
2	D	5	0	0	0	0
3	А	33	0	0	0	0
3	В	21	0	0	0	0
3	С	26	0	0	1	0
3	D	19	0	0	0	1
All	All	5247	4964	5084	34	14

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:49:LEU:O	1:D:50:ASP:OD1	1.83	0.94
1:D:50:ASP:OD1	1:D:53:SER:HB2	1.91	0.69
1:D:110:ASP:OD1	1:D:111:LYS:N	2.28	0.65
1:C:43:VAL:HG21	1:C:121:TYR:CZ	2.33	0.64
1:C:112:THR:HG22	1:C:113:ARG:H	1.63	0.63

The worst 5 of 14 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:LYS:NZ	1:C:141:GLU:O[1_655]	0.94	1.26



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:LYS:HZ3	1:C:141:GLU:O[1_655]	0.54	1.06
1:B:111:LYS:NZ	1:C:141:GLU:C[1_655]	1.40	0.80
1:B:111:LYS:HZ2	$1:C:141:GLU:C[1_{655}]$	0.83	0.77
1:B:111:LYS:HZ1	1:C:142:HIS:HA[1_655]	0.92	0.68

Continued from previous page...

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	156/193~(81%)	145~(93%)	11 (7%)	0	100 100
1	В	156/193~(81%)	150 (96%)	6 (4%)	0	100 100
1	С	155/193~(80%)	148 (96%)	7 (4%)	0	100 100
1	D	155/193~(80%)	146 (94%)	9~(6%)	0	100 100
All	All	622/772 (81%)	589~(95%)	33~(5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	146/177~(82%)	145~(99%)	1 (1%)	84 93
1	В	146/177~(82%)	146 (100%)	0	100 100
1	С	145/177~(82%)	145 (100%)	0	100 100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	145/177~(82%)	145 (100%)	0	100 100
All	All	582/708~(82%)	581 (100%)	1 (0%)	93 98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type	
1	А	36	ASN	

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	108	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol Type C	Chain	Res	Link	Bond lengths			Bond angles			
	Unam			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
2	SO4	С	201	-	4,4,4	0.36	0	$6,\!6,\!6$	0.09	0
2	SO4	D	201	-	4,4,4	0.34	0	$6,\!6,\!6$	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	158/193~(81%)	0.10	3 (1%) 66 49	38, 61, 116, 146	0
1	В	158/193~(81%)	0.05	1 (0%) 89 78	37, 59, 110, 132	0
1	С	157/193~(81%)	0.02	1 (0%) 89 78	32, 62, 101, 129	0
1	D	157/193~(81%)	0.10	1 (0%) 89 78	37, 68, 106, 138	0
All	All	630/772~(81%)	0.07	6 (0%) 82 68	32, 62, 109, 146	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	5	ASN	3.6
1	А	79	GLY	3.5
1	В	5	ASN	2.2
1	А	78	ASP	2.1
1	D	110	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	SO4	С	201	5/5	0.88	0.20	110,116,128,146	0
2	SO4	D	201	5/5	0.89	0.16	101,114,118,118	0

6.5 Other polymers (i)

There are no such residues in this entry.

