



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2021 – 03:17 pm BST

PDB ID : 7AQZ  
Title : Co-Crystal Structure of Variant Surface Glycoprotein VSG2 in complex with Nanobody VSG2(NB14)  
Authors : Stebbins, C.E.; Hempelmann, A.; VanStraaten, M.  
Deposited on : 2020-10-23  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	 96%
1	B	364	 96%
2	C	130	 95%
2	D	130	 68% 28%
3	E	4	 75% 25%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	4	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14807 atoms, of which 6579 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Variant surface glycoprotein MITAT 1.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	358	Total	C	H	N	O	S	0	11	0
			5375	1704	2635	478	547	11			
1	B	361	Total	C	H	N	O	S	0	14	0
			5550	1740	2752	491	556	11			

- Molecule 2 is a protein called Nanobody VSG2(NB14).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	C	124	Total	C	H	N	O	S	0	5	0
			1742	599	774	169	195	5			
2	D	93	Total	C	H	N	O	S	0	0	0
			991	353	398	108	128	4			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
3	E	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	F	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	B	1	Total	C	H	O	0	0
			18	6	5	7		
4	B	1	Total	C	H	O	0	1
			36	12	10	14		
4	D	1	Total	C	H	O	0	0
			18	6	5	7		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Na	0	0
			1	1		
5	D	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	395	Total	O	0	0
			395	395		
6	B	426	Total	O	0	0
			426	426		
6	C	114	Total	O	0	0
			114	114		
6	D	40	Total	O	0	0
			40	40		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Variant surface glycoprotein MITAT 1.2




- Molecule 1: Variant surface glycoprotein MITAT 1.2



- Molecule 2: Nanobody VSG2(NB14)



- Molecule 2: Nanobody VSG2(NB14)



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.54Å 95.95Å 124.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.24 – 1.30 74.24 – 1.30	Depositor EDS
% Data completeness (in resolution range)	98.3 (74.24-1.30) 98.3 (74.24-1.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.37 (at 1.30Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, $R_{free}$	0.151 , 0.172 0.151 , 0.172	Depositor DCC
$R_{free}$ test set	13427 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.278	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14807	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, NA, CIT, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.50	1/2781 (0.0%)	0.68	1/3761 (0.0%)
1	B	0.50	0/2845	0.66	1/3844 (0.0%)
2	C	0.54	0/985	0.71	0/1332
2	D	0.42	0/597	0.86	3/793 (0.4%)
All	All	0.50	1/7208 (0.0%)	0.69	5/9730 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	385	GLU	CD-OE2	-5.26	1.19	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	99	ASP	CB-CG-OD1	6.32	123.99	118.30
2	D	99	ASP	CB-CG-OD2	-5.98	112.92	118.30
1	B	130	ARG	NE-CZ-NH1	5.92	123.26	120.30
2	D	65	ARG	O-C-N	-5.68	113.55	123.20
1	A	378	MET	CG-SD-CE	-5.04	92.13	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	63	SER	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2740	2635	2702	6	0
1	B	2798	2752	2790	5	0
2	C	968	774	891	0	0
2	D	593	398	473	1	0
3	E	50	0	43	0	0
3	F	50	0	43	0	0
4	B	39	15	15	1	0
4	D	13	5	5	1	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	395	0	0	2	0
6	B	426	0	0	0	0
6	C	114	0	0	0	0
6	D	40	0	0	0	0
All	All	8228	6579	6962	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:97:HIS:CE1	1:B:297:LYS:HG2	2.35	0.61
1:A:172:LYS:NZ	6:A:501:HOH:O	1.97	0.60
1:A:194[B]:VAL:HG12	6:A:764:HOH:O	2.04	0.56
1:A:140:GLN:HB3	1:B:234[B]:VAL:HG12	1.99	0.45
1:B:258:LEU:HD22	1:B:270[B]:ILE:CD1	2.46	0.45
1:A:382:GLY:HA2	1:A:385:GLU:HG2	1.99	0.44
1:A:236[B]:MET:SD	1:A:243:VAL:HG21	2.58	0.43
1:B:89:LYS:HD2	1:B:326:TYR:CE1	2.54	0.43
4:D:502:CIT:O1	4:D:502:CIT:C6	2.66	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:306[B]:GLU:OE2	4:B:401:CIT:O3	2.35	0.42
1:A:382:GLY:O	1:A:385:GLU:HG2	2.20	0.42
2:D:22:CYS:HB3	2:D:79:VAL:CG1	2.50	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/364 (100%)	355 (97%)	10 (3%)	0	100	100
1	B	371/364 (102%)	361 (97%)	10 (3%)	0	100	100
2	C	125/130 (96%)	122 (98%)	3 (2%)	0	100	100
2	D	83/130 (64%)	78 (94%)	5 (6%)	0	100	100
All	All	944/988 (96%)	916 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/279 (100%)	280 (100%)	0	100	100
1	B	290/279 (104%)	286 (99%)	4 (1%)	67	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	98/105 (93%)	97 (99%)	1 (1%)	76	48
2	D	49/105 (47%)	49 (100%)	0	100	100
All	All	717/768 (93%)	712 (99%)	5 (1%)	86	61

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	83	HIS
1	B	217[A]	SER
1	B	217[B]	SER
1	B	312	LYS
2	C	124	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	3	GLN
2	C	13	GLN
2	C	39	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	E	1	1,3	14,14,15	0.72	0	17,19,21	0.93	0
3	NAG	E	2	3	14,14,15	0.69	1 (7%)	17,19,21	1.00	1 (5%)
3	BMA	E	3	3	11,11,12	0.43	0	15,15,17	0.72	0
3	MAN	E	4	3	11,11,12	0.32	0	15,15,17	0.56	0
3	NAG	F	1	1,3	14,14,15	0.85	1 (7%)	17,19,21	0.81	0
3	NAG	F	2	3	14,14,15	0.83	0	17,19,21	1.12	1 (5%)
3	BMA	F	3	3	11,11,12	0.66	0	15,15,17	0.74	0
3	MAN	F	4	3	11,11,12	0.37	0	15,15,17	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	2/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	NAG	O5-C1	-2.10	1.40	1.43
3	F	1	NAG	C1-C2	-2.02	1.49	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	2	NAG	O5-C1-C2	-3.00	106.55	111.29
3	E	2	NAG	C4-C3-C2	-2.20	107.80	111.02

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	4	MAN	O5-C5-C6-O6
3	E	4	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	CIT	B	402[A]	-	3,12,12	0.83	0	3,17,17	1.04	0
4	CIT	B	401	-	3,12,12	0.75	0	3,17,17	0.27	0
4	CIT	D	502	-	3,12,12	0.76	0	3,17,17	0.49	0
4	CIT	B	402[B]	-	3,12,12	1.30	0	3,17,17	2.58	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	B	402[A]	-	-	2/6/16/16	-
4	CIT	B	401	-	-	0/6/16/16	-
4	CIT	D	502	-	-	0/6/16/16	-
4	CIT	B	402[B]	-	-	1/6/16/16	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	402[B]	CIT	C3-C2-C1	4.26	121.81	114.98

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	402[A]	CIT	C2-C3-C4-C5
4	B	402[A]	CIT	C1-C2-C3-O7
4	B	402[B]	CIT	C1-C2-C3-O7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	401	CIT	1	0
4	D	502	CIT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	358/364 (98%)	-0.42	2 (0%) 89 88	12, 19, 39, 57	0
1	B	361/364 (99%)	-0.32	3 (0%) 86 86	13, 19, 33, 51	0
2	C	124/130 (95%)	-0.19	2 (1%) 72 73	15, 23, 42, 63	0
2	D	93/130 (71%)	0.85	15 (16%) 1 1	18, 45, 70, 80	0
All	All	936/988 (94%)	-0.23	22 (2%) 59 58	12, 20, 50, 80	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	106	ASP	5.2
2	D	60	TYR	4.1
2	D	108	THR	4.0
1	B	184	ALA	3.9
2	D	37	PHE	3.6
1	B	389	GLN	3.1
2	D	107	GLY	2.8
2	D	105	LYS	2.8
2	D	63	SER	2.6
2	D	47	PHE	2.6
2	D	61	ALA	2.4
1	B	292	ALA	2.4
2	C	106	ASP	2.3
2	D	83	MET	2.2
2	D	81	LEU	2.1
2	D	62	ASP	2.1
2	D	117	GLY	2.1
1	A	388	THR	2.1
2	D	93	VAL	2.1
1	A	352	VAL	2.0
2	D	66	GLY	2.0
2	C	124	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BMA	F	3	11/12	0.72	0.12	25,36,42,52	0
3	MAN	E	4	11/12	0.84	0.15	33,36,43,43	0
3	MAN	F	4	11/12	0.88	0.17	28,57,72,77	0
3	BMA	E	3	11/12	0.90	0.14	27,35,56,62	0
3	NAG	E	1	14/15	0.95	0.06	17,18,22,25	0
3	NAG	F	2	14/15	0.95	0.08	14,22,44,51	0
3	NAG	E	2	14/15	0.96	0.06	18,23,32,33	0
3	NAG	F	1	14/15	0.97	0.08	16,16,19,21	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CIT	B	401	13/13	0.75	0.29	32,40,58,63	0
4	CIT	B	402[A]	13/13	0.86	0.16	12,16,54,65	18
4	CIT	B	402[B]	13/13	0.86	0.16	12,16,19,20	18
4	CIT	D	502	13/13	0.92	0.16	29,40,66,85	0
5	NA	D	501	1/1	0.94	0.09	21,21,21,21	0
5	NA	C	201	1/1	1.00	0.14	14,14,14,14	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.