



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 7, 2022 – 11:13 AM EDT

PDB ID : 2ARK  
Title : Structure of a flavodoxin from *Aquifex aeolicus*  
Authors : Cuff, M.E.; Quartey, P.; Zhou, M.; Cymborowski, M.; Minor, W.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2005-08-19  
Resolution : 2.40 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	 4% 74% 20% 5% ..
1	B	188	 4% 77% 18% ..
1	C	188	 4% 78% 16% ..
1	D	188	 7% 73% 19% 6% ..
1	E	188	 4% 68% 23% 5% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	188	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	501	-	-	X	-
3	GOL	A	512	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flavodoxin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	187	1445	917	251	267	3	7	0	0	0
1	B	186	1439	914	250	265	3	7	0	0	0
1	C	186	1439	914	249	266	3	7	0	0	0
1	D	187	1445	917	251	267	3	7	0	0	0
1	E	185	1434	911	248	265	3	7	0	0	0
1	F	185	1431	910	248	263	3	7	0	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	cloning artifact	UNP O67866
A	-1	ASN	-	cloning artifact	UNP O67866
A	0	ALA	-	cloning artifact	UNP O67866
A	1	MSE	MET	modified residue	UNP O67866
A	18	MSE	MET	modified residue	UNP O67866
A	61	MSE	MET	modified residue	UNP O67866
A	68	MSE	MET	modified residue	UNP O67866
A	106	MSE	MET	modified residue	UNP O67866
A	111	MSE	MET	modified residue	UNP O67866
A	113	MSE	MET	modified residue	UNP O67866
B	-2	SER	-	cloning artifact	UNP O67866
B	-1	ASN	-	cloning artifact	UNP O67866
B	0	ALA	-	cloning artifact	UNP O67866
B	1	MSE	MET	modified residue	UNP O67866
B	18	MSE	MET	modified residue	UNP O67866
B	61	MSE	MET	modified residue	UNP O67866
B	68	MSE	MET	modified residue	UNP O67866

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	106	MSE	MET	modified residue	UNP O67866
B	111	MSE	MET	modified residue	UNP O67866
B	113	MSE	MET	modified residue	UNP O67866
C	-2	SER	-	cloning artifact	UNP O67866
C	-1	ASN	-	cloning artifact	UNP O67866
C	0	ALA	-	cloning artifact	UNP O67866
C	1	MSE	MET	modified residue	UNP O67866
C	18	MSE	MET	modified residue	UNP O67866
C	61	MSE	MET	modified residue	UNP O67866
C	68	MSE	MET	modified residue	UNP O67866
C	106	MSE	MET	modified residue	UNP O67866
C	111	MSE	MET	modified residue	UNP O67866
C	113	MSE	MET	modified residue	UNP O67866
D	-2	SER	-	cloning artifact	UNP O67866
D	-1	ASN	-	cloning artifact	UNP O67866
D	0	ALA	-	cloning artifact	UNP O67866
D	1	MSE	MET	modified residue	UNP O67866
D	18	MSE	MET	modified residue	UNP O67866
D	61	MSE	MET	modified residue	UNP O67866
D	68	MSE	MET	modified residue	UNP O67866
D	106	MSE	MET	modified residue	UNP O67866
D	111	MSE	MET	modified residue	UNP O67866
D	113	MSE	MET	modified residue	UNP O67866
E	-2	SER	-	cloning artifact	UNP O67866
E	-1	ASN	-	cloning artifact	UNP O67866
E	0	ALA	-	cloning artifact	UNP O67866
E	1	MSE	MET	modified residue	UNP O67866
E	18	MSE	MET	modified residue	UNP O67866
E	61	MSE	MET	modified residue	UNP O67866
E	68	MSE	MET	modified residue	UNP O67866
E	106	MSE	MET	modified residue	UNP O67866
E	111	MSE	MET	modified residue	UNP O67866
E	113	MSE	MET	modified residue	UNP O67866
F	-2	SER	-	cloning artifact	UNP O67866
F	-1	ASN	-	cloning artifact	UNP O67866
F	0	ALA	-	cloning artifact	UNP O67866
F	1	MSE	MET	modified residue	UNP O67866
F	18	MSE	MET	modified residue	UNP O67866
F	61	MSE	MET	modified residue	UNP O67866
F	68	MSE	MET	modified residue	UNP O67866
F	106	MSE	MET	modified residue	UNP O67866
F	111	MSE	MET	modified residue	UNP O67866

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	113	MSE	MET	modified residue	UNP O67866

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	69	Total O 69 69	0	0
4	B	95	Total O 95 95	0	0
4	C	55	Total O 55 55	0	0
4	D	75	Total O 75 75	0	0
4	E	42	Total O 42 42	0	0
4	F	84	Total O 84 84	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Flavodoxin

Chain A: 




- Molecule 1: Flavodoxin

Chain B: 





- Molecule 1: Flavodoxin

Chain C: 




- Molecule 1: Flavodoxin

Chain D: 










- Molecule 1: Flavodoxin



- Molecule 1: Flavodoxin



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.85Å 112.85Å 150.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.00 – 2.40 36.94 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (36.00-2.40) 99.6 (36.94-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.02 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.171 , 0.235 0.169 , 0.233	Depositor DCC
$R_{free}$ test set	2205 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.5	Xtrriage
Anisotropy	0.506	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9119	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.80	0/1464	0.89	6/1956 (0.3%)
1	B	0.75	0/1458	0.95	6/1948 (0.3%)
1	C	0.93	3/1458 (0.2%)	1.06	8/1948 (0.4%)
1	D	1.11	4/1464 (0.3%)	0.92	8/1956 (0.4%)
1	E	1.46	9/1454 (0.6%)	0.97	7/1944 (0.4%)
1	F	0.70	0/1450	0.83	4/1937 (0.2%)
All	All	0.99	16/8748 (0.2%)	0.94	39/11689 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	-2	SER	CB-OG	27.80	1.78	1.42
1	E	171	GLU	CD-OE2	25.95	1.54	1.25
1	E	169	ARG	CZ-NH2	24.22	1.64	1.33
1	E	171	GLU	CD-OE1	23.67	1.51	1.25
1	C	169	ARG	CZ-NH2	13.32	1.50	1.33

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	169	ARG	NE-CZ-NH1	-22.91	108.84	120.30
1	E	169	ARG	NE-CZ-NH2	15.00	127.80	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	169	ARG	NE-CZ-NH2	14.29	127.44	120.30
1	E	169	ARG	NE-CZ-NH1	-9.42	115.59	120.30
1	B	156	ARG	NE-CZ-NH1	8.98	124.79	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	171	GLU	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1445	0	1450	48	0
1	B	1439	0	1445	34	0
1	C	1439	0	1443	28	0
1	D	1445	0	1450	31	0
1	E	1434	0	1438	43	0
1	F	1431	0	1439	20	1
2	A	5	0	0	3	0
2	B	5	0	0	1	0
2	C	5	0	0	1	0
2	D	5	0	0	0	0
2	E	5	0	0	1	0
2	F	5	0	0	1	0
3	A	12	0	16	0	0
3	B	6	0	8	1	0
3	C	12	0	16	3	0
3	D	6	0	8	1	0
4	A	69	0	0	3	0
4	B	95	0	0	7	0
4	C	55	0	0	1	0
4	D	75	0	0	3	0
4	E	42	0	0	1	0
4	F	84	0	0	4	1
All	All	9119	0	8713	188	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 188 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:169:ARG:CD	1:E:169:ARG:CG	1.79	1.55
1:C:0:ALA:CA	1:C:0:ALA:N	1.68	1.50
1:F:1:MSE:CE	1:F:1:MSE:SE	2.14	1.46
1:C:61:MSE:SE	1:C:61:MSE:CE	2.17	1.42
1:A:61:MSE:CE	1:A:61:MSE:SE	2.17	1.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:17:LYS:NZ	4:F:553:HOH:O[6_765]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/188 (98%)	181 (98%)	2 (1%)	2 (1%)	14	20
1	B	184/188 (98%)	179 (97%)	5 (3%)	0	100	100
1	C	184/188 (98%)	181 (98%)	3 (2%)	0	100	100
1	D	185/188 (98%)	178 (96%)	7 (4%)	0	100	100
1	E	183/188 (97%)	180 (98%)	3 (2%)	0	100	100
1	F	183/188 (97%)	175 (96%)	7 (4%)	1 (0%)	29	41
All	All	1104/1128 (98%)	1074 (97%)	27 (2%)	3 (0%)	41	55

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	169	ARG
1	F	64	VAL
1	A	64	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/143 (104%)	136 (91%)	13 (9%)	10	15
1	B	148/143 (104%)	128 (86%)	20 (14%)	4	4
1	C	148/143 (104%)	131 (88%)	17 (12%)	5	7
1	D	149/143 (104%)	130 (87%)	19 (13%)	4	5
1	E	148/143 (104%)	130 (88%)	18 (12%)	5	6
1	F	147/143 (103%)	125 (85%)	22 (15%)	3	3
All	All	889/858 (104%)	780 (88%)	109 (12%)	4	6

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	75	VAL
1	E	39	VAL
1	F	119	VAL
1	D	118	LEU
1	D	158	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	60	ASN
1	F	60	ASN
1	F	101	ASN
1	B	60	ASN
1	B	101	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	F	506	-	4,4,4	1.31	0	6,6,6	1.28	1 (16%)
2	PO4	D	504	-	4,4,4	0.78	0	6,6,6	0.93	0
3	GOL	C	508	-	5,5,5	0.38	0	5,5,5	0.56	0
2	PO4	E	505	-	4,4,4	0.74	0	6,6,6	0.68	0
3	GOL	C	511	-	5,5,5	0.34	0	5,5,5	0.62	0
2	PO4	A	501	-	4,4,4	1.83	2 (50%)	6,6,6	0.69	0
3	GOL	A	512	-	5,5,5	0.44	0	5,5,5	0.31	0
3	GOL	A	509	-	5,5,5	0.49	0	5,5,5	0.73	0
3	GOL	B	510	-	5,5,5	0.41	0	5,5,5	0.58	0
3	GOL	D	507	-	5,5,5	0.28	0	5,5,5	0.75	0
2	PO4	C	503	-	4,4,4	0.75	0	6,6,6	0.71	0
2	PO4	B	502	-	4,4,4	1.26	0	6,6,6	1.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	508	-	-	4/4/4/4	-
3	GOL	C	511	-	-	2/4/4/4	-
3	GOL	A	512	-	-	2/4/4/4	-
3	GOL	A	509	-	-	4/4/4/4	-
3	GOL	B	510	-	-	4/4/4/4	-
3	GOL	D	507	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	PO4	P-O4	-2.21	1.48	1.54
2	A	501	PO4	P-O2	-2.20	1.48	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	506	PO4	O4-P-O1	-2.21	102.80	110.89

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	512	GOL	C1-C2-C3-O3
3	B	510	GOL	C1-C2-C3-O3
3	C	508	GOL	O1-C1-C2-C3
3	C	508	GOL	C1-C2-C3-O3
3	D	507	GOL	O1-C1-C2-C3

There are no ring outliers.

9 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	506	PO4	1	0
3	C	508	GOL	2	0
2	E	505	PO4	1	0
3	C	511	GOL	1	0
2	A	501	PO4	3	0
3	B	510	GOL	1	0
3	D	507	GOL	1	0
2	C	503	PO4	1	0
2	B	502	PO4	1	0



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/188 (95%)	-0.27	8 (4%) 34 33	19, 28, 37, 44	0
1	B	179/188 (95%)	-0.21	8 (4%) 33 31	18, 27, 35, 46	0
1	C	179/188 (95%)	0.02	7 (3%) 39 38	20, 28, 36, 48	0
1	D	180/188 (95%)	-0.04	13 (7%) 15 14	13, 28, 40, 51	0
1	E	178/188 (94%)	-0.06	7 (3%) 39 38	17, 28, 36, 49	0
1	F	178/188 (94%)	-0.29	7 (3%) 39 38	21, 28, 36, 52	0
All	All	1074/1128 (95%)	-0.14	50 (4%) 31 30	13, 28, 36, 52	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	103	VAL	4.3
1	A	103	VAL	4.2
1	E	185	ASP	4.1
1	C	103	VAL	3.9
1	D	0	ALA	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	510	6/6	0.68	0.25	73,75,76,76	0
3	GOL	C	511	6/6	0.69	0.34	59,61,61,62	0
3	GOL	A	512	6/6	0.73	0.99	61,64,64,65	0
3	GOL	A	509	6/6	0.87	0.38	43,46,49,49	0
3	GOL	D	507	6/6	0.87	0.34	35,39,41,42	0
2	PO4	C	503	5/5	0.88	0.17	48,50,52,53	0
3	GOL	C	508	6/6	0.89	0.27	35,38,38,38	0
2	PO4	E	505	5/5	0.93	0.15	32,35,36,37	0
2	PO4	B	502	5/5	0.95	0.18	34,36,39,40	5
2	PO4	F	506	5/5	0.95	0.24	33,35,37,38	5
2	PO4	A	501	5/5	0.95	0.22	18,23,24,25	5
2	PO4	D	504	5/5	0.95	0.13	48,50,51,52	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.