



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 07:44 pm GMT

PDB ID : 5AR1
Title : Crystal structure of Cdc11 from *Saccharomyces cerevisiae*
Authors : Brausemann, A.; Gerhardt, S.; Schott, A.K.; Einsle, O.; Grosse-Berkenbusch, A.; Johnsson, N.; Gronemeyer, T.
Deposited on : 2015-09-23
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION CONTROL PROTEIN 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	216	1643	1045	277	316	5	0	0	1

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	MET	-	expression tag	UNP P32458
A	-3	SER	-	expression tag	UNP P32458
A	-2	SER	-	expression tag	UNP P32458
A	-1	HIS	-	expression tag	UNP P32458
A	0	HIS	-	expression tag	UNP P32458
A	1	HIS	-	expression tag	UNP P32458
A	2	HIS	-	expression tag	UNP P32458
A	3	HIS	-	expression tag	UNP P32458
A	4	HIS	-	expression tag	UNP P32458
A	5	SER	-	expression tag	UNP P32458
A	6	SER	-	expression tag	UNP P32458
A	7	ALA	-	expression tag	UNP P32458
A	8	MET	-	expression tag	UNP P32458
A	9	ALA	-	expression tag	UNP P32458
A	10	SER	-	expression tag	UNP P32458
A	11	THR	-	expression tag	UNP P32458
A	12	ALA	-	expression tag	UNP P32458
A	13	GLU	-	expression tag	UNP P32458
A	14	ASN	-	expression tag	UNP P32458
A	15	LEU	-	expression tag	UNP P32458
A	16	TYR	-	expression tag	UNP P32458
A	17	PHE	-	expression tag	UNP P32458
A	18	GLN	-	expression tag	UNP P32458
A	19	GLY	-	expression tag	UNP P32458

4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	187.05Å 187.05Å 57.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.20 – 2.85 61.23 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.20-2.85) 99.9 (61.23-2.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 2.86Å)	Xtrriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.257 , 0.259 0.281 , 0.275	Depositor DCC
R_{free} test set	682 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	112.0	Xtrriage
Anisotropy	0.099	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 145.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1643	wwPDB-VP
Average B, all atoms (Å ²)	167.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1667	0.72	1/2261 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	210	SER	C-N-CA	5.13	134.52	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1643	0	1546	17	0
All	All	1643	0	1546	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:ILE:HD11	1:A:232:ASN:ND2	2.09	0.68
1:A:136:HIS:HA	1:A:283:THR:HG21	1.83	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ILE:HD11	1:A:197:LEU:HD13	1.89	0.55
1:A:283:THR:HA	1:A:287:LEU:HB2	1.92	0.52
1:A:219:TYR:O	1:A:222:THR:HB	2.13	0.48
1:A:23:PHE:HE1	1:A:25:VAL:HB	1.77	0.48
1:A:228:ILE:HD11	1:A:232:ASN:HD21	1.78	0.47
1:A:178:ARG:HH12	1:A:237:MET:HA	1.79	0.47
1:A:169:VAL:HG12	1:A:226:PHE:HB2	1.99	0.44
1:A:28:VAL:HG11	1:A:156:PHE:CZ	2.53	0.43
1:A:21:ILE:HG21	1:A:277:HIS:HE1	1.83	0.43
1:A:24:THR:H	1:A:136:HIS:CE1	2.37	0.43
1:A:72:THR:HA	1:A:85:LEU:O	2.19	0.42
1:A:167:ILE:HG22	1:A:200:TYR:HB3	2.03	0.41
1:A:287:LEU:HD12	1:A:287:LEU:HA	1.93	0.40
1:A:154:VAL:HG22	1:A:197:LEU:HD11	2.04	0.40
1:A:228:ILE:HA	1:A:233:GLU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	204/303 (67%)	177 (87%)	18 (9%)	9 (4%)	2 8

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	32	GLY
1	A	211	ASP
1	A	233	GLU
1	A	146	GLY
1	A	147	HIS
1	A	209	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	208	GLU
1	A	232	ASN
1	A	102	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	171/277 (62%)	149 (87%)	22 (13%)	4 11

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ILE
1	A	68	LEU
1	A	69	ARG
1	A	72	THR
1	A	103	PHE
1	A	107	SER
1	A	117	ILE
1	A	134	ARG
1	A	158	ARG
1	A	163	LEU
1	A	174	ASP
1	A	181	LEU
1	A	194	ARG
1	A	199	ILE
1	A	215	GLU
1	A	220	LEU
1	A	222	THR
1	A	224	LEU
1	A	276	LEU
1	A	287	LEU
1	A	289	GLU
1	A	296	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	165	ASN
1	A	184	ASN
1	A	232	ASN
1	A	277	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/303 (71%)	0.78	41 (18%) 1 1	103, 171, 220, 247	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	23	PHE	7.9
1	A	296	LEU	6.9
1	A	163	LEU	5.4
1	A	114	TYR	5.0
1	A	292	ARG	4.6
1	A	149	LEU	4.4
1	A	288	TYR	4.2
1	A	103	PHE	4.1
1	A	238	GLY	3.9
1	A	156	PHE	3.8
1	A	160	LEU	3.7
1	A	173	SER	3.4
1	A	290	ARG	3.4
1	A	297	SER	3.3
1	A	135	VAL	3.2
1	A	106	ILE	3.2
1	A	157	ILE	3.1
1	A	119	LEU	3.1
1	A	110	ILE	3.0
1	A	291	TYR	2.9
1	A	294	GLU	2.9
1	A	295	ALA	2.9
1	A	267	LEU	2.8
1	A	261	ILE	2.8
1	A	25	VAL	2.8
1	A	141	LEU	2.8
1	A	176	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	271	LEU	2.5
1	A	228	ILE	2.4
1	A	293	THR	2.4
1	A	101	PRO	2.3
1	A	168	PRO	2.3
1	A	69	ARG	2.2
1	A	265	VAL	2.1
1	A	166	ILE	2.1
1	A	220	LEU	2.1
1	A	161	GLY	2.1
1	A	237	MET	2.1
1	A	276	LEU	2.0
1	A	24	THR	2.0
1	A	226	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.