

Full wwPDB X-ray Structure Validation Report (i)

Mar 3, 2024 – 01:34 AM EST

PDB ID : 6ARR

Title : Aspergillus fumigatus Cytosolic Thiolase: Apo enzyme in complex with cesium

ions

Authors: Marshall, A.C.; Bond, C.S.; Bruning, J.B.

Deposited on : 2017-08-23

Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	399	88%	11%
1	В	399	91%	8% •
1	С	399	94%	5% •
1	D	399	94%	5% •



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 12895 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Acetyl-CoA acetyltransferase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	395	Total	С	N	О	S	0	6	0
1	A	390	2853	1800	490	552	11	0	0	
1	В	396	Total	С	N	О	S	0	2	0
1	Б	390	2851	1793	494	552	12	U		
1	С	396	Total	С	N	О	S	0	9	0
1		390	2883	1822	494	555	12	U	9	
1	1 D	D 20"	Total	С	N	О	S	0	1	0
1	D	395	2841	1790	488	551	12	U	4	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q4WCL5
В	0	GLY	-	expression tag	UNP Q4WCL5
С	0	GLY	-	expression tag	UNP Q4WCL5
D	0	GLY	-	expression tag	UNP Q4WCL5

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	В	1	Total Cl 1 1	0	0
2	С	2	Total Cl 2 2	0	0

• Molecule 3 is CESIUM ION (three-letter code: CS) (formula: Cs) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Cs 3 3	0	0
3	В	3	Total Cs 3 3	0	0
3	С	4	Total Cs 4 4	0	0
3	D	5	Total Cs 5 5	0	0

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	156	Total O 156 156	0	0
5	В	190	Total O 190 190	0	0
5	С	540	Total O 540 540	0	0

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	544	Total O 544 544	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetyl-CoA acetyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	71.41Å 105.27Å 110.45Å	Depositor
a, b, c, α , β , γ	90.00° 108.61° 90.00°	Depositor
Resolution (Å)	56.93 - 1.82	Depositor
Resolution (A)	104.67 - 1.82	EDS
% Data completeness	100.0 (56.93-1.82)	Depositor
(in resolution range)	100.0 (104.67-1.82)	EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.43 (at 1.82Å)	Xtriage
Refinement program	PHENIX dev_2747	Depositor
D.D.	0.177 , 0.209	Depositor
R, R_{free}	0.177 , 0.208	DCC
R_{free} test set	6801 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 55.9	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12895	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.30	0/2916	0.49	0/3960
1	В	0.30	0/2900	0.48	0/3932
1	С	0.47	0/2953	0.58	0/4006
1	D	0.46	0/2896	0.58	0/3932
All	All	0.39	0/11665	0.54	0/15830

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2853	0	2904	28	0
1	В	2851	0	2913	20	0
1	С	2883	0	2966	12	0
1	D	2841	0	2891	10	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	2	0	0	0	0
3	A	3	0	0	0	0
3	В	3	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	4	0	0	0	0
3	D	5	0	0	0	0
4	С	6	0	8	0	0
4	D	12	0	16	1	0
5	A	156	0	0	6	1
5	В	190	0	0	3	1
5	С	540	0	0	3	0
5	D	544	0	0	3	0
All	All	12895	0	11698	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:4:LEU:HD13	1:C:107:MET:HE1	1.57	0.85
1:B:112:ASP:HB3	1:B:269:LYS:HG3	1.62	0.79
1:A:299:ALA:HA	1:A:336:LEU:HD21	1.68	0.75
1:C:1:MET:HA	1:C:4:LEU:HD12	1.72	0.71
1:D:188:GLU:OE1	5:D:501:HOH:O	2.10	0.70
1:B:290[A]:GLN:HB2	1:B:293:LYS:HG3	1.74	0.69
1:C:195[B]:LYS:NZ	5:C:501:HOH:O	2.25	0.69
1:A:345:LEU:HD13	1:A:374:LYS:HE3	1.76	0.66
1:B:290[B]:GLN:HB3	1:B:293:LYS:HG3	1.78	0.64
1:B:166:GLU:HG2	1:B:241:ILE:HD12	1.82	0.62
1:A:270:LEU:HD12	1:A:275:LEU:HB2	1.83	0.60
1:D:362:ARG:O	1:D:365[B]:THR:HG22	2.03	0.58
1:A:253[B]:SER:OG	1:A:323:GLU:O	2.21	0.58
1:D:195:LYS:HE3	5:D:793:HOH:O	2.04	0.57
1:A:184:ILE:O	1:A:188:GLU:HG2	2.05	0.57
1:A:284:TRP:HB3	1:A:393:LEU:HD12	1.86	0.56
1:B:320:GLU:HG3	1:B:367:LEU:HB2	1.88	0.55
1:C:1:MET:N	5:C:507:HOH:O	2.40	0.55
1:A:45:ASP:OD1	1:A:45:ASP:N	2.40	0.55
1:A:187:TYR:HB3	1:A:228:LEU:HD22	1.87	0.55
1:C:362:ARG:O	1:C:365[B]:THR:HG22	2.07	0.54
1:C:342:LYS:NZ	5:C:502:HOH:O	2.27	0.54
1:A:189:LYS:O	5:A:501:HOH:O	2.19	0.54
1:B:179:GLN:NE2	5:B:507:HOH:O	2.40	0.53
1:A:177:GLU:O	1:A:181:GLU:HG2	2.08	0.53

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:282:LEU:HB3	1:A:309:ALA:HB1	1.89	0.53
1:B:269:LYS:HD3	1:B:273:LEU:HD23	1.91	0.52
1:C:201:GLU:OE1	1:C:372:LYS:NZ	2.31	0.52
1:B:157[B]:LYS:NZ	5:B:508:HOH:O	2.43	0.51
1:A:349:ALA:N	5:A:511:HOH:O	2.42	0.48
1:A:90:LYS:NZ	5:A:508:HOH:O	2.36	0.48
1:A:322:ASN:ND2	5:A:504:HOH:O	2.30	0.47
1:A:168:CYS:HA	5:A:549:HOH:O	2.14	0.47
1:B:132:LEU:HD21	1:D:141:GLN:HG3	1.96	0.47
1:C:84[A]:ILE:HD13	1:D:286:ASP:HB2	1.96	0.47
1:A:167:GLU:OE2	1:A:293:LYS:NZ	2.46	0.47
1:A:288:ALA:O	1:B:82:SER:HA	2.15	0.46
1:A:346:HIS:ND1	1:A:370:VAL:HG22	2.30	0.46
1:A:82:SER:HA	1:B:288:ALA:O	2.16	0.46
1:B:238:PRO:HA	1:B:248:THR:HG22	1.98	0.46
1:B:377:LYS:HD2	1:B:377:LYS:HA	1.69	0.46
1:B:11:SER:OG	1:B:40:ALA:O	2.23	0.46
1:A:132:LEU:HD21	1:C:141:GLN:HG3	1.97	0.45
1:A:187:TYR:CD1	1:A:228:LEU:HB2	2.52	0.45
1:A:278:VAL:O	1:A:397:LEU:HD12	2.17	0.45
1:D:131:ASN:N	1:D:131:ASN:OD1	2.51	0.44
1:C:323:GLU:HB3	1:C:350:VAL:HG23	1.98	0.44
1:A:238:PRO:HA	1:A:248:THR:HG22	2.00	0.44
1:B:162:GLY:HA3	1:B:240:PHE:CZ	2.53	0.44
1:A:374:LYS:HG2	5:A:506:HOH:O	2.18	0.43
1:D:8:TYR:OH	1:D:267:GLU:OE2	2.29	0.43
1:B:276:LYS:H	1:B:276:LYS:HG2	1.65	0.42
1:B:304:LYS:NZ	5:B:518:HOH:O	2.52	0.42
1:B:141:GLN:HG3	1:D:132:LEU:HD21	2.02	0.41
1:C:320:GLU:HG3	1:C:367:LEU:HB2	2.01	0.41
1:A:154:ASP:OD2	1:A:157:LYS:HB2	2.21	0.41
1:A:181:GLU:OE2	1:A:234:ARG:NH1	2.53	0.41
1:C:249:ALA:HB3	1:C:250:PRO:HD3	2.01	0.41
1:D:320:GLU:HG3	1:D:367:LEU:HB2	2.02	0.41
1:B:229:ASN:ND2	1:B:232:LYS:HD2	2.36	0.41
4:D:402:GOL:O3	5:D:503:HOH:O	2.22	0.41
1:A:323:GLU:HB3	1:A:350:VAL:HG23	2.02	0.41
1:A:171:ASP:OD2	1:A:293:LYS:HE3	2.21	0.40
1:B:381:ALA:O	1:B:392:ALA:HA	2.21	0.40
1:D:162:GLY:HA3	1:D:240:PHE:CZ	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-



metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
5:A:634:HOH:O	5:B:675:HOH:O[1_455]	2.12	0.08

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	399/399 (100%)	391 (98%)	7 (2%)	1 (0%)	41	27
1	В	394/399~(99%)	385 (98%)	8 (2%)	1 (0%)	41	27
1	С	401/399 (100%)	394 (98%)	6 (2%)	1 (0%)	47	33
1	D	395/399~(99%)	389 (98%)	5 (1%)	1 (0%)	41	27
All	All	$1589/1596 \ (100\%)$	1559 (98%)	26 (2%)	4 (0%)	41	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	VAL
1	В	91	VAL
1	С	91	VAL
1	D	91	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	294/298~(99%)	288 (98%)	6 (2%)	55 43
1	В	295/298~(99%)	293 (99%)	2 (1%)	84 80
1	C	300/298 (101%)	296 (99%)	4 (1%)	69 61
1	D	293/298 (98%)	290 (99%)	3 (1%)	76 70
All	All	1182/1192 (99%)	1167 (99%)	15 (1%)	69 61

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	PHE
1	A	157	LYS
1	A	180	ASP
1	A	186	THR
1	A	211	PHE
1	A	346	HIS
1	В	57	PHE
1	В	272	GLU
1	С	1	MET
1	С	57	PHE
1	С	92	CYS
1	С	393	LEU
1	D	57	PHE
1	D	92	CYS
1	D	377	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 19 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res Link		В	ond leng	$_{ m gths}$	Е	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	GOL	D	402	-	5,5,5	0.50	0	5,5,5	1.31	1 (20%)
4	GOL	D	401	-	5,5,5	0.97	0	5,5,5	1.06	1 (20%)
4	GOL	С	403	-	5,5,5	0.58	0	5,5,5	1.03	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	402	-	-	0/4/4/4	-
4	GOL	D	401	-	-	2/4/4/4	-
4	GOL	С	403	-	-	0/4/4/4	ı

There are no bond length outliers.

All (3) bond angle outliers are listed below:

	Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
ſ	4	D	402	GOL	C3-C2-C1	-2.64	101.43	111.70
	4	D	401	GOL	C3-C2-C1	-2.08	103.62	111.70
	4	С	403	GOL	C3-C2-C1	-2.03	103.82	111.70

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	401	GOL	C1-C2-C3-O3
4	D	401	GOL	O2-C2-C3-O3



There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	402	GOL	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ} {>} 2$	$OWAB(A^2)$	Q<0.9
1	A	395/399 (98%)	0.22	22 (5%) 24 19	21, 50, 75, 111	0
1	В	396/399 (99%)	-0.13	8 (2%) 65 61	22, 43, 65, 106	0
1	С	396/399 (99%)	-0.54	2 (0%) 91 89	8, 16, 34, 76	0
1	D	395/399 (98%)	-0.58	4 (1%) 82 80	8, 15, 31, 79	0
All	All	1582/1596 (99%)	-0.26	36 (2%) 60 56	8, 27, 67, 111	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	215	PRO	4.5
1	A	210	GLY	4.2
1	A	235	ALA	4.0
1	A	165	ALA	3.9
1	A	211	PHE	3.6
1	A	245	GLY	3.5
1	A	236	ILE	3.5
1	С	2	SER	3.3
1	В	209	PRO	3.1
1	D	209	PRO	2.9
1	A	370	VAL	2.9
1	С	1	MET	2.9
1	A	169	ALA	2.9
1	В	398	LEU	2.9
1	A	243	GLY	2.8
1	A	212	ARG	2.7
1	A	173	GLY	2.7
1	A	341[A]	GLU	2.7
1	В	210	GLY	2.6
1	В	242	PRO	2.6
1	A	181	GLU	2.5

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	317	ASP	2.5
1	A	184	ILE	2.4
1	В	244	SER	2.4
1	D	1	MET	2.4
1	A	342	LYS	2.4
1	D	2	SER	2.4
1	A	44	VAL	2.3
1	A	296	THR	2.3
1	A	172	HIS	2.2
1	D	215	PRO	2.1
1	В	241	ILE	2.1
1	A	325	PHE	2.1
1	A	174	PHE	2.1
1	В	214	LYS	2.0
1	A	398	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	GOL	D	402	6/6	0.90	0.29	20,20,20,20	0
4	GOL	С	403	6/6	0.92	0.12	20,20,20,20	0
4	GOL	D	401	6/6	0.95	0.07	21,30,43,48	0
3	CS	D	407	1/1	0.95	0.06	70,70,70,70	1
2	CL	A	401	1/1	0.96	0.08	27,27,27,27	1
2	CL	В	401	1/1	0.96	0.06	27,27,27,27	1
3	CS	В	404	1/1	0.97	0.06	89,89,89,89	1
3	CS	В	403	1/1	0.98	0.08	65,65,65,65	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\AA^2)$	Q<0.9
2	CL	С	402	1/1	0.98	0.07	22,22,22,22	1
2	CL	С	401	1/1	0.99	0.07	19,19,19,19	0
3	CS	A	402	1/1	0.99	0.09	54,54,54,54	0
3	CS	A	403	1/1	0.99	0.05	73,73,73,73	1
3	CS	С	406	1/1	1.00	0.08	25,25,25,25	1
3	CS	С	407	1/1	1.00	0.06	41,41,41,41	1
3	CS	D	403	1/1	1.00	0.10	14,14,14,14	0
3	CS	D	404	1/1	1.00	0.10	20,20,20,20	1
3	CS	D	405	1/1	1.00	0.10	25,25,25,25	1
3	CS	D	406	1/1	1.00	0.09	26,26,26,26	1
3	CS	A	404	1/1	1.00	0.07	72,72,72,72	0
3	CS	В	402	1/1	1.00	0.08	45,45,45,45	0
3	CS	С	404	1/1	1.00	0.09	16,16,16,16	0
3	CS	С	405	1/1	1.00	0.10	25,25,25,25	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around CS D 407:












Electron density around CL A 401:













Electron density around CL B 401:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)

















Electron density around CS B 403:










Electron density around CL C 402: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around CL C 401:












Electron density around CS A 402:













Electron density around CS A 403:









Electron density around CS C 406:












Electron density around CS C 407:







Electron density around CS D 403: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around CS D 404: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around CS D 405: $2 \mathrm{mF}_o\text{-DF}_c \text{ (at } 0.7 \text{ rmsd) in gray} \\ \mathrm{mF}_o\text{-DF}_c \text{ (at } 3 \text{ rmsd) in purple (negative)}$

and green (positive)














Electron density around CS A 404:









Electron density around CS B 402: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

