



wwPDB EM Validation Summary Report ⓘ

Mar 3, 2024 – 03:48 PM EST

PDB ID : 6AWD
EMDB ID : EMD-7016
Title : Structure of 30S (S1 depleted) ribosomal subunit and RNA polymerase complex
Authors : Demo, G.; Rasouly, A.; Vasilyev, N.; Loveland, A.B.; Diaz-Avalos, R.; Grigorieff, N.; Nudler, E.; Korostelev, A.A.
Deposited on : 2017-09-05
Resolution : 8.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 8.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1539	
2	01	229	
2	02	229	
3	03	1340	
4	04	1369	
5	05	59	
6	E	218	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	F	206	 52% 46%
8	G	205	 45% 50%
9	H	157	 52% 47%
10	I	100	 53% 45%
11	J	151	 48% 50%
12	K	129	 47% 49%
13	L	127	 49% 46% 5%
14	M	98	 40% 57%
15	N	116	 51% 47%
16	O	123	 53% 45%
17	P	114	 50% 47%
18	Q	100	 45% 51%
19	R	88	 57% 43%
20	S	82	 46% 49% 5%
21	T	80	 60% 36%
22	U	65	 43% 55%
23	V	79	 43% 53%
24	W	85	 47% 51%
25	X	65	 5% 57% 34% 8%

2 Entry composition [i](#)

There are 25 unique types of molecules in this entry. The entry contains 76452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
1	A	1539	33012	14725	6052	10697	1538	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	01	227	1753	1091	311	345	6	0	0
2	02	227	1757	1094	311	346	6	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	03	1327	10272	6441	1787	2001	43	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	04	1345	10372	6508	1853	1962	49	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	05	59	458	283	86	88	1	0	0

- Molecule 6 is a protein called 30S ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	E	218	Total	C	N	O	S	0	0
			1705	1081	305	312	7		

- Molecule 7 is a protein called 30S ribosomal protein S3.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	F	206	Total	C	N	O	S	0	0
			1625	1028	305	289	3		

- Molecule 8 is a protein called 30S ribosomal protein S4.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	G	205	Total	C	N	O	S	0	0
			1643	1026	315	298	4		

- Molecule 9 is a protein called 30S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	H	157	Total	C	N	O	S	0	0
			1157	719	218	214	6		

- Molecule 10 is a protein called 30S ribosomal protein S6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	I	100	Total	C	N	O	S	0	0
			818	515	148	149	6		

- Molecule 11 is a protein called 30S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	J	151	Total	C	N	O	S	0	0
			1182	735	227	216	4		

- Molecule 12 is a protein called 30S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	K	129	Total	C	N	O	S	0	0
			979	616	173	184	6		

- Molecule 13 is a protein called 30S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	L	127	Total	C	N	O	S	0	0
			1022	634	206	179	3		

- Molecule 14 is a protein called 30S ribosomal protein S10.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	M	98	Total	C	N	O	S	0	0
			787	493	150	143	1		

- Molecule 15 is a protein called 30S ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	N	116	Total	C	N	O	S	0	0
			870	535	173	159	3		

- Molecule 16 is a protein called 30S ribosomal protein S12.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	O	123	Total	C	N	O	S	0	0
			955	590	196	165	4		

- Molecule 17 is a protein called 30S ribosomal protein S13.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	P	114	Total	C	N	O	S	0	0
			884	546	178	157	3		

- Molecule 18 is a protein called 30S ribosomal protein S14.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	Q	100	Total	C	N	O	S	0	0
			805	499	164	139	3		

- Molecule 19 is a protein called 30S ribosomal protein S15.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	R	88	Total	C	N	O	S	0	0
			714	439	144	130	1		

- Molecule 20 is a protein called 30S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	S	82	Total	C	N	O	S	0	0
			649	406	128	114	1		

- Molecule 21 is a protein called 30S ribosomal protein S17.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	T	80	Total	C	N	O	S	0	0
			649	411	121	114	3		

- Molecule 22 is a protein called 30S ribosomal protein S18.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	U	65	Total	C	N	O	S	0	0
			536	339	100	96	1		

- Molecule 23 is a protein called 30S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	V	79	Total	C	N	O	S	0	0
			638	408	120	108	2		

- Molecule 24 is a protein called 30S ribosomal protein S20.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	W	85	Total	C	N	O	S	0	0
			665	411	137	114	3		


- Molecule 25 is a protein called 30S ribosomal protein S21.


Mol	Chain	Residues	Atoms					AltConf	Trace
25	X	65	Total	C	N	O	S	0	0
			545	335	117	92	1		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 16S rRNA







• Molecule 2: DNA-directed RNA polymerase subunit alpha






• Molecule 3: DNA-directed RNA polymerase subunit beta








● Molecule 4: DNA-directed RNA polymerase subunit beta'









• Molecule 5: DNA-directed RNA polymerase subunit omega




• Molecule 6: 30S ribosomal protein S2




• Molecule 7: 30S ribosomal protein S3



• Molecule 8: 30S ribosomal protein S4






• Molecule 9: 30S ribosomal protein S5




• Molecule 10: 30S ribosomal protein S6



• Molecule 11: 30S ribosomal protein S7




• Molecule 12: 30S ribosomal protein S8





• Molecule 13: 30S ribosomal protein S9







- Molecule 14: 30S ribosomal protein S10



- Molecule 15: 30S ribosomal protein S11



- Molecule 16: 30S ribosomal protein S12




- Molecule 17: 30S ribosomal protein S13



- Molecule 18: 30S ribosomal protein S14






- Molecule 19: 30S ribosomal protein S15



- Molecule 20: 30S ribosomal protein S16




- Molecule 21: 30S ribosomal protein S17



- Molecule 22: 30S ribosomal protein S18




- Molecule 23: 30S ribosomal protein S19



- Molecule 24: 30S ribosomal protein S20





• Molecule 25: 30S ribosomal protein S21



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	21123	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40.0	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	22000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	13.228	Depositor
Minimum map value	-3.576	Depositor
Average map value	-0.054	Depositor
Map value standard deviation	0.680	Depositor
Recommended contour level	2.0	Depositor
Map size (\AA)	528.64, 528.64, 528.64	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.888, 1.888, 1.888	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/36963	0.66	0/57662
2	01	0.24	0/1774	0.60	0/2405
2	02	0.25	0/1779	0.59	0/2411
3	03	0.28	0/10433	0.62	1/14101 (0.0%)
4	04	0.28	0/10528	0.61	2/14224 (0.0%)
5	05	0.24	0/460	0.54	0/620
6	E	0.36	0/1736	0.77	1/2338 (0.0%)
7	F	0.25	0/1652	0.56	0/2225
8	G	0.27	0/1665	0.60	0/2227
9	H	0.26	0/1170	0.57	0/1573
10	I	0.27	0/836	0.58	0/1128
11	J	0.26	0/1196	0.60	0/1602
12	K	0.28	0/989	0.63	0/1326
13	L	0.26	0/1034	0.58	0/1375
14	M	0.26	0/797	0.66	1/1077 (0.1%)
15	N	0.28	0/886	0.66	1/1195 (0.1%)
16	O	0.27	0/969	0.71	0/1300
17	P	0.25	0/893	0.65	0/1193
18	Q	0.29	0/817	0.63	0/1088
19	R	0.26	0/722	0.64	0/964
20	S	0.33	0/659	0.65	0/884
21	T	0.30	0/658	0.65	0/881
22	U	0.28	0/545	0.68	0/731
23	V	0.29	0/653	0.64	0/877
24	W	0.29	0/671	0.60	0/888
25	X	0.38	0/551	0.74	0/728
All	All	0.27	0/81036	0.64	6/117023 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	26	LYS	N-CA-C	-8.64	87.66	111.00
14	M	58	ASN	N-CA-C	-6.29	94.01	111.00
4	04	642	ASP	N-CA-C	5.62	126.19	111.00
3	03	516	ASP	CB-CG-OD2	5.24	123.02	118.30
15	N	93	GLU	N-CA-C	-5.13	97.15	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	25	C	Sidechain
1	A	872	A	Sidechain
1	A	9	G	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	33012	0	16618	423	0
2	01	1753	0	1780	106	0
2	02	1757	0	1778	96	0
3	03	10272	0	10138	518	0
4	04	10372	0	10512	558	0
5	05	458	0	476	23	0
6	E	1705	0	1732	180	0
7	F	1625	0	1699	78	0
8	G	1643	0	1710	111	0
9	H	1157	0	1199	70	0
10	I	818	0	808	48	0
11	J	1182	0	1240	74	0
12	K	979	0	1034	63	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	L	1022	0	1070	64	0
14	M	787	0	828	57	0
15	N	870	0	878	46	0
16	O	955	0	1019	43	0
17	P	884	0	944	59	0
18	Q	805	0	847	59	0
19	R	714	0	737	31	0
20	S	649	0	666	40	0
21	T	649	0	691	27	0
22	U	536	0	552	32	0
23	V	638	0	665	49	0
24	W	665	0	714	51	0
25	X	545	0	579	25	0
All	All	76452	0	60914	2743	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 2743 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:E:26:LYS:O	6:E:29:PRO:HD2	1.44	1.17
6:E:196:VAL:HG21	6:E:199:VAL:HG13	1.32	1.11
4:04:19:ALA:HB1	4:04:1343:GLU:HB3	1.27	1.08
6:E:11:LYS:O	6:E:13:GLY:N	1.88	1.07
4:04:639:VAL:HG12	4:04:640:GLY:H	1.20	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	01	225/229 (98%)	193 (86%)	24 (11%)	8 (4%)	3	25
2	02	225/229 (98%)	179 (80%)	35 (16%)	11 (5%)	2	20
3	03	1323/1340 (99%)	1036 (78%)	233 (18%)	54 (4%)	3	23
4	04	1339/1369 (98%)	1099 (82%)	206 (15%)	34 (2%)	5	32
5	05	57/59 (97%)	48 (84%)	9 (16%)	0	100	100
6	E	216/218 (99%)	156 (72%)	37 (17%)	23 (11%)	0	8
7	F	204/206 (99%)	180 (88%)	22 (11%)	2 (1%)	15	55
8	G	203/205 (99%)	151 (74%)	42 (21%)	10 (5%)	2	20
9	H	155/157 (99%)	120 (77%)	30 (19%)	5 (3%)	4	26
10	I	98/100 (98%)	78 (80%)	18 (18%)	2 (2%)	7	38
11	J	149/151 (99%)	118 (79%)	26 (17%)	5 (3%)	3	26
12	K	127/129 (98%)	105 (83%)	17 (13%)	5 (4%)	3	23
13	L	125/127 (98%)	101 (81%)	18 (14%)	6 (5%)	2	21
14	M	96/98 (98%)	82 (85%)	10 (10%)	4 (4%)	3	22
15	N	114/116 (98%)	92 (81%)	16 (14%)	6 (5%)	2	19
16	O	121/123 (98%)	88 (73%)	23 (19%)	10 (8%)	1	12
17	P	112/114 (98%)	90 (80%)	17 (15%)	5 (4%)	2	22
18	Q	98/100 (98%)	66 (67%)	24 (24%)	8 (8%)	1	12
19	R	86/88 (98%)	72 (84%)	12 (14%)	2 (2%)	6	34
20	S	80/82 (98%)	58 (72%)	16 (20%)	6 (8%)	1	13
21	T	78/80 (98%)	61 (78%)	10 (13%)	7 (9%)	1	11
22	U	63/65 (97%)	46 (73%)	13 (21%)	4 (6%)	1	17
23	V	77/79 (98%)	60 (78%)	15 (20%)	2 (3%)	5	31
24	W	83/85 (98%)	69 (83%)	13 (16%)	1 (1%)	13	50
25	X	63/65 (97%)	45 (71%)	10 (16%)	8 (13%)	0	5
All	All	5517/5614 (98%)	4393 (80%)	896 (16%)	228 (4%)	5	23

5 of 228 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	01	161	SER
2	01	168	ILE
2	02	138	ALA
2	02	168	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	03	596	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	01	194/197 (98%)	191 (98%)	3 (2%)	65 80
2	02	194/197 (98%)	190 (98%)	4 (2%)	53 72
3	03	1098/1155 (95%)	1075 (98%)	23 (2%)	53 72
4	04	1103/1141 (97%)	1090 (99%)	13 (1%)	71 83
5	05	49/49 (100%)	49 (100%)	0	100 100
6	E	180/180 (100%)	166 (92%)	14 (8%)	12 36
7	F	170/170 (100%)	167 (98%)	3 (2%)	59 77
8	G	172/172 (100%)	164 (95%)	8 (5%)	26 51
9	H	119/119 (100%)	117 (98%)	2 (2%)	60 78
10	I	87/87 (100%)	86 (99%)	1 (1%)	73 84
11	J	124/124 (100%)	123 (99%)	1 (1%)	81 89
12	K	104/104 (100%)	101 (97%)	3 (3%)	42 64
13	L	105/105 (100%)	104 (99%)	1 (1%)	76 86
14	M	86/86 (100%)	86 (100%)	0	100 100
15	N	89/89 (100%)	85 (96%)	4 (4%)	27 52
16	O	103/103 (100%)	99 (96%)	4 (4%)	32 56
17	P	92/92 (100%)	91 (99%)	1 (1%)	73 84
18	Q	83/83 (100%)	80 (96%)	3 (4%)	35 59
19	R	76/76 (100%)	75 (99%)	1 (1%)	69 81
20	S	65/65 (100%)	64 (98%)	1 (2%)	65 80
21	T	74/74 (100%)	72 (97%)	2 (3%)	44 65
22	U	56/56 (100%)	55 (98%)	1 (2%)	59 77
23	V	70/70 (100%)	69 (99%)	1 (1%)	67 80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
24	W	65/65 (100%)	62 (95%)	3 (5%)	27	52
25	X	55/55 (100%)	51 (93%)	4 (7%)	14	39
All	All	4613/4714 (98%)	4512 (98%)	101 (2%)	54	71

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	F	2	GLN
11	J	78	ARG
25	X	47	ARG
7	F	163	ARG
8	G	187	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 93 such sidechains are listed below:

Mol	Chain	Res	Type
9	H	131	ASN
13	L	30	ASN
9	H	145	ASN
11	J	96	ASN
15	N	37	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1538/1539 (99%)	182 (11%)	3 (0%)

5 of 182 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	6	G
1	A	9	G
1	A	22	G
1	A	31	G
1	A	32	A

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	69	G
1	A	343	U
1	A	1399	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-7016. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 148



Y Index: 147




Z Index: 147


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 1882 nm^3 ; this corresponds to an approximate mass of 1700 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.123\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-7016 and PDB model 6AWD. Per-residue inclusion information can be found in section [3](#) on page [8](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 2.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.0).























































9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (2.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8680	 0.0700
01	 0.7340	 0.0450
02	 0.6820	 0.0520
03	 0.6420	 0.0520
04	 0.6410	 0.0530
05	 0.1430	 0.0260
A	 0.9640	 0.0820
E	 0.9680	 0.0910
F	 0.9960	 0.0960
G	 0.9960	 0.0570
H	 0.9980	 0.0960
I	 0.9950	 0.0990
J	 0.9920	 0.0570
K	 0.9990	 0.0910
L	 0.9980	 0.0490
M	 0.9990	 0.0710
N	 0.9990	 0.0750
O	 0.9760	 0.0720
P	 0.9930	 0.0620
Q	 1.0000	 0.0320
R	 1.0000	 0.0850
S	 0.9920	 0.0700
T	 0.9970	 0.0910
U	 1.0000	 0.0750
V	 0.9940	 0.0430
W	 0.9860	 0.0690
X	 0.9420	 0.0230

