



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2020 – 08:23 pm BST

PDB ID : 2AZX
Title : Charged and uncharged tRNAs adopt distinct conformations when complexed with human tryptophanyl-tRNA synthetase
Authors : Yang, X.L.; Otero, F.J.; Ewalt, K.L.; Liu, J.; Swairjo, M.A.; Kohrer, C.; RajBhandary, U.L.; Skene, R.J.; McRee, D.E.; Schimmel, P.
Deposited on : 2005-09-12
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)
RNA backbone	3102	1227 (3.10-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	75	
1	D	75	
2	A	477	
2	B	477	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 9445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 72-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	72	Total	C	N	O	P	0	0	0
			1541	686	276	507	72			
1	D	72	Total	C	N	O	P	0	0	0
			1541	686	276	507	72			

- Molecule 2 is a protein called Tryptophanyl-tRNA synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	A	387	Total	C	N	O	S	Se	0	0	0
			3114	1994	527	578	5	10			
2	B	388	Total	C	N	O	S	Se	0	0	0
			3119	1997	528	579	5	10			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	42	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	48	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	143	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	169	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	195	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	213	GLY	SER	VARIANT	UNP P23381
A	214	ASP	TYR	VARIANT	UNP P23381
A	241	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	243	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	319	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	350	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	401	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	425	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	461	MSE	MET	MODIFIED RESIDUE	UNP P23381
A	472	HIS	-	EXPRESSION TAG	UNP P23381
A	473	HIS	-	EXPRESSION TAG	UNP P23381

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	474	HIS	-	EXPRESSION TAG	UNP P23381
A	475	HIS	-	EXPRESSION TAG	UNP P23381
A	476	HIS	-	EXPRESSION TAG	UNP P23381
A	477	HIS	-	EXPRESSION TAG	UNP P23381
B	1	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	42	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	48	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	143	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	169	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	195	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	213	GLY	SER	VARIANT	UNP P23381
B	214	ASP	TYR	VARIANT	UNP P23381
B	241	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	243	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	319	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	350	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	401	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	425	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	461	MSE	MET	MODIFIED RESIDUE	UNP P23381
B	472	HIS	-	EXPRESSION TAG	UNP P23381
B	473	HIS	-	EXPRESSION TAG	UNP P23381
B	474	HIS	-	EXPRESSION TAG	UNP P23381
B	475	HIS	-	EXPRESSION TAG	UNP P23381
B	476	HIS	-	EXPRESSION TAG	UNP P23381
B	477	HIS	-	EXPRESSION TAG	UNP P23381

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is TRYPTOPHAN (three-letter code: TRP) (formula: C₁₁H₁₂N₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	A	1	Total	C	N	O	0	0
			15	11	2	2		
4	A	1	Total	C	N	O	0	0
			15	11	2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	2	Total	O	0	0
			2	2		
7	D	2	Total	O	0	0
			2	2		
7	A	9	Total	O	0	0
			9	9		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	9	Total	O	0	0
			9	9		



MSE	ASP	M152	S245	Q347	I442	PRO	CYS	G246	Q347	A443
ASN	PRO	K153	F247	T348	A444	ASN	PRO	F247	K349	E444
GLY	PRO	K154	Y248	M350	H445	GLY	PRO	Y248	K351	H446
PRO	ASN	Y159	K249	S351	A447	PRO	ASN	K249	A352	R448
ALA	PRO	R162	N250	A352	R448	ALA	PRO	N250	S353	R449
SER	ALA	G163	V251	D354	P463	SER	ALA	V251	P355	R464
LEU	PRO	P164	I254	D354	P463	LEU	PRO	I254	N356	K465
LEU	THR	S165	Q255	P355	K465	LEU	THR	Q255	S357	L466
GLU	SER	S166	T259	S357	S467	GLU	SER	S166	S358	D468
LEU	ASN	E167	Q262	S358	F469	LEU	ASN	E167	T364	PHE
PHE	HIS	A168	K264	T364	D469	PHE	HIS	A168	Q367	GLN
ASN	ASN	M169	F269	I367	PHE	ASN	ASN	M169	I367	GLN
ASN	PRO	H173	S279	V372	GLN	ASN	PRO	H173	V372	THR
SER	ASP	L174	F280	N373	HIS	SER	ASP	L174	N373	HIS
ILE	ALA	I175	P281	K374	HIS	ILE	ALA	I175	K374	HIS
THR	THR	P176	A282	G380	HIS	THR	THR	P176	G380	HIS
GLN	GLU	K181	I283	R381	HIS	GLN	GLU	K181	R381	HIS
GLY	GLU	W182	P287	D382	HIS	GLY	GLU	W182	D382	HIS
LEU	LEU	V186	F289	T383	HIS	LEU	LEU	V186	T383	HIS
VAL	VAL	F187	F289	I384	HIS	VAL	VAL	F187	I384	HIS
ARG	ARG	M188	P294	E385	HIS	ARG	ARG	M188	E385	HIS
SER	SER	P190	Q295	E386	HIS	SER	SER	P190	E386	HIS
LEU	LEU	Q194	R298	H387	HIS	LEU	LEU	Q194	H387	HIS
LEU	LEU	M195	D299	R388	HIS	LEU	LEU	M195	R388	HIS
LEU	LEU	I196	R300	F400	HIS	LEU	LEU	I196	F400	HIS
LEU	ASP	D197	I311	M401	HIS	LEU	ASP	D197	M401	HIS
LEU	GLU	E199	D812	Y402	HIS	LEU	GLU	E199	Y402	HIS
LEU	GLU	K200	D813	L403	HIS	LEU	GLU	K200	L403	HIS
LEU	ASP	M204	Q813	T404	HIS	LEU	ASP	M204	T404	HIS
LEU	ALA	D205	D814	F405	HIS	LEU	ALA	D205	F405	HIS
LEU	VAL	L206	P815	F406	HIS	LEU	VAL	L206	F406	HIS
LEU	LYS	T207	P815	E408	HIS	LEU	LYS	T207	E408	HIS
LEU	LYS	L208	D822	L407	HIS	LEU	LYS	L208	L407	HIS
LEU	LEU	D209	V823	D409	HIS	LEU	LEU	D209	D409	HIS
LEU	VAL	Q210	A824	D410	HIS	LEU	VAL	Q210	D410	HIS
LEU	SER	P825	P830	L413	HIS	LEU	SER	P825	L413	HIS
LEU	LEU	M218	P830	K418	HIS	LEU	LEU	M218	K418	HIS
LEU	LYS	H129	A833	M425	HIS	LEU	LYS	H129	M425	HIS
LEU	SER	G226	L334	L426	HIS	LEU	SER	G226	L426	HIS
LEU	TYR	R133	L335	T427	HIS	LEU	TYR	R133	T427	HIS
LEU	LYS	F137	H336	K431	HIS	LEU	LYS	F137	K431	HIS
LEU	ALA	F138	S337	I435	HIS	LEU	ALA	F138	I435	HIS
LEU	ALA	S139	T338	Q439	HIS	LEU	ALA	S139	Q439	HIS
LEU	ALA	H140	F339	P440	HIS	LEU	ALA	H140	P440	HIS
LEU	GLY	D239	F840	L441	HIS	LEU	GLY	D239	L441	HIS
LEU	GLY	Y240	P841			LEU	GLY	Y240		
LEU	ASP	M143	Q844			LEU	ASP	M143		
LEU	TYR	I144				LEU	TYR	I144		
LEU	LYS	L147				LEU	LYS	L147		

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	114.06Å 132.62Å 246.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 45.27 – 2.79	Depositor EDS
% Data completeness (in resolution range)	89.0 (50.00-2.80) 91.9 (45.27-2.79)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.77Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.205 , 0.252 0.205 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	67.1	Xtrriage
Anisotropy	0.295	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9445	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.36	1/1722 (0.1%)	0.70	0/2682
1	D	0.40	1/1722 (0.1%)	0.71	1/2682 (0.0%)
2	A	0.41	0/3179	0.65	0/4276
2	B	0.44	0/3184	0.67	0/4283
All	All	0.41	2/9807 (0.0%)	0.68	1/13923 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	501	G	OP3-P	-7.40	1.52	1.61
1	D	501	G	OP3-P	-7.17	1.52	1.61

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	560	U	N1-C1'-C2'	5.06	120.58	114.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	560	U	Sidechain
1	D	560	U	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1541	0	777	48	0
1	D	1541	0	777	43	0
2	A	3114	0	3061	138	0
2	B	3119	0	3063	116	0
3	A	20	0	0	1	0
3	B	30	0	0	2	0
4	A	30	0	18	1	0
4	B	15	0	9	0	0
5	B	1	0	0	0	0
6	B	12	0	16	2	0
7	A	9	0	0	1	0
7	B	9	0	0	0	0
7	C	2	0	0	0	0
7	D	2	0	0	0	0
All	All	9445	0	7721	328	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 328 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:465:LYS:HD3	2:B:465:LYS:H	1.02	1.16
2:A:364:THR:H	2:A:367:GLN:NE2	1.53	1.07
1:D:518:G:H2'	1:D:557:G:N2	1.75	1.01
1:D:558:A:H4'	1:D:559:A:OP1	1.57	0.99
2:A:143:MSE:HE1	2:A:306:LEU:HD11	1.42	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	385/477 (81%)	358 (93%)	25 (6%)	2 (0%)	29	61
2	B	386/477 (81%)	358 (93%)	21 (5%)	7 (2%)	8	28
All	All	771/954 (81%)	716 (93%)	46 (6%)	9 (1%)	13	39

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	168	ALA
2	B	165	SER
2	B	168	ALA
2	B	467	SER
2	B	468	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	340/402 (85%)	322 (95%)	18 (5%)	22	54
2	B	340/402 (85%)	322 (95%)	18 (5%)	22	54
All	All	680/804 (85%)	644 (95%)	36 (5%)	22	54

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	382	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	103	LEU
2	B	385	GLU
2	B	86	ASP
2	B	143	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
2	A	367	GLN
2	A	389	GLN
2	B	387	HIS
2	A	387	HIS
2	B	118	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	C	71/75 (94%)	12 (16%)	4 (5%)
1	D	71/75 (94%)	19 (26%)	5 (7%)
All	All	142/150 (94%)	31 (21%)	9 (6%)

5 of 31 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	C	508	U
1	C	509	G
1	C	510	G
1	C	516	U
1	C	518	G

5 of 9 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	D	507	G
1	D	558	A
1	D	518	G
1	C	518	G
1	D	516	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 1 is monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	703	-	4,4,4	0.28	0	6,6,6	0.18	0
3	SO4	A	704	-	4,4,4	0.26	0	6,6,6	0.18	0
3	SO4	A	708	-	4,4,4	0.29	0	6,6,6	0.10	0
3	SO4	B	706	-	4,4,4	0.26	0	6,6,6	0.15	0
3	SO4	A	707	-	4,4,4	0.29	0	6,6,6	0.07	0
3	SO4	B	710	-	4,4,4	0.26	0	6,6,6	0.07	0
6	GOL	B	801	-	5,5,5	0.39	0	5,5,5	0.35	0
3	SO4	B	705	-	4,4,4	0.28	0	6,6,6	0.14	0
3	SO4	B	701	-	4,4,4	0.29	0	6,6,6	0.07	0
3	SO4	B	709	-	4,4,4	0.28	0	6,6,6	0.06	0
6	GOL	B	802	-	5,5,5	0.38	0	5,5,5	0.30	0
3	SO4	A	702	-	4,4,4	0.26	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	802	-	-	0/4/4/4	-
6	GOL	B	801	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	703	SO4	1	0
3	A	704	SO4	1	0
6	B	801	GOL	2	0
3	B	705	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	C	72/75 (96%)	1.49	24 (33%) 0 0	81, 127, 155, 167	0
1	D	72/75 (96%)	0.91	17 (23%) 0 0	57, 107, 188, 200	0
2	A	377/477 (79%)	-0.07	0 100 100	39, 60, 84, 97	0
2	B	378/477 (79%)	0.01	0 100 100	39, 54, 72, 98	1 (0%)
All	All	899/1104 (81%)	0.17	41 (4%) 32 22	39, 59, 134, 200	1 (0%)

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	570	G	5.9
1	D	502	A	5.7
1	C	520	U	5.3
1	C	516	U	5.1
1	D	572	C	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	B	710	5/5	0.74	0.30	155,155,156,156	0
5	MG	B	901	1/1	0.74	0.28	82,82,82,82	0
6	GOL	B	802	6/6	0.83	0.44	93,94,95,96	0
4	TRP	A	602	15/15	0.84	0.41	103,106,112,113	0
3	SO4	B	709	5/5	0.90	0.16	131,131,131,132	0
3	SO4	A	708	5/5	0.90	0.22	122,123,124,124	0
6	GOL	B	801	6/6	0.90	0.27	64,67,67,71	0
3	SO4	B	706	5/5	0.93	0.14	112,114,114,114	0
3	SO4	A	707	5/5	0.93	0.18	125,125,126,126	0
3	SO4	A	702	5/5	0.94	0.19	100,102,103,103	0
3	SO4	B	705	5/5	0.95	0.17	103,104,104,105	0
4	TRP	A	601	15/15	0.96	0.23	51,53,61,62	0
3	SO4	A	704	5/5	0.96	0.23	76,78,79,80	0
4	TRP	B	603	15/15	0.96	0.28	43,45,55,55	0
3	SO4	B	701	5/5	0.97	0.23	77,79,80,80	0
3	SO4	B	703	5/5	0.98	0.12	72,73,74,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.