wwPDB X-ray Structure Validation Summary Report (i)

Jun 16, 2024 – 11:07 PM EDT

PDB ID	:	3B0V
Title	:	tRNA-dihydrouridine synthase from Thermus thermophilus in complex with tRNA
Authors	:	Yu, F.; Tanaka, Y.; Yamashita, K.; Nakamura, A.; Yao, M.; Tanaka, I.
Deposited on	:	2011-06-14
Resolution	:	3.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1161 (3.60-3.44)
Clashscore	141614	1244 (3.60-3.44)
Ramachandran outliers	138981	1206 (3.60-3.44)
Sidechain outliers	138945	1207 (3.60-3.44)
RSRZ outliers	127900	1080 (3.60-3.44)
RNA backbone	3102	1003 (4.02-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	76	75%; 26%, 46%, 22%, • •
1	B	76	79%; 26%, 46%, 24%, •
2	C	363	73%, 13%, •, 13%
2	D	363	%; 70%, 16%, •, 13%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8204 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called tRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	73	1561	695	280	513	73	0	0	0
1	B	73	1561	695	280	513	73	0	0	0

- Molecule 2 is a protein called tRNA-dihydrouridine synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	C	315	2510	1583	476	439	2	10	0	0	0
2	D	315	2510	1583	476	439	2	10	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-20	MSE	-	EXPRESSION TAG	UNP Q5SMC7
C	-19	GLY	-	EXPRESSION TAG	UNP Q5SMC7
C	-18	SER	-	EXPRESSION TAG	UNP Q5SMC7
C	-17	SER	-	EXPRESSION TAG	UNP Q5SMC7
C	-16	HIS	-	EXPRESSION TAG	UNP Q5SMC7
C	-15	HIS	-	EXPRESSION TAG	UNP Q5SMC7
C	-14	HIS	-	EXPRESSION TAG	UNP Q5SMC7
C	-13	HIS	-	EXPRESSION TAG	UNP Q5SMC7
C	-12	HIS	-	EXPRESSION TAG	UNP Q5SMC7
C	-11	HIS	-	EXPRESSION TAG	UNP Q5SMC7
C	-10	SER	-	EXPRESSION TAG	UNP Q5SMC7
C	-9	GLN	-	EXPRESSION TAG	UNP Q5SMC7
C	-8	ASP	-	EXPRESSION TAG	UNP Q5SMC7
C	-7	PRO	-	EXPRESSION TAG	UNP Q5SMC7
C	-6	LEU	-	EXPRESSION TAG	UNP Q5SMC7
C	-5	VAL	-	EXPRESSION TAG	UNP Q5SMC7
C	-4	PRO	-	EXPRESSION TAG	UNP Q5SMC7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	ARG	-	EXPRESSION TAG	UNP Q5SMC7
C	-2	GLY	-	EXPRESSION TAG	UNP Q5SMC7
C	-1	SER	-	EXPRESSION TAG	UNP Q5SMC7
C	0	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-20	MSE	-	EXPRESSION TAG	UNP Q5SMC7
D	-19	GLY	-	EXPRESSION TAG	UNP Q5SMC7
D	-18	SER	-	EXPRESSION TAG	UNP Q5SMC7
D	-17	SER	-	EXPRESSION TAG	UNP Q5SMC7
D	-16	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-15	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-14	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-13	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-12	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-11	HIS	-	EXPRESSION TAG	UNP Q5SMC7
D	-10	SER	-	EXPRESSION TAG	UNP Q5SMC7
D	-9	GLN	-	EXPRESSION TAG	UNP Q5SMC7
D	-8	ASP	-	EXPRESSION TAG	UNP Q5SMC7
D	-7	PRO	-	EXPRESSION TAG	UNP Q5SMC7
D	-6	LEU	-	EXPRESSION TAG	UNP Q5SMC7
D	-5	VAL	-	EXPRESSION TAG	UNP Q5SMC7
D	-4	PRO	-	EXPRESSION TAG	UNP Q5SMC7
D	-3	ARG	-	EXPRESSION TAG	UNP Q5SMC7
D	-2	GLY	-	EXPRESSION TAG	UNP Q5SMC7
D	-1	SER	-	EXPRESSION TAG	UNP Q5SMC7
D	0	HIS	-	EXPRESSION TAG	UNP Q5SMC7

- Molecule 3 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	C	1	31	17	4	9	1	0	0
			Total	C	N	O	P		
3	D	1	31	17	4	9	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA

Chain A:





- Molecule 1: tRNA

Chain B:





- Molecule 2: tRNA-dihydrouridine synthase

Chain C:



- Molecule 2: tRNA-dihydrouridine synthase

Chain D:


MSE GLY SER SER HIS HIS HIS HIS HIS HIS SER GLN ASP PRO LEU VAL PRO ARG GLY SER HIS
M1 L2 D3 P4
S7 V8 A9
V12
R20 F21 L22 V23 R24 Q25
R31 L32
D39
L43
G92 C93 P94
K97
L107 L108 L109 D110 L111
V114 R115
T130 V131 K132 M133 R134 L135 G136 L137
E141
M153
A156
V160 F161 V162
R166 S167 A168
L172
I180 P181 P182 L183
P196
N203
I206
V222 M223 L224
F233
A238
V242
R251
L262
V266
T270 P271 P272
L276
R286
P287 K288
L291 W292 R293 R294
R310 L311 M312 E313 E314 E315
VAL GLY GLU GLU GLY GLU LYS GLU LYS PRO GLY PRO ARG GLY GLN ARG GLU ALA ALA PRO GLY PRO ALA ARG GLU GLY VAL

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α, β, γ	118.89Å 118.89Å 319.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.60 – 3.51 48.60 – 3.51	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.60-3.51) 99.2 (48.60-3.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$< I/\sigma(I) >$ [1]	3.11 (at 3.48Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.300 , 0.320 0.267 , 0.287	Depositor DCC
R_{free} test set	2323 reflections (7.92%)	wwPDB-VP
Wilson B-factor (Å^2)	102.9	Xtriage
Anisotropy	0.764	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 99.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.40$, $< L^2 > = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.82	EDS
Total number of atoms	8204	wwPDB-VP
Average B, all atoms (Å^2)	148.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, H2U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.29	1/1722 (0.1%)	0.42	0/2682
1	B	0.29	1/1722 (0.1%)	0.42	0/2682
2	C	0.20	0/2546	0.37	0/3416
2	D	0.20	0/2546	0.37	0/3416
All	All	0.24	2/8536 (0.0%)	0.39	0/12196

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	G	OP3-P	-10.61	1.48	1.61
1	A	1	G	OP3-P	-10.58	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1561	0	790	53	0
1	B	1561	0	790	56	0
2	C	2510	0	2586	32	0
2	D	2510	0	2586	42	0
3	C	31	0	19	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	31	0	19	1	0
All	All	8204	0	6790	172	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 172 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:H2U:C5	2:C:93:CYS:SG	2.01	1.47
1:A:43:A:H3'	1:A:44:G:H5"	1.16	1.10
1:A:43:A:H3'	1:A:44:G:C5'	1.82	1.09
1:B:43:A:H3'	1:B:44:G:H5"	1.18	1.08
1:B:43:A:H3'	1:B:44:G:C5'	1.83	1.07

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	313/363 (86%)	297 (95%)	16 (5%)	0	100	100
2	D	313/363 (86%)	297 (95%)	16 (5%)	0	100	100
All	All	626/726 (86%)	594 (95%)	32 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	261/287 (91%)	254 (97%)	7 (3%)	44	73
2	D	261/287 (91%)	256 (98%)	5 (2%)	57	80
All	All	522/574 (91%)	510 (98%)	12 (2%)	50	77

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	107	LEU
2	D	111	LEU
2	D	311	LEU
2	D	294	ARG
2	C	259	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	72/76 (94%)	20 (27%)	1 (1%)
1	B	72/76 (94%)	20 (27%)	1 (1%)
All	All	144/152 (94%)	40 (27%)	2 (1%)

5 of 40 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	10	G
1	A	13	C
1	A	17	U
1	A	18	G
1	A	19	G

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	22	G
1	B	22	G

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	H2U	B	20	1	18,21,22	0.71	0	19,30,33	1.89	3 (15%)
1	H2U	A	20	1	18,21,22	0.72	0	19,30,33	1.92	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	H2U	B	20	1	-	0/7/38/39	0/2/2/2
1	H2U	A	20	1	-	0/7/38/39	0/2/2/2

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	20	H2U	N3-C2-N1	5.43	122.10	116.65
1	B	20	H2U	N3-C2-N1	5.29	121.96	116.65
1	A	20	H2U	C5-C4-N3	4.47	121.45	116.69
1	B	20	H2U	C5-C4-N3	4.44	121.41	116.69
1	B	20	H2U	O2-C2-N1	-3.58	118.80	123.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	20	H2U	3	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	FMN	D	400	-	33,33,33	1.05	2 (6%)	48,50,50	1.24	9 (18%)
3	FMN	C	400	-	33,33,33	1.04	2 (6%)	48,50,50	1.24	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMN	D	400	-	-	1/18/18/18	0/3/3/3
3	FMN	C	400	-	-	1/18/18/18	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	400	FMN	C4A-N5	3.53	1.38	1.30
3	C	400	FMN	C4A-N5	3.43	1.38	1.30
3	D	400	FMN	C10-N1	2.54	1.38	1.33
3	C	400	FMN	C10-N1	2.47	1.38	1.33

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	400	FMN	C4-N3-C2	-3.29	119.80	125.64
3	D	400	FMN	C4-N3-C2	-3.18	119.99	125.64
3	D	400	FMN	C4A-C4-N3	2.64	119.98	113.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	400	FMN	C4A-C4-N3	2.62	119.93	113.25
3	D	400	FMN	C9A-C5A-N5	-2.55	119.75	122.45

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	400	FMN	C4'-C5'-O5'-P
3	C	400	FMN	C4'-C5'-O5'-P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	400	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FMN D 400



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN C 400



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	72/76 (94%)	3.90	57 (79%)	0	0	154, 207, 270, 282	0
1	B	72/76 (94%)	4.30	60 (83%)	0	0	153, 207, 271, 281	0
2	C	305/363 (84%)	0.40	1 (0%)	94	89	79, 110, 133, 163	0
2	D	305/363 (84%)	0.47	5 (1%)	72	59	80, 110, 134, 162	0
All	All	754/878 (85%)	1.13	123 (16%)	1	1	79, 114, 232, 282	0

The worst 5 of 123 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	73	A	13.9
1	A	70	G	12.4
1	B	71	G	12.1
1	A	72	C	11.0
1	B	72	C	9.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	H2U	A	20	20/21	0.90	0.30	114,118,127,129	0
1	H2U	B	20	20/21	0.92	0.31	113,120,126,127	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	FMN	D	400	31/31	0.93	0.36	90,99,105,107	0
3	FMN	C	400	31/31	0.94	0.38	91,100,104,105	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FMN D 400:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around FMN C 400:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.