



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 04:28 am BST

PDB ID : 4B0F  
Title : Heptameric core complex structure of C4b-binding (C4BP) protein from human  
Authors : Schmelz, S.; Hofmeyer, T.; Kolmar, H.; Heinz, D.W.  
Deposited on : 2012-07-02  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	65	
1	B	65	
1	C	65	
1	D	65	
1	E	65	
1	F	65	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	65	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '5%', a large green segment labeled '74%', a yellow segment labeled '11%', and a grey segment at the end labeled '14%'. A small black dot is visible on the grey segment.</p>

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 2941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C4B-BINDING PROTEIN ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	57	436	270	72	90	2	2	0	0	0
1	B	55	424	263	70	87	2	2	0	0	0
1	C	49	383	237	64	78	2	2	0	0	0
1	D	55	430	266	73	87	2	2	0	0	0
1	E	55	415	257	69	85	2	2	0	0	0
1	F	53	411	254	68	85	2	2	0	0	0
1	G	56	436	271	72	89	2	2	0	0	0

There are 49 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	533	SER	-	expression tag	UNP P04003
A	534	ALA	-	expression tag	UNP P04003
A	535	GLY	-	expression tag	UNP P04003
A	536	ALA	-	expression tag	UNP P04003
A	537	HIS	-	expression tag	UNP P04003
A	538	ALA	-	expression tag	UNP P04003
A	539	GLY	-	expression tag	UNP P04003
B	533	SER	-	expression tag	UNP P04003
B	534	ALA	-	expression tag	UNP P04003
B	535	GLY	-	expression tag	UNP P04003
B	536	ALA	-	expression tag	UNP P04003
B	537	HIS	-	expression tag	UNP P04003
B	538	ALA	-	expression tag	UNP P04003
B	539	GLY	-	expression tag	UNP P04003
C	533	SER	-	expression tag	UNP P04003

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	534	ALA	-	expression tag	UNP P04003
C	535	GLY	-	expression tag	UNP P04003
C	536	ALA	-	expression tag	UNP P04003
C	537	HIS	-	expression tag	UNP P04003
C	538	ALA	-	expression tag	UNP P04003
C	539	GLY	-	expression tag	UNP P04003
D	533	SER	-	expression tag	UNP P04003
D	534	ALA	-	expression tag	UNP P04003
D	535	GLY	-	expression tag	UNP P04003
D	536	ALA	-	expression tag	UNP P04003
D	537	HIS	-	expression tag	UNP P04003
D	538	ALA	-	expression tag	UNP P04003
D	539	GLY	-	expression tag	UNP P04003
E	533	SER	-	expression tag	UNP P04003
E	534	ALA	-	expression tag	UNP P04003
E	535	GLY	-	expression tag	UNP P04003
E	536	ALA	-	expression tag	UNP P04003
E	537	HIS	-	expression tag	UNP P04003
E	538	ALA	-	expression tag	UNP P04003
E	539	GLY	-	expression tag	UNP P04003
F	533	SER	-	expression tag	UNP P04003
F	534	ALA	-	expression tag	UNP P04003
F	535	GLY	-	expression tag	UNP P04003
F	536	ALA	-	expression tag	UNP P04003
F	537	HIS	-	expression tag	UNP P04003
F	538	ALA	-	expression tag	UNP P04003
F	539	GLY	-	expression tag	UNP P04003
G	533	SER	-	expression tag	UNP P04003
G	534	ALA	-	expression tag	UNP P04003
G	535	GLY	-	expression tag	UNP P04003
G	536	ALA	-	expression tag	UNP P04003
G	537	HIS	-	expression tag	UNP P04003
G	538	ALA	-	expression tag	UNP P04003
G	539	GLY	-	expression tag	UNP P04003

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0
2	C	1	Total Cl 1 1	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	C	1	Total O 1 1	0	0
3	D	1	Total O 1 1	0	0
3	E	1	Total O 1 1	0	0
3	G	1	Total O 1 1	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN

Chain A: 



- Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN

Chain B: 



- Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN

Chain C: 



- Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN

Chain D: 



- Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN

Chain E: 




- Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN

Chain F: 



● Molecule 1: C4B-BINDING PROTEIN ALPHA CHAIN





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.85Å 73.85Å 204.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.50 – 2.80 46.50 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.50-2.80) 100.0 (46.50-2.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.26 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.212 , 0.248 0.217 , 0.256	Depositor DCC
$R_{free}$ test set	734 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.7	Xtrriage
Anisotropy	0.015	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 56.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2941	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.35	0/437	0.53	0/587
1	B	0.36	0/425	0.54	0/569
1	C	0.34	0/384	0.46	0/512
1	D	0.39	0/431	0.55	0/576
1	E	0.35	0/416	0.49	0/559
1	F	0.36	0/412	0.58	1/551 (0.2%)
1	G	0.54	1/439 (0.2%)	0.73	2/590 (0.3%)
All	All	0.39	1/2944 (0.0%)	0.56	3/3944 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	540	TRP	CD2-CE3	5.45	1.48	1.40

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	540	TRP	CE2-CD2-CG	6.21	112.27	107.30
1	G	540	TRP	CG-CD2-CE3	-5.82	128.66	133.90
1	F	545	GLY	C-N-CA	-5.47	108.02	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	436	0	428	3	0
1	B	424	0	420	4	0
1	C	383	0	383	2	0
1	D	430	0	431	4	0
1	E	415	0	405	2	0
1	F	411	0	407	8	0
1	G	436	0	426	3	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
All	All	2941	0	2900	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:580:GLN:NE2	1:F:584:GLN:OE1	1.99	0.94
1:A:589:ARG:O	1:A:593:LEU:HD13	1.84	0.76
1:B:589:ARG:O	1:B:593:LEU:HD13	1.87	0.74
1:D:594:ASP:O	1:D:597:LEU:HB2	1.90	0.70
1:D:584:GLN:OE1	1:F:580:GLN:NE2	2.29	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	55/65 (85%)	53 (96%)	2 (4%)	0	100	100
1	B	53/65 (82%)	52 (98%)	1 (2%)	0	100	100
1	C	47/65 (72%)	46 (98%)	1 (2%)	0	100	100
1	D	53/65 (82%)	50 (94%)	3 (6%)	0	100	100
1	E	53/65 (82%)	50 (94%)	2 (4%)	1 (2%)	8	26
1	F	51/65 (78%)	49 (96%)	2 (4%)	0	100	100
1	G	54/65 (83%)	52 (96%)	2 (4%)	0	100	100
All	All	366/455 (80%)	352 (96%)	13 (4%)	1 (0%)	41	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	594	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/54 (91%)	48 (98%)	1 (2%)	55	84
1	B	48/54 (89%)	47 (98%)	1 (2%)	53	84
1	C	44/54 (82%)	43 (98%)	1 (2%)	50	82
1	D	49/54 (91%)	44 (90%)	5 (10%)	7	22
1	E	46/54 (85%)	45 (98%)	1 (2%)	52	83
1	F	47/54 (87%)	45 (96%)	2 (4%)	29	62
1	G	49/54 (91%)	46 (94%)	3 (6%)	18	48
All	All	332/378 (88%)	318 (96%)	14 (4%)	30	63

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	593	LEU
1	D	597	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	G	563	GLU
1	D	591	SER
1	F	594	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	55/65 (84%)	0.07	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	43, 70, 119, 143	0
1	B	53/65 (81%)	-0.08	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	43, 72, 112, 143	0
1	C	47/65 (72%)	-0.05	1 (2%) <span style="border: 1px solid gray; padding: 2px;">63</span> <span style="border: 1px solid gray; padding: 2px;">54</span>	41, 67, 111, 146	0
1	D	53/65 (81%)	0.18	5 (9%) <span style="border: 1px solid red; padding: 2px;">8</span> <span style="border: 1px solid red; padding: 2px;">4</span>	41, 72, 125, 144	0
1	E	53/65 (81%)	-0.09	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	43, 72, 130, 150	0
1	F	51/65 (78%)	0.02	3 (5%) <span style="border: 1px solid red; padding: 2px;">22</span> <span style="border: 1px solid red; padding: 2px;">14</span>	42, 71, 141, 147	0
1	G	54/65 (83%)	0.13	3 (5%) <span style="border: 1px solid red; padding: 2px;">24</span> <span style="border: 1px solid red; padding: 2px;">16</span>	42, 71, 139, 181	0
All	All	366/455 (80%)	0.03	12 (3%) <span style="border: 1px solid gray; padding: 2px;">46</span> <span style="border: 1px solid gray; padding: 2px;">36</span>	41, 72, 133, 181	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	592	THR	3.3
1	C	590	GLN	3.2
1	G	588	ALA	3.1
1	G	540	TRP	2.9
1	F	590	GLN	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	B	1001	1/1	0.82	0.28	96,96,96,96	0
2	CL	C	1001	1/1	0.91	0.33	100,100,100,100	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.