



wwPDB EM Validation Summary Report ⓘ

Nov 6, 2022 – 09:40 AM EST

PDB ID : 6B1T
EMDB ID : EMD-7034
Title : Improved cryoEM structure of human adenovirus type 5 with atomic details of minor proteins VI and VII
Authors : Dai, X.H.; Wu, L.; Sun, R.; Zhou, Z.H.
Deposited on : 2017-09-18
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	952	
1	B	952	
1	C	952	
1	D	952	
1	E	952	
1	F	952	
1	G	952	
1	H	952	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	952	
1	J	952	
1	K	952	
1	L	952	
2	M	571	
3	N	585	
4	O	227	
4	P	227	
5	Q	140	
5	R	140	
5	S	140	
5	T	140	
6	U	33	
6	V	33	
6	Y	33	
7	W	11	
8	X	217	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 100134 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hexon protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	908	Total 7286	C 4639	N 1231	O 1380	S 36	0	0
1	B	910	Total 7299	C 4647	N 1233	O 1383	S 36	0	0
1	C	910	Total 7302	C 4650	N 1232	O 1383	S 37	0	0
1	D	906	Total 7273	C 4631	N 1229	O 1378	S 35	0	0
1	E	914	Total 7325	C 4662	N 1237	O 1390	S 36	0	0
1	F	910	Total 7300	C 4647	N 1233	O 1384	S 36	0	0
1	G	909	Total 7293	C 4643	N 1232	O 1382	S 36	0	0
1	H	913	Total 7319	C 4659	N 1236	O 1387	S 37	0	0
1	I	910	Total 7300	C 4647	N 1233	O 1384	S 36	0	0
1	J	913	Total 7318	C 4658	N 1236	O 1388	S 36	0	0
1	K	910	Total 7299	C 4646	N 1233	O 1384	S 36	0	0
1	L	912	Total 7311	C 4654	N 1235	O 1386	S 36	0	0

- Molecule 2 is a protein called Penton protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	M	455	Total 3642	C 2309	N 630	O 691	S 12	0	0

- Molecule 3 is a protein called Pre-hexon-linking protein IIIa.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	N	288	Total	C	N	O	S	0	0
			2236	1385	410	437	4		

- Molecule 4 is a protein called Pre-hexon-linking protein VIII.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	O	180	Total	C	N	O	S	0	0
			1383	870	244	264	5		
4	P	180	Total	C	N	O	S	0	0
			1383	870	244	264	5		

- Molecule 5 is a protein called Hexon-interlacing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	Q	119	Total	C	N	O	S	0	0
			874	541	155	176	2		
5	R	93	Total	C	N	O	S	0	0
			698	432	123	141	2		
5	S	86	Total	C	N	O	S	0	0
			647	397	115	133	2		
5	T	88	Total	C	N	O	S	0	0
			663	409	118	134	2		

- Molecule 6 is a protein called Pre-protein VI.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	U	29	Total	C	N	O	S	0	0
			216	132	42	40	2		
6	V	29	Total	C	N	O	S	0	0
			216	132	42	40	2		
6	Y	28	Total	C	N	O	S	0	0
			208	128	40	38	2		

- Molecule 7 is a protein called Pre-histone-like nucleoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	W	11	Total	C	N	O	S	0	0
			83	55	15	12	1		


- Molecule 8 is a protein called Pre-protein VI.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
8	X	33	260	158	49	53	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hexon protein




- Molecule 1: Hexon protein





• Molecule 1: Hexon protein







• Molecule 1: Hexon protein








• Molecule 1: Hexon protein





• Molecule 1: Hexon protein







• Molecule 1: Hexon protein








• Molecule 1: Hexon protein





• Molecule 1: Hexon protein









• Molecule 1: Hexon protein




• Molecule 1: Hexon protein




● Molecule 1: Hexon protein






● Molecule 2: Penton protein




● Molecule 3: Pre-hexon-linking protein IIIA







• Molecule 4: Pre-hexon-linking protein VIII




• Molecule 4: Pre-hexon-linking protein VIII




• Molecule 5: Hexon-interlacing protein







• Molecule 5: Hexon-interlacing protein




• Molecule 5: Hexon-interlacing protein




• Molecule 5: Hexon-interlacing protein




• Molecule 6: Pre-protein VI




• Molecule 6: Pre-protein VI






• Molecule 6: Pre-protein VI



• Molecule 7: Pre-histone-like nucleoprotein



• Molecule 8: Pre-protein VI



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	53000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	16.748	Depositor
Minimum map value	-11.582	Depositor
Average map value	0.004	Depositor
Map value standard deviation	0.876	Depositor
Recommended contour level	3.0	Depositor
Map size (Å)	1088.0, 1088.0, 1088.0	wwPDB
Map dimensions	1280, 1280, 1280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.85, 0.85, 0.85	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/7481	0.52	0/10170
1	B	0.32	0/7495	0.52	0/10189
1	C	0.33	0/7498	0.52	0/10194
1	D	0.35	0/7468	0.52	0/10153
1	E	0.34	0/7521	0.53	0/10227
1	F	0.35	0/7495	0.53	0/10190
1	G	0.36	0/7488	0.54	0/10180
1	H	0.36	0/7515	0.54	0/10217
1	I	0.35	0/7495	0.53	0/10190
1	J	0.35	0/7514	0.53	0/10217
1	K	0.35	0/7494	0.53	0/10188
1	L	0.34	0/7507	0.53	0/10207
2	M	0.30	0/3734	0.51	0/5091
3	N	0.30	0/2271	0.52	0/3090
4	O	0.37	0/1422	0.52	0/1942
4	P	0.36	0/1422	0.53	0/1942
5	Q	0.27	0/884	0.55	0/1204
5	R	0.26	0/706	0.52	0/962
5	S	0.26	0/653	0.52	0/888
5	T	0.26	0/669	0.49	0/909
6	U	0.39	0/222	0.55	0/298
6	V	0.33	0/222	0.50	0/298
6	Y	0.32	0/214	0.51	0/287
7	W	0.33	0/85	0.40	0/110
8	X	0.33	0/265	0.52	0/359
All	All	0.34	0/102740	0.53	0/139702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7286	0	6998	126	0
1	B	7299	0	7011	103	0
1	C	7302	0	7020	112	0
1	D	7273	0	6984	133	0
1	E	7325	0	7036	117	0
1	F	7300	0	7012	133	0
1	G	7293	0	7004	96	0
1	H	7319	0	7033	92	0
1	I	7300	0	7011	88	0
1	J	7318	0	7029	109	0
1	K	7299	0	7010	114	0
1	L	7311	0	7022	96	0
2	M	3642	0	3562	34	0
3	N	2236	0	2207	15	0
4	O	1383	0	1326	11	0
4	P	1383	0	1326	20	0
5	Q	874	0	890	17	0
5	R	698	0	713	7	0
5	S	647	0	652	6	0
5	T	663	0	676	13	0
6	U	216	0	196	38	0
6	V	216	0	196	44	0
6	Y	208	0	190	37	0
7	W	83	0	84	17	0
8	X	260	0	256	35	0
All	All	100134	0	96444	1256	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 1256 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:GLN:NE2	6:Y:22:TRP:CZ3	1.83	1.43
1:D:790:ARG:HH21	8:X:137:ARG:NE	1.25	1.34

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:GLN:NE2	6:Y:22:TRP:HZ3	1.16	1.31
4:P:203:GLY:O	4:P:208:TYR:CE1	1.82	1.31
1:D:872:ARG:NH2	7:W:23:GLY:CA	1.94	1.30

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	898/952 (94%)	860 (96%)	38 (4%)	0	100	100
1	B	900/952 (94%)	856 (95%)	43 (5%)	1 (0%)	51	83
1	C	900/952 (94%)	859 (95%)	40 (4%)	1 (0%)	51	83
1	D	896/952 (94%)	856 (96%)	40 (4%)	0	100	100
1	E	904/952 (95%)	866 (96%)	38 (4%)	0	100	100
1	F	900/952 (94%)	865 (96%)	34 (4%)	1 (0%)	51	83
1	G	899/952 (94%)	859 (96%)	40 (4%)	0	100	100
1	H	903/952 (95%)	867 (96%)	36 (4%)	0	100	100
1	I	900/952 (94%)	865 (96%)	35 (4%)	0	100	100
1	J	903/952 (95%)	869 (96%)	34 (4%)	0	100	100
1	K	900/952 (94%)	864 (96%)	36 (4%)	0	100	100
1	L	902/952 (95%)	867 (96%)	35 (4%)	0	100	100
2	M	451/571 (79%)	426 (94%)	25 (6%)	0	100	100
3	N	284/585 (48%)	276 (97%)	8 (3%)	0	100	100
4	O	176/227 (78%)	171 (97%)	5 (3%)	0	100	100
4	P	176/227 (78%)	171 (97%)	4 (2%)	1 (1%)	25	64
5	Q	115/140 (82%)	109 (95%)	6 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	R	89/140 (64%)	86 (97%)	3 (3%)	0	100	100
5	S	82/140 (59%)	79 (96%)	3 (4%)	0	100	100
5	T	84/140 (60%)	81 (96%)	3 (4%)	0	100	100
6	U	27/33 (82%)	27 (100%)	0	0	100	100
6	V	27/33 (82%)	27 (100%)	0	0	100	100
6	Y	26/33 (79%)	24 (92%)	2 (8%)	0	100	100
7	W	9/11 (82%)	9 (100%)	0	0	100	100
8	X	29/217 (13%)	28 (97%)	1 (3%)	0	100	100
All	All	12380/13921 (89%)	11867 (96%)	509 (4%)	4 (0%)	100	100

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	P	204	PRO
1	C	668	PRO
1	F	668	PRO
1	B	668	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	793/829 (96%)	793 (100%)	0	100	100
1	B	795/829 (96%)	794 (100%)	1 (0%)	93	98
1	C	796/829 (96%)	796 (100%)	0	100	100
1	D	792/829 (96%)	792 (100%)	0	100	100
1	E	798/829 (96%)	798 (100%)	0	100	100
1	F	795/829 (96%)	795 (100%)	0	100	100
1	G	794/829 (96%)	794 (100%)	0	100	100
1	H	797/829 (96%)	796 (100%)	1 (0%)	93	98

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	795/829 (96%)	795 (100%)	0	100	100
1	J	797/829 (96%)	797 (100%)	0	100	100
1	K	795/829 (96%)	795 (100%)	0	100	100
1	L	796/829 (96%)	796 (100%)	0	100	100
2	M	412/489 (84%)	412 (100%)	0	100	100
3	N	241/500 (48%)	241 (100%)	0	100	100
4	O	147/186 (79%)	147 (100%)	0	100	100
4	P	147/186 (79%)	147 (100%)	0	100	100
5	Q	97/112 (87%)	97 (100%)	0	100	100
5	R	81/112 (72%)	81 (100%)	0	100	100
5	S	75/112 (67%)	75 (100%)	0	100	100
5	T	76/112 (68%)	76 (100%)	0	100	100
6	U	22/26 (85%)	22 (100%)	0	100	100
6	V	22/26 (85%)	22 (100%)	0	100	100
6	Y	21/26 (81%)	21 (100%)	0	100	100
7	W	8/8 (100%)	8 (100%)	0	100	100
8	X	30/184 (16%)	30 (100%)	0	100	100
All	All	10922/12027 (91%)	10920 (100%)	2 (0%)	100	100

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	896	ASN
1	H	414	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 131 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	471	ASN
1	L	840	GLN
6	U	5	ASN
1	E	840	GLN
1	E	799	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-7034. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 617



Y Index: 329



Z Index: 1017


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 18707 nm^3 ; this corresponds to an approximate mass of 16898 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8 Fourier-Shell correlation


This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-7034 and PDB model 6B1T. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)




9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3.0).





















































9.4 Atom inclusion [i](#)



At the recommended contour level, 70% of all backbone atoms, 62% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (3.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6249	 0.5520
A	 0.5994	 0.5510
B	 0.5757	 0.5450
C	 0.5903	 0.5470
D	 0.6655	 0.5620
E	 0.6390	 0.5570
F	 0.6468	 0.5590
G	 0.6732	 0.5640
H	 0.6755	 0.5640
I	 0.6766	 0.5620
J	 0.6474	 0.5590
K	 0.6453	 0.5580
L	 0.6512	 0.5610
M	 0.4688	 0.5150
N	 0.5257	 0.5100
O	 0.6832	 0.5510
P	 0.7135	 0.5560
Q	 0.3965	 0.4600
R	 0.4217	 0.4780
S	 0.3370	 0.4600
T	 0.3426	 0.4700
U	 0.6019	 0.5750
V	 0.6445	 0.5760
W	 0.6667	 0.5800
X	 0.3160	 0.5320
Y	 0.6798	 0.5610

