



# wwPDB X-ray Structure Validation Summary Report

Aug 16, 2023 – 11:34 PM EDT

PDB ID : 2B2X  
Title : VLA1 RdeltaH I-domain complexed with a quadruple mutant of the AQC2 Fab  
Authors : Clark, L.A.; Boriack-Sjodin, P.A.; Eldredge, J.; Fitch, C.; Friedman, B.; Hanf, K.J.; Jarpe, M.; Liparoto, S.F.; Li, Y.; Lugovskoy, A.  
Deposited on : 2005-09-19  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	 4% 48% 33% 16%
1	B	223	 6% 48% 29% 16%
2	H	226	 72% 21% 7%
2	I	226	 72% 18% 7%
3	L	213	 2% 67% 28% 3%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	M	213	 <p>2% 66% 31% ..</p>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	188	1490	940	258	288	4	0	0	0
1	B	176	1397	884	240	270	3	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	125	GLY	-	cloning artifact	UNP P18614
A	126	SER	-	cloning artifact	UNP P18614
A	217	VAL	GLY	engineered mutation	UNP P18614
A	218	GLN	ARG	engineered mutation	UNP P18614
A	219	ARG	GLN	engineered mutation	UNP P18614
A	222	ARG	LEU	engineered mutation	UNP P18614
A	341	LEU	-	cloning artifact	UNP P18614
A	342	GLU	-	cloning artifact	UNP P18614
A	343	ARG	-	cloning artifact	UNP P18614
A	344	PRO	-	cloning artifact	UNP P18614
A	345	HIS	-	cloning artifact	UNP P18614
A	346	ARG	-	cloning artifact	UNP P18614
A	347	ASP	-	cloning artifact	UNP P18614
B	125	GLY	-	cloning artifact	UNP P18614
B	126	SER	-	cloning artifact	UNP P18614
B	217	VAL	GLY	engineered mutation	UNP P18614
B	218	GLN	ARG	engineered mutation	UNP P18614
B	219	ARG	GLN	engineered mutation	UNP P18614
B	222	ARG	LEU	engineered mutation	UNP P18614
B	341	LEU	-	cloning artifact	UNP P18614
B	342	GLU	-	cloning artifact	UNP P18614
B	343	ARG	-	cloning artifact	UNP P18614
B	344	PRO	-	cloning artifact	UNP P18614
B	345	HIS	-	cloning artifact	UNP P18614
B	346	ARG	-	cloning artifact	UNP P18614

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	347	ASP	-	cloning artifact	UNP P18614

- Molecule 2 is a protein called Antibody AQC2 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	210	Total	C	N	O	S	0	0	0
			1575	1000	260	308	7			
2	I	210	Total	C	N	O	S	0	0	0
			1575	1000	260	308	7			

- Molecule 3 is a protein called Antibody AQC2 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	210	Total	C	N	O	S	0	0	0
			1636	1026	274	330	6			
3	M	210	Total	C	N	O	S	0	0	0
			1636	1026	274	330	6			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	13	Total	O	0	0
			13	13		
5	H	73	Total	O	0	0
			73	73		
5	L	30	Total	O	0	0
			30	30		
5	B	5	Total	O	0	0
			5	5		
5	I	77	Total	O	0	0
			77	77		
5	M	28	Total	O	0	0
			28	28		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrin alpha-1



- Molecule 1: Integrin alpha-1




- Molecule 2: Antibody AQC2 Fab




- Molecule 2: Antibody AQC2 Fab

Chain I:  72% 18% 7%




- Molecule 3: Antibody AQC2 Fab

Chain L:  2% 67% 28%



- Molecule 3: Antibody AQC2 Fab

Chain M:  2% 66% 31%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.12Å 43.68Å 153.88Å 90.00° 104.10° 90.00°	Depositor
Resolution (Å)	35.00 – 2.20 49.75 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.3 (35.00-2.20) 94.2 (49.75-2.20)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.04 (at 2.20Å)	Xtrriage
Refinement program	CNS, CNX	Depositor
R, $R_{free}$	0.238 , 0.272 0.238 , 0.272	Depositor DCC
$R_{free}$ test set	3380 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.6	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 43.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9537	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7301e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/1510	0.56	0/2036
1	B	0.27	0/1414	0.55	0/1903
2	H	0.39	0/1616	0.71	0/2208
2	I	0.37	0/1616	0.70	0/2208
3	L	0.35	0/1680	0.63	0/2288
3	M	0.33	0/1680	0.61	0/2288
All	All	0.34	0/9516	0.63	0/12931

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1490	0	1497	67	0
1	B	1397	0	1395	75	0
2	H	1575	0	1537	39	0
2	I	1575	0	1537	38	0
3	L	1636	0	1561	53	0
3	M	1636	0	1561	56	0
4	A	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
5	A	13	0	0	0	0
5	B	5	0	0	0	0
5	H	73	0	0	0	0
5	I	77	0	0	1	0
5	L	30	0	0	1	0
5	M	28	0	0	0	0
All	All	9537	0	9088	315	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 315 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:191:GLU:HA	3:L:215:ARG:HH12	1.25	0.98
1:B:234:ARG:HH11	1:B:234:ARG:HB3	1.27	0.97
2:H:161:SER:H	2:H:201:ASN:HD21	1.00	0.95
2:I:161:SER:H	2:I:201:ASN:HD21	1.08	0.94
3:M:6:GLN:H	3:M:99:GLN:NE2	1.64	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	184/223 (82%)	164 (89%)	18 (10%)	2 (1%)	14 12
1	B	166/223 (74%)	147 (89%)	17 (10%)	2 (1%)	13 10
2	H	206/226 (91%)	201 (98%)	5 (2%)	0	100 100
2	I	206/226 (91%)	202 (98%)	4 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	L	208/213 (98%)	198 (95%)	9 (4%)	1 (0%)	29	31
3	M	208/213 (98%)	195 (94%)	12 (6%)	1 (0%)	29	31
All	All	1178/1324 (89%)	1107 (94%)	65 (6%)	6 (0%)	29	31

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	287	GLY
1	B	236	GLU
1	A	333	GLU
3	L	83	ALA
1	B	173	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/194 (84%)	157 (96%)	7 (4%)	29	36
1	B	153/194 (79%)	148 (97%)	5 (3%)	38	49
2	H	177/190 (93%)	170 (96%)	7 (4%)	31	40
2	I	177/190 (93%)	166 (94%)	11 (6%)	18	21
3	L	187/190 (98%)	179 (96%)	8 (4%)	29	36
3	M	187/190 (98%)	179 (96%)	8 (4%)	29	36
All	All	1045/1148 (91%)	999 (96%)	46 (4%)	28	35

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	38	ARG
2	I	154	PRO
2	I	43	LYS
2	I	113	LEU
2	I	164	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	288	HIS
3	M	36	GLN
1	B	314	HIS
2	I	169	HIS
3	M	99	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	188/223 (84%)	0.40	8 (4%) 35 33	28, 59, 77, 83	0
1	B	176/223 (78%)	0.66	14 (7%) 12 11	33, 66, 91, 96	0
2	H	210/226 (92%)	0.06	0 100 100	22, 31, 49, 58	0
2	I	210/226 (92%)	0.01	0 100 100	24, 34, 50, 61	0
3	L	210/213 (98%)	0.17	4 (1%) 66 65	22, 42, 74, 86	0
3	M	210/213 (98%)	0.16	5 (2%) 59 56	24, 41, 68, 86	0
All	All	1204/1324 (90%)	0.23	31 (2%) 56 53	22, 42, 78, 96	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	200	LEU	4.3
1	B	179	GLY	3.7
1	B	147	LEU	3.4
3	M	192	ARG	3.4
1	B	334	ARG	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	B	401	1/1	0.88	0.16	54,54,54,54	0
4	MG	A	400	1/1	0.92	0.14	47,47,47,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.