



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 04:38 am GMT

PDB ID : 4B5S  
Title : Crystal structures of divalent metal dependent pyruvate aldolase, HpaI, in complex with pyruvate  
Authors : Coincon, M.; Wang, W.; Seah, S.Y.K.; Sygusch, J.  
Deposited on : 2012-08-07  
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	 2% 92% 8%
1	B	251	 2% 94% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3GR	A	1252[B]	X	-	-	-
4	PO4	A	1254	-	-	X	-
7	GOL	B	1252[A]	-	-	-	X
7	GOL	B	1252[B]	-	-	-	X

## 2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 8843 atoms, of which 3864 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-HYDROXY-2-OXO-HEPTANE-1,7-DIOATE ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	251	Total 3838	C 1212	H 1927	N 337	O 356	S 6	0	1	0
1	B	251	Total 3852	C 1216	H 1935	N 338	O 357	S 6	0	2	0

- Molecule 2 is D-Glyceraldehyde (three-letter code: 3GR) (formula: C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	Total 14	C 6	H 2	O 6	0	1

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Co 1	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Co	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is PYRUVIC ACID (three-letter code: PYR) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mg 1 1	0	0
6	B	1	Total Mg 1 1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 12 6 6	0	1

- Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	530	Total O 530 530	0	0
8	B	561	Total O 561 561	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-HYDROXY-2-OXO-HEPTANE-1,7-DIOATE ALDOLASE

Chain A: 



- Molecule 1: 4-HYDROXY-2-OXO-HEPTANE-1,7-DIOATE ALDOLASE

Chain B: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	151.74Å 151.74Å 151.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.55 – 1.68 47.98 – 1.68	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.55-1.68) 96.0 (47.98-1.68)	Depositor EDS
$R_{merge}$	0.02	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 1.68Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.151 , 0.179 0.154 , 0.180	Depositor DCC
$R_{free}$ test set	9873 reflections (7.49%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.2	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 66.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.029 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	8843	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO, PYR, PO4, GOL, MG, 3GR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.83	0/1952	0.79	0/2661
1	B	0.88	0/1961	0.86	1/2673 (0.0%)
All	All	0.86	0/3913	0.83	1/5334 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	130	ARG	NE-CZ-NH2	-5.89	117.35	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1911	1927	1929	18	0
1	B	1917	1935	1937	8	0
2	A	12	2	12	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	0	2	0
4	B	10	0	0	0	0
5	A	6	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	6	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	B	12	0	16	2	0
8	A	530	0	0	10	5
8	B	561	0	0	2	7
All	All	4979	3864	3894	28	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:SER:O	8:A:2351:HOH:O	1.90	0.89
4:A:1254:PO4:O2	8:A:2434:HOH:O	1.90	0.88
1:B:26:TYR:HB2	7:B:1252[A]:GOL:H32	1.63	0.78
4:A:1254:PO4:O1	8:A:2527:HOH:O	2.03	0.76
1:A:2:GLU:OE1	8:A:2018:HOH:O	2.06	0.73
1:A:23:SER:HB2	7:B:1252[B]:GOL:H32	1.70	0.72
1:B:121:ALA:O	8:B:2334:HOH:O	2.09	0.70
1:A:123:ALA:HB3	8:A:2351:HOH:O	1.92	0.69
1:B:73[B]:TRP:CE2	1:B:75:ASP:HB2	2.31	0.66
1:A:120:SER:HA	8:A:2351:HOH:O	1.96	0.65
1:A:73[B]:TRP:CE2	1:A:75:ASP:HB2	2.31	0.65
1:A:75:ASP:OD2	8:A:2225:HOH:O	2.13	0.65
1:A:26:TYR:HB2	2:A:1252[A]:3GR:H1C1	1.81	0.61
1:B:51[A]:GLN:NE2	8:B:2163:HOH:O	2.32	0.60
1:A:121:ALA:O	8:A:2354:HOH:O	2.18	0.53
1:A:120:SER:CA	8:A:2351:HOH:O	2.56	0.52
1:A:158:PRO:HG3	1:A:201:GLN:HE22	1.75	0.52
1:A:221:ARG:HG3	8:A:2490:HOH:O	2.11	0.51
1:B:171:ILE:O	1:B:173:PRO:HD3	2.16	0.46
1:A:133:ASP:HB2	1:A:137:LYS:HE3	1.98	0.45
1:B:30:LEU:C	1:B:30:LEU:HD23	2.38	0.44
1:A:122:LEU:HD23	1:A:122:LEU:HA	1.83	0.44
1:A:84:ASP:O	1:A:123:ALA:HB1	2.19	0.42
1:A:30:LEU:C	1:A:30:LEU:HD23	2.41	0.41
1:A:79:ILE:O	1:A:83:LEU:HG	2.21	0.41
1:B:133:ASP:HB2	1:B:137:LYS:HE3	2.03	0.41
1:A:213:ILE:HG23	1:A:213:ILE:O	2.21	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:79:ILE:O	1:B:83:LEU:HG	2.22	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:2178:HOH:O	8:B:2168:HOH:O[5_555]	1.93	0.27
8:A:2166:HOH:O	8:B:2168:HOH:O[5_555]	2.09	0.11
8:A:2095:HOH:O	8:B:2163:HOH:O[5_555]	2.11	0.09
8:B:2225:HOH:O	8:B:2234:HOH:O[5_555]	2.12	0.08
8:A:2149:HOH:O	8:B:2202:HOH:O[5_555]	2.13	0.07
8:A:2154:HOH:O	8:B:2109:HOH:O[5_555]	2.13	0.07
8:B:2154:HOH:O	8:B:2162:HOH:O[5_555]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/251 (100%)	246 (98%)	4 (2%)	0	100	100
1	B	251/251 (100%)	248 (99%)	3 (1%)	0	100	100
All	All	501/502 (100%)	494 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/195 (100%)	196 (100%)	0	100	100
1	B	197/195 (101%)	194 (98%)	3 (2%)	65	48
All	All	393/390 (101%)	390 (99%)	3 (1%)	81	72

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	122	LEU
1	B	185	ASN
1	B	186	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	201	GLN
1	B	185	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GOL	B	1252[B]	-	5,5,5	0.26	0	5,5,5	0.27	0
4	PO4	B	1255	-	4,4,4	0.77	0	6,6,6	0.68	0
2	3GR	A	1252[A]	-	4,5,5	3.07	1 (25%)	4,5,5	1.06	1 (25%)
7	GOL	B	1252[A]	-	5,5,5	0.25	0	5,5,5	0.27	0
4	PO4	A	1255	-	4,4,4	0.89	0	6,6,6	0.42	0
5	PYR	B	1256	6,3	5,5,5	2.76	3 (60%)	3,6,6	1.54	1 (33%)
5	PYR	A	1256	6,3	5,5,5	2.77	3 (60%)	3,6,6	1.57	1 (33%)
4	PO4	B	1254	-	4,4,4	0.81	0	6,6,6	0.68	0
4	PO4	A	1254	-	4,4,4	0.95	0	6,6,6	0.49	0
2	3GR	A	1252[B]	-	4,5,5	3.06	1 (25%)	4,5,5	1.05	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	B	1252[B]	-	-	3/4/4/4	-
2	3GR	A	1252[A]	-	-	3/3/4/4	-
7	GOL	B	1252[A]	-	-	0/4/4/4	-
5	PYR	B	1256	6,3	-	0/4/4/4	-
5	PYR	A	1256	6,3	-	0/4/4/4	-
2	3GR	A	1252[B]	-	1/1/1/2	3/3/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1252[A]	3GR	O1-C1	5.94	1.43	1.19
2	A	1252[B]	3GR	O1-C1	5.93	1.43	1.19
5	B	1256	PYR	CA-C	-4.13	1.39	1.54
5	A	1256	PYR	CA-C	-3.97	1.40	1.54
5	A	1256	PYR	O3-CA	3.51	1.31	1.23
5	B	1256	PYR	O3-CA	3.47	1.30	1.23
5	A	1256	PYR	O-C	3.16	1.31	1.22
5	B	1256	PYR	O-C	2.97	1.30	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1256	PYR	OXT-C-CA	2.28	120.20	113.97
5	B	1256	PYR	OXT-C-CA	2.06	119.60	113.97
2	A	1252[A]	3GR	O3-C3-C2	-2.02	109.46	112.40
2	A	1252[B]	3GR	O3-C3-C2	-2.01	109.47	112.40

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1252[B]	3GR	C2

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1252[A]	3GR	O2-C2-C3-O3
2	A	1252[A]	3GR	O1-C1-C2-C3
2	A	1252[B]	3GR	O2-C2-C3-O3
2	A	1252[B]	3GR	C1-C2-C3-O3
2	A	1252[B]	3GR	O1-C1-C2-C3
2	A	1252[A]	3GR	C1-C2-C3-O3
7	B	1252[B]	GOL	O2-C2-C3-O3
7	B	1252[B]	GOL	C1-C2-C3-O3
7	B	1252[B]	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	1252[B]	GOL	1	0
2	A	1252[A]	3GR	1	0
7	B	1252[A]	GOL	1	0
4	A	1254	PO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	251/251 (100%)	-0.19	4 (1%) 72 75	15, 26, 48, 75	0
1	B	251/251 (100%)	-0.26	4 (1%) 72 75	15, 24, 46, 81	0
All	All	502/502 (100%)	-0.22	8 (1%) 72 75	15, 25, 48, 81	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	51[A]	GLN	4.2
1	B	73[A]	TRP	4.2
1	A	184	GLY	3.7
1	A	73[A]	TRP	3.2
1	B	132	PRO	2.7
1	B	184	GLY	2.5
1	A	132	PRO	2.1
1	A	50	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	GOL	B	1252[A]	6/6	0.53	0.43	24,26,36,45	6
7	GOL	B	1252[B]	6/6	0.53	0.43	29,33,36,37	6
2	3GR	A	1252[A]	6/6	0.61	0.40	23,32,42,51	7
2	3GR	A	1252[B]	6/6	0.61	0.40	28,30,38,43	7
5	PYR	B	1256	6/6	0.82	0.12	32,36,46,47	0
4	PO4	A	1255	5/5	0.88	0.21	50,50,61,66	5
4	PO4	B	1255	5/5	0.91	0.26	70,76,97,108	0
4	PO4	A	1254	5/5	0.92	0.13	36,40,51,55	5
5	PYR	A	1256	6/6	0.93	0.09	32,36,42,54	0
4	PO4	B	1254	5/5	0.93	0.15	42,45,53,57	5
3	CO	B	1253	1/1	0.96	0.05	31,31,31,31	1
6	MG	B	1257	1/1	0.96	0.05	22,22,22,22	1
6	MG	A	1257	1/1	0.97	0.05	55,55,55,55	1
3	CO	A	1253	1/1	0.97	0.05	21,21,21,21	1

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.