



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2023 – 02:48 PM JST

PDB ID : 5B7Y
Title : Crystal Structure of Hyperthermophilic *Thermotoga maritima* L-Ketose-3-Epimerase with Co₂⁺
Authors : Cao, T.P.; Shin, S.M.; Lee, D.W.; Lee, S.H.
Deposited on : 2016-06-10
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1611 (1.34-1.30)
Clashscore	141614	1667 (1.34-1.30)
Ramachandran outliers	138981	1615 (1.34-1.30)
Sidechain outliers	138945	1615 (1.34-1.30)
RSRZ outliers	127900	1580 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	
1	B	290	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	1PG	A	302	-	-	X	-

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein TM_0416.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	269	Total	C	N	O	S	0	0	0
			2136	1361	370	396	9			
1	B	268	Total	C	N	O	S	0	0	0
			2128	1355	369	395	9			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q9WYP7
A	-18	GLY	-	expression tag	UNP Q9WYP7
A	-17	SER	-	expression tag	UNP Q9WYP7
A	-16	SER	-	expression tag	UNP Q9WYP7
A	-15	HIS	-	expression tag	UNP Q9WYP7
A	-14	HIS	-	expression tag	UNP Q9WYP7
A	-13	HIS	-	expression tag	UNP Q9WYP7
A	-12	HIS	-	expression tag	UNP Q9WYP7
A	-11	HIS	-	expression tag	UNP Q9WYP7
A	-10	HIS	-	expression tag	UNP Q9WYP7
A	-9	SER	-	expression tag	UNP Q9WYP7
A	-8	SER	-	expression tag	UNP Q9WYP7
A	-7	GLY	-	expression tag	UNP Q9WYP7
A	-6	LEU	-	expression tag	UNP Q9WYP7
A	-5	VAL	-	expression tag	UNP Q9WYP7
A	-4	PRO	-	expression tag	UNP Q9WYP7
A	-3	ARG	-	expression tag	UNP Q9WYP7
A	-2	GLY	-	expression tag	UNP Q9WYP7
A	-1	SER	-	expression tag	UNP Q9WYP7
A	0	HIS	-	expression tag	UNP Q9WYP7
B	-19	MET	-	expression tag	UNP Q9WYP7
B	-18	GLY	-	expression tag	UNP Q9WYP7
B	-17	SER	-	expression tag	UNP Q9WYP7
B	-16	SER	-	expression tag	UNP Q9WYP7
B	-15	HIS	-	expression tag	UNP Q9WYP7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP Q9WYP7
B	-13	HIS	-	expression tag	UNP Q9WYP7
B	-12	HIS	-	expression tag	UNP Q9WYP7
B	-11	HIS	-	expression tag	UNP Q9WYP7
B	-10	HIS	-	expression tag	UNP Q9WYP7
B	-9	SER	-	expression tag	UNP Q9WYP7
B	-8	SER	-	expression tag	UNP Q9WYP7
B	-7	GLY	-	expression tag	UNP Q9WYP7
B	-6	LEU	-	expression tag	UNP Q9WYP7
B	-5	VAL	-	expression tag	UNP Q9WYP7
B	-4	PRO	-	expression tag	UNP Q9WYP7
B	-3	ARG	-	expression tag	UNP Q9WYP7
B	-2	GLY	-	expression tag	UNP Q9WYP7
B	-1	SER	-	expression tag	UNP Q9WYP7
B	0	HIS	-	expression tag	UNP Q9WYP7

- Molecule 2 is 2-(2-{2-[2-(2-METHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHOXY)-ETHANOL (three-letter code: 1PG) (formula: C₁₁H₂₄O₆).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C O	0	0
			17	11 6		
2	A	1	Total	C O	0	0
			17	11 6		
2	B	1	Total	C O	0	0
			17	11 6		

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Co 1 1	0	0
3	B	1	Total Co 1 1	0	0

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 6 2	0	0
4	A	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	349	Total O 349 349	0	0

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	327	Total 327	O 327	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uncharacterized protein TM_0416

Chain A: 



- Molecule 1: Uncharacterized protein TM_0416

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.28Å 55.26Å 58.75Å 107.14° 102.35° 92.20°	Depositor
Resolution (Å)	35.15 – 1.32 35.15 – 1.32	Depositor EDS
% Data completeness (in resolution range)	91.9 (35.15-1.32) 91.9 (35.15-1.32)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.87 (at 1.32Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.171 , 0.189 0.171 , 0.188	Depositor DCC
R_{free} test set	6405 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtrriage
Anisotropy	0.013	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5033	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO, MPD, 1PG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2173	0.57	1/2930 (0.0%)
1	B	0.33	0/2165	0.54	0/2919
All	All	0.34	0/4338	0.55	1/5849 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	13	ALA	C-N-CA	6.05	136.83	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2136	0	2170	43	1
1	B	2128	0	2159	19	0
2	A	34	0	48	20	1
2	B	17	0	24	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	0	28	1	0
4	B	24	0	42	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	349	0	0	9	0
5	B	327	0	0	6	1
All	All	5033	0	4471	68	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263:LYS:HZ2	2:A:302:1PG:H82	1.14	1.10
1:A:263:LYS:NZ	2:A:302:1PG:H102	1.74	1.01
1:A:263:LYS:HZ2	2:A:302:1PG:H102	1.24	0.98
1:A:263:LYS:NZ	2:A:302:1PG:H82	1.81	0.93
1:B:252:GLU:HG3	5:B:573:HOH:O	1.69	0.91
1:B:232:LYS:NZ	1:B:268:LYS:O	2.03	0.90
1:A:136:ARG:NH2	5:A:403:HOH:O	2.12	0.83
1:A:263:LYS:HZ3	2:A:302:1PG:H61	1.41	0.82
1:A:90:LYS:NZ	5:A:401:HOH:O	2.05	0.78
2:B:301:1PG:O6	2:B:301:1PG:H13	1.84	0.77
1:A:263:LYS:CE	2:A:302:1PG:H102	2.15	0.76
1:B:264:GLU:HA	1:B:267:ILE:HD12	1.68	0.76
1:A:32:ARG:NH1	5:A:405:HOH:O	2.19	0.74
1:B:145:LYS:HZ1	2:B:301:1PG:H22	1.56	0.70
1:A:263:LYS:HZ2	2:A:302:1PG:C10	2.01	0.69
1:B:145:LYS:HZ3	2:B:301:1PG:H41	1.56	0.69
1:A:263:LYS:HZ3	2:A:302:1PG:H42	1.57	0.69
1:A:263:LYS:HE2	2:A:302:1PG:H21	1.74	0.68
1:A:86:GLU:OE2	5:A:402:HOH:O	2.11	0.68
1:A:127:GLU:OE1	1:A:167:ARG:NH2	2.28	0.67
1:A:263:LYS:HE2	2:A:302:1PG:C2	2.25	0.66
4:B:305:MPD:O2	4:B:305:MPD:O4	2.14	0.66
1:A:13:ALA:N	1:A:14:PHE:HB3	2.12	0.65
1:B:145:LYS:NZ	2:B:301:1PG:H41	2.11	0.65
1:A:263:LYS:HE3	2:A:302:1PG:H102	1.79	0.64
1:A:237:ASN:OD1	5:A:404:HOH:O	2.16	0.63
1:A:225:ARG:NH2	5:A:409:HOH:O	2.32	0.63
1:A:266:ILE:O	1:A:269:LEU:HB2	1.98	0.63
1:A:263:LYS:NZ	2:A:302:1PG:H61	2.14	0.62
1:A:266:ILE:HA	1:A:269:LEU:HD22	1.85	0.58
1:A:263:LYS:HZ2	2:A:302:1PG:C8	2.01	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:ASP:HB3	1:A:14:PHE:CD2	2.39	0.57
1:B:123:TYR:OH	1:B:164:ASP:OD2	2.18	0.57
1:A:263:LYS:NZ	2:A:302:1PG:H42	2.19	0.56
1:A:13:ALA:H	1:A:14:PHE:HB3	1.70	0.55
1:B:145:LYS:HZ1	2:B:301:1PG:C2	2.21	0.53
1:B:263:LYS:O	1:B:267:ILE:HG13	2.08	0.53
1:A:11:ASP:HB3	1:A:14:PHE:HD2	1.74	0.52
1:A:25:LYS:NZ	1:A:252:GLU:OE2	2.43	0.50
4:B:304:MPD:O2	5:B:402:HOH:O	2.19	0.50
1:A:225:ARG:CZ	5:A:409:HOH:O	2.59	0.50
1:A:263:LYS:NZ	2:A:302:1PG:H22	2.27	0.49
1:A:13:ALA:O	5:A:406:HOH:O	2.20	0.49
1:B:52:GLU:O	1:B:55:ILE:HG13	2.13	0.49
1:A:82:HIS:CE1	1:A:87:ILE:HD12	2.48	0.49
1:A:100:GLU:HB2	4:A:305:MPD:H52	1.94	0.49
1:B:19:PHE:HD2	1:B:20:LYS:HD2	1.79	0.48
1:B:267:ILE:HG22	1:B:268:LYS:HG2	1.97	0.47
1:A:267:ILE:HD12	2:A:302:1PG:H101	1.97	0.46
4:B:303:MPD:HM1	5:B:628:HOH:O	2.15	0.46
1:B:263:LYS:NZ	5:B:403:HOH:O	2.22	0.46
4:B:304:MPD:O2	4:B:304:MPD:O4	2.29	0.46
1:B:90:LYS:NZ	5:B:401:HOH:O	2.05	0.45
1:B:56:LEU:HA	1:B:59:GLU:HG2	1.99	0.44
1:A:13:ALA:HB3	1:A:14:PHE:CB	2.47	0.44
1:A:263:LYS:HZ1	2:A:302:1PG:H22	1.82	0.44
2:A:302:1PG:H42	2:A:302:1PG:H22	1.79	0.44
1:B:180:LEU:HD13	1:B:206:HIS:HB3	1.99	0.44
1:B:145:LYS:NZ	2:B:301:1PG:H22	2.30	0.43
1:A:169:LEU:HD11	1:A:179:ILE:HG13	2.01	0.43
1:A:136:ARG:HD3	5:A:672:HOH:O	2.19	0.43
1:A:263:LYS:HZ1	2:A:302:1PG:H13	1.85	0.42
1:A:145:LYS:NZ	2:A:301:1PG:H51	2.34	0.42
1:B:54:LYS:O	1:B:58:GLU:HG3	2.21	0.41
1:A:269:LEU:HD12	1:A:269:LEU:HA	1.87	0.41
1:A:264:GLU:OE2	1:A:268:LYS:NZ	2.50	0.41
1:B:32:ARG:NH2	5:B:411:HOH:O	2.42	0.40
1:A:13:ALA:CA	1:A:14:PHE:HB3	2.51	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:GLU:OE2	2:A:302:1PG:O6[1_455]	1.99	0.21
5:B:625:HOH:O	5:B:722:HOH:O[1_545]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/290 (92%)	263 (98%)	2 (1%)	2 (1%)	22	3
1	B	266/290 (92%)	263 (99%)	2 (1%)	1 (0%)	34	10
All	All	533/580 (92%)	526 (99%)	4 (1%)	3 (1%)	25	5

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	PHE
1	A	159	ILE
1	B	159	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/247 (93%)	229 (100%)	0	100	100
1	B	228/247 (92%)	227 (100%)	1 (0%)	91	76
All	All	457/494 (92%)	456 (100%)	1 (0%)	93	79

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	268	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1PG	B	301	-	16,16,16	0.74	0	15,15,15	1.38	2 (13%)
4	MPD	B	303	-	7,7,7	0.34	0	9,10,10	0.72	0
4	MPD	B	305	-	7,7,7	0.27	0	9,10,10	0.17	0
4	MPD	A	304	-	7,7,7	0.24	0	9,10,10	0.39	0
4	MPD	A	305	-	7,7,7	0.21	0	9,10,10	0.19	0
2	1PG	A	301	-	16,16,16	0.75	0	15,15,15	0.22	0
2	1PG	A	302	-	16,16,16	0.37	0	15,15,15	0.96	1 (6%)
4	MPD	B	304	-	7,7,7	0.30	0	9,10,10	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PG	B	301	-	-	4/14/14/14	-
4	MPD	B	303	-	-	0/5/5/5	-
4	MPD	B	305	-	-	5/5/5/5	-
4	MPD	A	304	-	-	0/5/5/5	-
4	MPD	A	305	-	-	3/5/5/5	-
2	1PG	A	301	-	-	3/14/14/14	-
2	1PG	A	302	-	-	9/14/14/14	-
4	MPD	B	304	-	-	1/5/5/5	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	1PG	O2-C4-C5	3.65	126.87	110.39
2	B	301	1PG	C6-O3-C5	3.19	127.11	113.29
2	A	302	1PG	O1-C2-C3	2.59	130.82	111.07

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	305	MPD	C2-C3-C4-O4
2	A	302	1PG	C2-C3-O2-C4
2	B	301	1PG	O2-C4-C5-O3
2	A	302	1PG	O1-C2-C3-O2
2	A	301	1PG	O3-C6-C7-O4
2	A	301	1PG	O2-C4-C5-O3
2	A	302	1PG	C11-C10-O5-C9
2	A	301	1PG	O5-C10-C11-O6
2	A	302	1PG	O5-C10-C11-O6
2	B	301	1PG	C2-C3-O2-C4
2	A	302	1PG	C3-C2-O1-C1
4	B	304	MPD	O2-C2-C3-C4
4	B	305	MPD	O2-C2-C3-C4
2	B	301	1PG	C4-C5-O3-C6
4	A	305	MPD	C2-C3-C4-C5

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	305	MPD	CM-C2-C3-C4
4	B	305	MPD	C1-C2-C3-C4
4	B	305	MPD	CM-C2-C3-C4
2	B	301	1PG	O1-C2-C3-O2
4	B	305	MPD	C2-C3-C4-C5
2	A	302	1PG	O3-C6-C7-O4
2	A	302	1PG	O4-C8-C9-O5
2	A	302	1PG	C8-C9-O5-C10
2	A	302	1PG	O2-C4-C5-O3
4	B	305	MPD	C2-C3-C4-O4

There are no ring outliers.

7 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	1PG	6	0
4	B	303	MPD	1	0
4	B	305	MPD	1	0
4	A	305	MPD	1	0
2	A	301	1PG	1	0
2	A	302	1PG	19	1
4	B	304	MPD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	269/290 (92%)	0.65	13 (4%) 30 31	10, 15, 29, 47	2 (0%)
1	B	268/290 (92%)	0.79	24 (8%) 9 7	10, 17, 34, 47	1 (0%)
All	All	537/580 (92%)	0.72	37 (6%) 16 15	10, 16, 32, 47	3 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	PHE	14.7
1	B	267	ILE	13.8
1	A	269	LEU	10.0
1	B	14	PHE	8.5
1	B	268	LYS	7.2
1	A	17	LEU	5.9
1	B	55	ILE	5.7
1	B	47	ILE	5.5
1	A	13	ALA	5.5
1	A	267	ILE	4.0
1	A	15	ASP	3.9
1	B	266	ILE	3.8
1	B	225	ARG	3.4
1	B	13	ALA	3.2
1	B	12	ALA	3.1
1	B	265	LEU	3.0
1	B	48	VAL	2.9
1	B	264	GLU	2.9
1	B	52	GLU	2.8
1	A	55	ILE	2.7
1	B	59	GLU	2.7
1	B	25	LYS	2.7
1	B	51	ASN	2.6
1	A	225	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	86	GLU	2.5
1	A	52	GLU	2.5
1	B	61	ASN	2.5
1	B	58	GLU	2.4
1	A	16	ALA	2.3
1	A	47	ILE	2.3
1	A	12	ALA	2.2
1	B	180	LEU	2.2
1	B	49	ASP	2.2
1	A	249	GLY	2.1
1	B	46	SER	2.1
1	B	263	LYS	2.1
1	B	249	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPD	A	305	8/8	0.69	0.29	19,34,37,37	0
4	MPD	B	304	8/8	0.69	0.18	24,30,36,41	0
4	MPD	B	305	8/8	0.74	0.30	28,33,38,39	0
4	MPD	B	303	8/8	0.76	0.17	27,30,37,41	0
2	1PG	A	302	17/17	0.77	0.18	16,31,41,43	0
2	1PG	A	301	17/17	0.80	0.18	25,32,40,42	0
4	MPD	A	304	8/8	0.83	0.17	26,28,33,34	0
2	1PG	B	301	17/17	0.91	0.14	12,21,29,39	0
3	CO	B	302	1/1	1.00	0.08	11,11,11,11	0
3	CO	A	303	1/1	1.00	0.08	10,10,10,10	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 1PG A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.