



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2023 – 10:22 am BST

PDB ID : 8B9M  
Title : Cysteine Synthase from Leishmania Infantum  
Authors : Sowerby, K.V.; Pohl, E.  
Deposited on : 2022-10-06  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.33  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.33


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	 2% 75% 11% 14%
1	B	352	 2% 78% 8% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	401	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9886 atoms, of which 4821 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	304	4731	1474	2410	390	440	17	135	15	0
1	B	306	4720	1472	2395	389	449	15	132	13	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A4ID39
A	2	GLY	-	expression tag	UNP A4ID39
A	3	SER	-	expression tag	UNP A4ID39
A	4	SER	-	expression tag	UNP A4ID39
A	5	HIS	-	expression tag	UNP A4ID39
A	6	HIS	-	expression tag	UNP A4ID39
A	7	HIS	-	expression tag	UNP A4ID39
A	8	HIS	-	expression tag	UNP A4ID39
A	9	HIS	-	expression tag	UNP A4ID39
A	10	HIS	-	expression tag	UNP A4ID39
A	11	SER	-	expression tag	UNP A4ID39
A	12	SER	-	expression tag	UNP A4ID39
A	13	GLY	-	expression tag	UNP A4ID39
A	14	LEU	-	expression tag	UNP A4ID39
A	15	VAL	-	expression tag	UNP A4ID39
A	16	PRO	-	expression tag	UNP A4ID39
A	17	ARG	-	expression tag	UNP A4ID39
A	18	GLY	-	expression tag	UNP A4ID39
A	19	SER	-	expression tag	UNP A4ID39
B	1	MET	-	initiating methionine	UNP A4ID39
B	2	GLY	-	expression tag	UNP A4ID39
B	3	SER	-	expression tag	UNP A4ID39
B	4	SER	-	expression tag	UNP A4ID39
B	5	HIS	-	expression tag	UNP A4ID39
B	6	HIS	-	expression tag	UNP A4ID39

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	7	HIS	-	expression tag	UNP A4ID39
B	8	HIS	-	expression tag	UNP A4ID39
B	9	HIS	-	expression tag	UNP A4ID39
B	10	HIS	-	expression tag	UNP A4ID39
B	11	SER	-	expression tag	UNP A4ID39
B	12	SER	-	expression tag	UNP A4ID39
B	13	GLY	-	expression tag	UNP A4ID39
B	14	LEU	-	expression tag	UNP A4ID39
B	15	VAL	-	expression tag	UNP A4ID39
B	16	PRO	-	expression tag	UNP A4ID39
B	17	ARG	-	expression tag	UNP A4ID39
B	18	GLY	-	expression tag	UNP A4ID39
B	19	SER	-	expression tag	UNP A4ID39

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cl 2 2	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	2	0
			14	3	8	3		
3	B	1	Total	C	H	O	2	0
			14	3	8	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	176	Total	O	0	0
			176	176		
4	B	228	Total	O	0	0
			228	228		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cysteine synthase



- Molecule 1: Cysteine synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.88Å 87.76Å 138.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.17 – 1.75 74.06 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.6 (74.17-1.75) 99.6 (74.06-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, $R_{free}$	0.164 , 0.205 0.177 , 0.214	Depositor DCC
$R_{free}$ test set	3030 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.3	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 43.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9886	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/2397	0.73	0/3245
1	B	0.46	0/2396	0.73	1/3244 (0.0%)
All	All	0.45	0/4793	0.73	1/6489 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	72	ARG	NE-CZ-NH2	-6.11	117.25	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	284	ARG	Sidechain
1	B	307	ARG	Sidechain
1	B	336	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2321	2410	2405	24	1
1	B	2325	2395	2391	37	1
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	B	12	16	16	10	0
4	A	176	0	0	7	0
4	B	228	0	0	17	0
All	All	5065	4821	4812	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:VAL:HA	4:B:581:HOH:O	1.52	1.10
1:B:202:ALA:HB1	3:B:401:GOL:H32	1.17	1.10
1:B:202:ALA:CB	3:B:401:GOL:H32	1.94	0.98
1:B:203:GLY:N	3:B:401:GOL:H31	1.84	0.93
1:B:202:ALA:HB1	3:B:401:GOL:C3	2.02	0.90
1:B:332:TYR:HB2	4:B:508:HOH:O	1.72	0.89
1:A:84:GLU:OE1	1:A:86:LYS:HE2	1.75	0.87
1:B:336:ARG:HD2	4:B:694:HOH:O	1.76	0.84
1:A:232:THR:O	1:A:232:THR:HG22	1.85	0.76
1:B:203:GLY:N	3:B:401:GOL:C3	2.50	0.74
1:B:259:LEU:O	1:B:261:ARG:N	2.22	0.71
1:B:203:GLY:H	3:B:401:GOL:H31	1.55	0.70
1:A:282[B]:LEU:HG	1:A:288:VAL:HB	1.74	0.69
1:B:203:GLY:H	3:B:401:GOL:C3	2.06	0.69
1:B:219:LYS:HD2	4:B:603:HOH:O	1.95	0.66
1:A:232:THR:HB	1:A:271:ALA:HA	1.77	0.66
1:B:51[B]:ASN:H	1:B:51[B]:ASN:HD22	1.43	0.64
1:A:232:THR:O	1:A:232:THR:CG2	2.46	0.64
1:B:51[B]:ASN:H	1:B:51[B]:ASN:ND2	1.97	0.62
1:A:129:ARG:HD2	4:A:566:HOH:O	2.00	0.60

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51[A]:ASN:ND2	4:B:502:HOH:O	2.35	0.60
1:A:129:ARG:NH2	4:A:502:HOH:O	2.35	0.60
1:B:170:ASP:HB2	4:B:520:HOH:O	2.02	0.58
1:B:203:GLY:CA	3:B:401:GOL:H31	2.33	0.58
1:B:282[A]:LEU:HG	1:B:288:VAL:HB	1.89	0.55
1:A:129:ARG:HD3	4:A:657:HOH:O	2.07	0.54
1:B:210:LEU:CD2	4:B:581:HOH:O	2.54	0.54
1:A:260:ASP:O	1:A:261:ARG:NH1	2.39	0.53
1:A:132:LEU:O	1:A:135:ILE:HG12	2.09	0.53
1:A:32:GLN:O	1:B:47[A]:ASN:OD1	2.27	0.52
1:A:125:MET:HE3	4:A:637:HOH:O	2.08	0.52
1:B:149:MET:SD	4:B:630:HOH:O	2.59	0.52
1:B:231[A]:PRO:HB2	4:B:600:HOH:O	2.09	0.52
1:B:231[B]:PRO:HB2	4:B:600:HOH:O	2.09	0.52
1:B:35[A]:ASP:OD1	1:B:109:LEU:HD22	2.10	0.51
1:B:160:VAL:HG13	4:B:596:HOH:O	2.09	0.51
1:A:160:VAL:HG21	1:A:168:LEU:HG	1.95	0.49
1:A:303:LYS:NZ	4:A:501:HOH:O	2.34	0.48
1:B:210:LEU:HD23	4:B:581:HOH:O	2.13	0.48
1:A:47:ASN:ND2	1:B:32:GLN:O	2.47	0.47
1:A:216:ALA:O	1:A:220:MET:HG2	2.15	0.46
1:B:170:ASP:HB3	4:B:612:HOH:O	2.16	0.46
1:B:231[A]:PRO:HB3	4:B:627:HOH:O	2.15	0.45
1:B:231[B]:PRO:HB3	4:B:627:HOH:O	2.15	0.45
1:A:304[A]:ILE:HD13	1:A:307:ARG:CZ	2.46	0.45
1:A:191[B]:GLU:HG2	4:A:548:HOH:O	2.16	0.44
1:B:73:LEU:HD13	1:B:177:ASN:HA	1.99	0.44
1:B:70:LYS:HE3	1:B:207:GLY:HA3	1.99	0.44
1:B:232:THR:O	1:B:271:ALA:HA	2.18	0.44
1:A:97:SER:O	1:A:149[B]:MET:HE1	2.18	0.44
1:B:202:ALA:HB1	3:B:401:GOL:C2	2.48	0.43
1:A:350:LEU:C	1:A:351:GLN:HG2	2.37	0.43
1:A:204:VAL:CG1	1:A:228:ALA:HB1	2.48	0.43
1:B:259:LEU:C	1:B:261:ARG:N	2.72	0.42
1:A:136:PHE:HD1	4:B:509:HOH:O	1.99	0.42
1:A:87:LEU:HD21	1:A:167:VAL:HG21	2.01	0.42
1:B:202:ALA:HB1	3:B:401:GOL:H2	2.00	0.42
1:A:277:GLU:HG3	4:A:596:HOH:O	2.21	0.41
1:B:149:MET:HG3	4:B:630:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:CYS:H	1:B:351:GLN:HE22[4_455]	1.29	0.31

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/352 (90%)	308 (98%)	7 (2%)	0	100	100
1	B	315/352 (90%)	307 (98%)	6 (2%)	2 (1%)	25	10
All	All	630/704 (90%)	615 (98%)	13 (2%)	2 (0%)	41	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	260	ASP
1	B	261	ARG

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/283 (89%)	247 (98%)	6 (2%)	49	26
1	B	254/283 (90%)	253 (100%)	1 (0%)	91	87
All	All	507/566 (90%)	500 (99%)	7 (1%)	65	52

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	SER
1	A	99	SER
1	A	128	GLU
1	A	231	PRO
1	A	333	ARG
1	A	351	GLN
1	B	149	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	40	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	B	402	-	5,5,5	0.20	0	5,5,5	0.55	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	401	-	5,5,5	0.63	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	402	-	-	2/4/4/4	-
3	GOL	B	401	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	GOL	C1-C2-C3-O3
3	B	401	GOL	O1-C1-C2-C3
3	B	402	GOL	O1-C1-C2-C3
3	B	401	GOL	O1-C1-C2-O2
3	B	401	GOL	O2-C2-C3-O3
3	B	402	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	GOL	10	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	304/352 (86%)	-0.07	6 (1%) 65 72	14, 25, 49, 93	0
1	B	306/352 (86%)	-0.10	7 (2%) 60 67	14, 22, 42, 106	0
All	All	610/704 (86%)	-0.09	13 (2%) 63 71	14, 23, 47, 106	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	21	ALA	14.5
1	B	22	ALA	8.1
1	A	22	ALA	6.4
1	B	259	LEU	6.3
1	A	260	ASP	6.0
1	B	260	ASP	5.2
1	A	99	SER	4.8
1	A	23	PRO	4.3
1	B	23	PRO	4.2
1	A	233	GLU	3.5
1	B	233	GLU	3.4
1	B	206	THR	2.9
1	A	164	PRO	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	402	6/6	0.85	0.17	20,35,43,53	2
3	GOL	B	401	6/6	0.87	0.27	18,30,37,41	2
2	CL	A	401	1/1	0.99	0.07	23,23,23,23	0
2	CL	A	402	1/1	0.99	0.09	19,19,19,19	0
2	CL	B	403	1/1	1.00	0.10	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.