

Full wwPDB X-ray Structure Validation Report (i)

Dec 18, 2023 – 09:13 am GMT

PDB ID : 4BBD

Title : THE STRUCTURE OF VACCINIA VIRUS N1 R58Y MUTANT

Authors: Maluquer de Motes, C.; Cooray, S.; McGourty, K.; Ren, H.; Bahar, M.W.;

Stuart, D.I.; Grimes, J.M.; Graham, S.C.; Smith, G.L.

Deposited on : 2012-09-21

Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 5682 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called N1L.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	114	Total	С	N	О	S	0	0	0
1	A	114	955	607	157	184	7	0	0	U
1	В	114	Total	С	N	О	S	0	0	0
1	Ъ	114	955	607	157	184	7	0		U
1	С	114	Total	С	N	О	S	0	0	0
1		114	955	607	157	184	7	0		
1	D	109	Total	С	N	О	S	0	0	0
1	D	109	907	576	151	173	7	0		U
1	E	114	Total	С	N	О	S	0	0	0
1	l Li	114	955	607	157	184	7			U
1	F	F 114	Total	С	N	О	S	0	0	0
			955	607	157	184	7		0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	118	LEU	-	expression tag	UNP Q49PX0
A	119	GLU	-	expression tag	UNP Q49PX0
A	120	HIS	-	expression tag	UNP Q49PX0
A	121	HIS	-	expression tag	UNP Q49PX0
A	122	HIS	-	expression tag	UNP Q49PX0
A	123	HIS	-	expression tag	UNP Q49PX0
A	124	HIS	-	expression tag	UNP Q49PX0
A	125	HIS	-	expression tag	UNP Q49PX0
A	40	SER	CYS	engineered mutation	UNP Q49PX0
A	58	TYR	ARG	engineered mutation	UNP Q49PX0
В	118	LEU	-	expression tag	UNP Q49PX0
В	119	GLU	-	expression tag	UNP Q49PX0
В	120	HIS	-	expression tag	UNP Q49PX0
В	121	HIS		expression tag	UNP Q49PX0
В	122	HIS	-	expression tag	UNP Q49PX0
В	123	HIS	-	expression tag	UNP Q49PX0
В	124	HIS	-	expression tag	UNP Q49PX0

Continued on next page...



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	125	HIS	-	expression tag	UNP Q49PX0
В	40	SER	CYS	engineered mutation	UNP Q49PX0
В	58	TYR	ARG	engineered mutation	UNP Q49PX0
С	118	LEU	-	expression tag	UNP Q49PX0
С	119	GLU	-	expression tag	UNP Q49PX0
С	120	HIS	-	expression tag	UNP Q49PX0
С	121	HIS	-	expression tag	UNP Q49PX0
С	122	HIS	-	expression tag	UNP Q49PX0
С	123	HIS	-	expression tag	UNP Q49PX0
С	124	HIS	-	expression tag	UNP Q49PX0
С	125	HIS	-	expression tag	UNP Q49PX0
С	40	SER	CYS	engineered mutation	UNP Q49PX0
С	58	TYR	ARG	engineered mutation	UNP Q49PX0
D	118	LEU	-	expression tag	UNP Q49PX0
D	119	GLU	-	expression tag	UNP Q49PX0
D	120	HIS	-	expression tag	UNP Q49PX0
D	121	HIS	-	expression tag	UNP Q49PX0
D	122	HIS	-	expression tag	UNP Q49PX0
D	123	HIS	-	expression tag	UNP Q49PX0
D	124	HIS	-	expression tag	UNP Q49PX0
D	125	HIS	-	expression tag	UNP Q49PX0
D	40	SER	CYS	engineered mutation	UNP Q49PX0
D	58	TYR	ARG	engineered mutation	UNP Q49PX0
Е	118	LEU	-	expression tag	UNP Q49PX0
Е	119	GLU	-	expression tag	UNP Q49PX0
Е	120	HIS	-	expression tag	UNP Q49PX0
Е	121	HIS	_	expression tag	UNP Q49PX0
Е	122	HIS	-	expression tag	UNP Q49PX0
Е	123	HIS	-	expression tag	UNP Q49PX0
Е	124	HIS	-	expression tag	UNP Q49PX0
Е	125	HIS	-	expression tag	UNP Q49PX0
E	40	SER	CYS	engineered mutation	UNP Q49PX0
Е	58	TYR	ARG	engineered mutation	UNP Q49PX0
F	118	LEU	-	expression tag	UNP Q49PX0
F	119	GLU	-	expression tag	UNP Q49PX0
F	120	HIS	-	expression tag	UNP Q49PX0
F	121	HIS	-	expression tag	UNP Q49PX0
F	122	HIS	-	expression tag	UNP Q49PX0
F	123	HIS	-	expression tag	UNP Q49PX0
F	124	HIS	-	expression tag	UNP Q49PX0
F	125	HIS	-	expression tag	UNP Q49PX0
F	40	SER	CYS	engineered mutation	UNP Q49PX0

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	58	TYR	ARG	engineered mutation	UNP Q49PX0

 $\operatorname{MolProbity}$ and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	70.56Å 110.20 Å 72.93 Å	Depositor
a, b, c, α , β , γ	90.00° 111.35° 90.00°	Depositor
Resolution (Å)	42.79 - 3.00	Depositor
% Data completeness	(Not available) (42.79-3.00)	Depositor
(in resolution range)		-
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.21 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.176 , 0.200	Depositor
Wilson B-factor $(Å^2)$	70.0	Xtriage
Anisotropy	0.261	Xtriage
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
Total number of atoms	5682	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

