



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 20, 2024 – 12:08 pm GMT

PDB ID : 7BE1  
Title : X-ray structure of Hen Egg White Lysozyme with dirhodium tetraacetate (3)  
Authors : Loreto, D.; Merlino, A.; Ferraro, G.  
Deposited on : 2020-12-22  
Resolution : 1.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	129	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ACT	AAA	216	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
6	ACT	AAA	218	-	-	X	-
6	ACT	AAA	220	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 1282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	129	1065	649	210	196	10	0	8	0

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	AAA	2	Total	Na	0	0
			2	2		

- Molecule 3 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	1
			8	2	6		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0
3	AAA	1	Total N O 4 1 3	0	0

- Molecule 4 is Rhodium (three-letter code: RH) (formula: Rh) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	2	Total Rh 2 2	0	1

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	AAA	1	Total C O 4 2 2	0	0
5	AAA	1	Total C O 4 2 2	0	0
5	AAA	1	Total C O 4 2 2	0	0

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0

- Molecule 7 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	AAA	124	Total 125	O 125	0	2

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysozyme

Chain AAA:  4% 88% 11%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.36Å 78.36Å 37.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.41 – 1.40 55.41 – 1.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (55.41-1.40) 100.0 (55.41-1.40)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.39 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.190 , 0.209 0.196 , 0.221	Depositor DCC
$R_{free}$ test set	1212 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.2	Xtrriage
Anisotropy	0.016	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1282	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, EDO, ACT, RH, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.80	0/1091	0.93	0/1470

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1065	0	1016	14	1
2	AAA	2	0	0	0	0
3	AAA	48	0	0	1	0
4	AAA	2	0	0	0	0
5	AAA	12	0	16	1	0
6	AAA	28	0	21	5	0
7	AAA	125	0	0	5	0
All	All	1282	0	1053	18	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:14[A]:ARG:NH2	3:AAA:222:NO3:O3	2.18	0.74
1:AAA:73:ARG:HD2	6:AAA:218:ACT:H1	1.69	0.73
1:AAA:77[A]:ASN:ND2	7:AAA:301:HOH:O	2.25	0.68
1:AAA:46:ASN:OD1	1:AAA:59[A]:ASN:ND2	2.28	0.67
1:AAA:73:ARG:NH2	7:AAA:302:HOH:O	2.21	0.64

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:129:LEU:O	1:AAA:129:LEU:O[8_555]	2.14	0.06

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	134/129 (104%)	133 (99%)	1 (1%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	113/105 (108%)	111 (98%)	2 (2%)	59 28

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	125	ARG
1	AAA	129	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 4 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NO3	AAA	221	-	1,3,3	0.44	0	0,3,3	-	-
3	NO3	AAA	205	-	1,3,3	0.97	0	0,3,3	-	-
6	ACT	AAA	220	-	3,3,3	1.05	0	3,3,3	0.91	0
3	NO3	AAA	203[A]	-	1,3,3	0.60	0	0,3,3	-	-
6	ACT	AAA	217	-	3,3,3	1.86	1 (33%)	3,3,3	0.81	0
3	NO3	AAA	204	-	1,3,3	0.07	0	0,3,3	-	-
6	ACT	AAA	215	4	3,3,3	0.90	0	3,3,3	0.73	0
6	ACT	AAA	216	-	3,3,3	0.84	0	3,3,3	0.85	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NO3	AAA	223	-	1,3,3	0.51	0	0,3,3	-	-
3	NO3	AAA	209	-	1,3,3	0.06	0	0,3,3	-	-
6	ACT	AAA	224	-	3,3,3	1.09	0	3,3,3	0.75	0
5	EDO	AAA	211	2	3,3,3	0.08	0	2,2,2	0.37	0
5	EDO	AAA	213	-	3,3,3	0.15	0	2,2,2	0.25	0
6	ACT	AAA	219	4	3,3,3	0.93	0	3,3,3	0.87	0
3	NO3	AAA	206	-	1,3,3	0.17	0	0,3,3	-	-
3	NO3	AAA	207	-	1,3,3	0.35	0	0,3,3	-	-
3	NO3	AAA	222	-	1,3,3	0.37	0	0,3,3	-	-
6	ACT	AAA	218	-	3,3,3	0.84	0	3,3,3	0.88	0
3	NO3	AAA	208	-	1,3,3	0.88	0	0,3,3	-	-
5	EDO	AAA	212	-	3,3,3	0.17	0	2,2,2	0.33	0
3	NO3	AAA	203[B]	-	1,3,3	1.72	0	0,3,3	-	-
3	NO3	AAA	202	2	1,3,3	0.65	0	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	AAA	211	2	-	0/1/1/1	-
5	EDO	AAA	212	-	-	1/1/1/1	-
5	EDO	AAA	213	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	AAA	217	ACT	OXT-C	-2.88	1.16	1.30

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	AAA	212	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	AAA	220	ACT	2	0
6	AAA	216	ACT	3	0
5	AAA	211	EDO	1	0
3	AAA	222	NO3	1	0
6	AAA	218	ACT	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	129/129 (100%)	0.43	5 (3%) 39 39	7, 14, 26, 39	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	71	GLY	3.5
1	AAA	125	ARG	2.5
1	AAA	62	TRP	2.4
1	AAA	129	LEU	2.1
1	AAA	121	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NO3	AAA	209	4/4	0.45	0.35	43,54,54,54	0
5	EDO	AAA	212	4/4	0.49	0.23	57,58,60,60	0


*Continued on next page...*


*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	ACT	AAA	219	4/4	0.53	0.23	32,33,33,36	4
6	ACT	AAA	216	4/4	0.64	0.22	50,50,51,53	0
6	ACT	AAA	224	4/4	0.68	0.30	39,43,44,48	0
6	ACT	AAA	215	4/4	0.73	0.19	41,42,44,46	0
3	NO3	AAA	204	4/4	0.77	0.27	21,21,25,36	0
5	EDO	AAA	213	4/4	0.77	0.26	33,34,37,37	0
6	ACT	AAA	218	4/4	0.79	0.21	60,61,62,66	0
3	NO3	AAA	203[A]	4/4	0.79	0.22	15,19,20,21	4
3	NO3	AAA	203[B]	4/4	0.79	0.22	16,18,19,21	4
3	NO3	AAA	206	4/4	0.83	0.25	25,29,33,43	0
3	NO3	AAA	207	4/4	0.83	0.17	23,36,37,41	0
3	NO3	AAA	205	4/4	0.89	0.14	27,28,29,31	0
3	NO3	AAA	208	4/4	0.89	0.17	23,31,32,33	0
5	EDO	AAA	211	4/4	0.91	0.10	22,24,24,28	0
3	NO3	AAA	202	4/4	0.91	0.13	18,23,23,25	0
3	NO3	AAA	223	4/4	0.93	0.16	44,46,46,46	4
4	RH	AAA	210	1/1	0.93	0.07	23,23,23,23	1
3	NO3	AAA	221	4/4	0.94	0.10	15,16,17,19	0
3	NO3	AAA	222	4/4	0.94	0.30	35,35,35,37	4
6	ACT	AAA	220	4/4	0.94	0.24	30,39,42,45	0
2	NA	AAA	225	1/1	0.94	0.12	24,24,24,24	0
2	NA	AAA	201	1/1	0.96	0.12	13,13,13,13	0
6	ACT	AAA	217	4/4	0.96	0.25	18,26,27,27	0
4	RH	AAA	214[A]	1/1	0.96	0.07	26,26,26,26	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.