



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2023 – 10:15 pm BST

PDB ID : 8BEY
Title : Structure of the Lysinibacillus sphaericus Tpp49Aa1 pesticidal protein at pH 7
Authors : Williamson, L.J.; Rizkallah, P.J.; Berry, C.; Oberthur, D.; Galchenkova, M.; Yefanov, O.; Bean, R.
Deposited on : 2022-10-22
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	
1	B	416	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cry49Aa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	416	3421	2151	617	642	11	0	8	0
1	B	416	3456	2171	628	647	10	0	12	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	242	242	242	0	0
2	B	238	238	238	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.65Å 83.11Å 156.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.98 – 1.62 24.97 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.98-1.62) 100.0 (24.97-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.176 , 0.195 0.185 , 0.201	Depositor DCC
R_{free} test set	6927 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtrriage
Anisotropy	0.264	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.109 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7357	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.01	2/3519 (0.1%)	0.79	1/4782 (0.0%)
1	B	0.99	1/3557 (0.0%)	0.81	4/4834 (0.1%)
All	All	1.00	3/7076 (0.0%)	0.80	5/9616 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82	ILE	C-O	5.52	1.33	1.23
1	B	82	ILE	C-O	5.25	1.33	1.23
1	A	92	ARG	C-O	5.03	1.32	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	331	ARG	NE-CZ-NH2	7.11	123.86	120.30
1	B	331	ARG	NE-CZ-NH1	-6.31	117.15	120.30
1	B	309	ASN	CB-CA-C	5.16	120.72	110.40
1	A	424	GLU	CB-CA-C	-5.09	100.22	110.40
1	B	424	GLU	CB-CA-C	-5.01	100.39	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3421	0	3325	18	0
1	B	3456	0	3362	14	0
2	A	242	0	0	5	0
2	B	238	0	0	6	0
All	All	7357	0	6687	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:407[B]:ARG:NH1	1:B:436[B]:ASP:OD2	1.94	1.00
1:B:357[B]:ASN:OD1	2:B:501:HOH:O	1.93	0.86
1:B:69[A]:ASN:OD1	2:B:502:HOH:O	1.98	0.81
1:A:262[B]:ASN:ND2	2:A:502:HOH:O	2.18	0.76
1:B:262[B]:ASN:ND2	2:B:503:HOH:O	2.26	0.69
1:A:79:PRO:HD3	1:A:97[B]:HIS:ND1	2.13	0.64
1:A:95:VAL:HG21	1:A:141:ASN:HB2	1.81	0.61
1:A:367:VAL:HG13	1:A:423:LEU:CD1	2.34	0.57
1:A:357[A]:ASN:OD1	2:A:501:HOH:O	2.17	0.56
1:B:319[A]:MET:HE2	2:B:629:HOH:O	2.06	0.54
1:A:319[A]:MET:HE2	2:A:652:HOH:O	2.08	0.52
1:A:367:VAL:HG13	1:A:423:LEU:HD11	1.93	0.50
1:A:51:ASN:HD21	1:B:350:ASN:HA	1.78	0.49
1:A:79:PRO:CD	1:A:97[B]:HIS:ND1	2.76	0.47
1:A:156:HIS:HD2	2:A:718:HOH:O	1.96	0.47
1:B:165:ASN:C	1:B:165:ASN:HD22	2.18	0.47
1:B:342[B]:ARG:HG3	1:B:343:GLN:N	2.31	0.46
1:A:91:CYS:SG	1:A:184:TRP:CG	3.10	0.45
1:A:232:ARG:HH11	1:A:232:ARG:HB3	1.82	0.44
1:A:116:PHE:HB3	1:A:124:PHE:CD1	2.52	0.44
1:B:415:TYR:HA	1:B:416:PRO:C	2.37	0.44
1:A:415:TYR:HA	1:A:416:PRO:C	2.37	0.43
1:A:95:VAL:HG13	1:A:140:ALA:HB3	1.99	0.43
1:B:116:PHE:HB3	1:B:124:PHE:CD1	2.53	0.43
1:B:91:CYS:SG	1:B:184:TRP:CG	3.12	0.42
1:B:137:PHE:CE2	1:B:176:THR:HG21	2.54	0.41
1:B:219:PRO:HD3	2:B:528:HOH:O	2.19	0.41
1:A:75:LEU:HD21	1:A:190:TRP:HH2	1.86	0.41
1:B:318:LYS:NZ	2:B:514:HOH:O	2.54	0.41
1:A:137:PHE:CE2	1:A:176:THR:HG21	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:PRO:HD3	2:A:529:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/416 (101%)	413 (98%)	9 (2%)	0	100	100
1	B	426/416 (102%)	417 (98%)	9 (2%)	0	100	100
All	All	848/832 (102%)	830 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/371 (102%)	374 (99%)	5 (1%)	69	49
1	B	383/371 (103%)	374 (98%)	9 (2%)	50	24
All	All	762/742 (103%)	748 (98%)	14 (2%)	59	34

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	196	ARG
1	A	232	ARG
1	A	253	ARG
1	A	360	TYR
1	B	60	GLN
1	B	69[A]	ASN
1	B	69[B]	ASN
1	B	165	ASN
1	B	196	ARG
1	B	253	ARG
1	B	289	ILE
1	B	309	ASN
1	B	360	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	ASN
1	A	270	ASN
1	A	274	GLN
1	A	336	ASN
1	B	68	ASN
1	B	123	ASN
1	B	165	ASN
1	B	270	ASN
1	B	284	GLN
1	B	314	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/416 (100%)	0.11	25 (6%) 21 19	18, 31, 60, 82	0
1	B	416/416 (100%)	0.07	18 (4%) 35 31	17, 31, 57, 73	0
All	All	832/832 (100%)	0.09	43 (5%) 27 24	17, 31, 57, 82	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	ASN	4.8
1	A	200	PRO	4.7
1	B	49	ASN	4.7
1	B	107	THR	4.4
1	A	107	THR	4.4
1	B	200	PRO	4.2
1	A	141	ASN	4.1
1	A	184	TRP	4.0
1	A	201	GLY	3.9
1	B	464	ASN	3.9
1	B	201	GLY	3.5
1	A	199	ASN	3.4
1	B	193	ILE	3.3
1	B	141	ASN	3.2
1	B	456	HIS	3.2
1	B	100	LEU	3.1
1	A	142	GLY	2.9
1	B	72	PHE	2.8
1	A	84	GLY	2.8
1	A	93	ILE	2.6
1	B	104	LEU	2.6
1	B	199	ASN	2.6
1	A	456	HIS	2.6
1	A	464	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	457	GLU	2.5
1	A	88	ASP	2.5
1	A	104	LEU	2.5
1	A	76	ASN	2.4
1	B	139	ILE	2.4
1	B	146	ILE	2.4
1	A	69	ASN	2.3
1	A	108	GLY	2.3
1	A	50	LEU	2.2
1	A	146	ILE	2.2
1	B	184	TRP	2.2
1	B	197	VAL	2.2
1	A	100	LEU	2.1
1	A	202	ALA	2.1
1	B	135	LEU	2.1
1	A	68	ASN	2.1
1	A	457	GLU	2.0
1	A	97[A]	HIS	2.0
1	A	190	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.