



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2022 – 06:21 pm GMT

PDB ID : 7BIC
Title : Crystal structure of human GSTA1-1 bound to allyl-isothiocyanate
Authors : Schwartz, M.; Neiers, F.
Deposited on : 2021-01-12
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.26
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	 5% 76% 14% • 9%
1	B	222	 3% 76% 17% • 5%
1	C	222	 6% 75% 15% • 9%
1	D	222	 5% 72% 21% 7%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6681 atoms, of which 6 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione S-transferase A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	201	Total 1626	C 1057	N 272	O 290	S 7	0	0	0
1	B	210	Total 1697	C 1101	N 282	O 306	S 8	0	0	0
1	C	203	Total 1636	C 1061	N 273	O 294	S 8	0	0	0
1	D	207	Total 1674	C 1086	N 278	O 302	S 8	0	0	0

- Molecule 2 is N-prop-2-en-1-ylthioformamide (three-letter code: 9AI) (formula: C₄H₇NS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	S		
2	B	1	Total 12	C 4	H 6	N 1	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	10	Total O 10 10	0	0
3	B	9	Total O 9 9	0	0
3	C	11	Total O 11 11	0	0
3	D	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutathione S-transferase A1




- Molecule 1: Glutathione S-transferase A1



- Molecule 1: Glutathione S-transferase A1



- Molecule 1: Glutathione S-transferase A1



K129	M130	R131	L148	R155	L160	V161	E162	L163	L164	Y165	Y166	D171	S172	S173	L174	I175	K182	T186	R187	I188	K196	F197	L198	G199	P200	R204	E210	LYS	SER	LEU	GLU	GLU	ALA	ARG	ARG	LYS	ILE	PHE	ARG	PHE
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	100.11Å 94.65Å 104.51Å 90.00° 92.01° 90.00°	Depositor
Resolution (Å)	47.33 – 2.46 47.33 – 2.46	Depositor EDS
% Data completeness (in resolution range)	98.6 (47.33-2.46) 99.0 (47.33-2.46)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.17_3644	Depositor
R, R_{free}	0.230 , 0.277 0.230 , 0.275	Depositor DCC
R_{free} test set	1765 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	50.1	Xtrriage
Anisotropy	0.563	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.001 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6681	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 63.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.6953e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9AI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1657	0.69	0/2226
1	B	0.51	0/1731	0.65	0/2329
1	C	0.59	0/1667	0.73	0/2241
1	D	0.50	0/1708	0.75	0/2299
All	All	0.52	0/6763	0.71	0/9095

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1626	0	1698	35	0
1	B	1697	0	1766	34	0
1	C	1636	0	1696	27	0
1	D	1674	0	1738	34	0
2	B	6	6	6	0	0
3	A	10	0	0	0	0
3	B	9	0	0	0	0
3	C	11	0	0	0	0
3	D	6	0	0	2	0
All	All	6675	6	6904	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:LYS:HA	1:A:120:LYS:HE2	1.28	1.09
1:D:102:LEU:HA	1:D:105:MET:HE3	1.40	1.02
1:D:113:PRO:HB2	1:D:116:GLU:OE1	1.70	0.90
1:B:183:ALA:O	1:B:187:ARG:HG3	1.73	0.87
1:A:89:ARG:NE	1:B:89:ARG:HH12	1.81	0.78
1:D:102:LEU:HA	1:D:105:MET:CE	2.16	0.73
1:B:141:LYS:HG3	1:B:180:LEU:CD1	2.19	0.73
1:D:116:GLU:HB3	1:D:120:LYS:NZ	2.04	0.72
1:C:109:LEU:N	1:C:110:PRO:HD2	2.06	0.70
1:A:117:LYS:CA	1:A:120:LYS:HE2	2.16	0.68
1:D:109:LEU:HB3	1:D:110:PRO:HD3	1.76	0.67
1:B:141:LYS:HG3	1:B:180:LEU:HD11	1.78	0.66
1:A:175:ILE:HG12	1:A:182:LYS:HG2	1.79	0.65
1:B:213:LEU:HD23	1:B:213:LEU:O	1.95	0.65
1:A:109:LEU:CB	1:A:110:PRO:CD	2.75	0.65
1:D:116:GLU:HB3	1:D:120:LYS:HZ3	1.63	0.63
1:B:210:GLU:O	1:B:213:LEU:HB3	1.99	0.62
1:C:111:VAL:HG23	1:C:111:VAL:O	1.99	0.61
1:B:102:LEU:HD23	1:B:163:LEU:HD21	1.82	0.61
1:B:82:TYR:CE1	1:B:89:ARG:HG3	2.36	0.60
1:C:123:LEU:HD11	1:C:127:LYS:HE2	1.83	0.59
1:D:182:LYS:O	1:D:186:THR:HG23	2.03	0.59
1:D:82:TYR:CE1	1:D:89:ARG:HG3	2.38	0.58
1:A:125:LYS:HG2	1:A:174:LEU:HD11	1.84	0.58
1:D:102:LEU:HD23	1:D:163:LEU:HD21	1.84	0.57
1:A:17:GLU:HG2	1:A:166:TYR:OH	2.04	0.57
1:C:175:ILE:HG12	1:C:182:LYS:HG2	1.86	0.57
1:B:141:LYS:HG3	1:B:180:LEU:HD12	1.85	0.57
1:C:17:GLU:HG2	1:C:166:TYR:OH	2.06	0.55
1:C:102:LEU:HD21	1:C:128:ILE:HG12	1.90	0.54
1:A:39:GLU:O	1:A:43:LYS:HG3	2.07	0.54
1:A:86:ILE:HD12	1:B:74:TYR:HE1	1.72	0.54
1:A:109:LEU:HB3	1:A:110:PRO:HD3	1.90	0.54
1:A:87:LYS:HD2	1:B:63:MET:HE1	1.89	0.54
1:C:90:ALA:HA	3:D:305:HOH:O	2.08	0.54
1:C:198:LEU:O	1:C:204:ARG:NH2	2.36	0.53
1:A:128:ILE:O	1:A:133:PHE:HB2	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:102:LEU:HD21	1:D:128:ILE:HG12	1.89	0.53
1:C:123:LEU:O	1:C:123:LEU:HG	2.07	0.52
1:C:102:LEU:HD22	1:C:132:TYR:CD2	2.45	0.52
1:D:90:ALA:HA	3:D:303:HOH:O	2.09	0.52
1:D:66:VAL:O	1:D:67:GLN:HB2	2.10	0.51
1:A:121:LEU:CD2	1:A:125:LYS:HD3	2.41	0.51
1:B:72:LEU:HB3	1:B:155:ARG:NH2	2.25	0.51
1:C:86:ILE:HG12	1:D:74:TYR:CD1	2.46	0.51
1:D:17:GLU:HG2	1:D:166:TYR:OH	2.10	0.51
1:D:175:ILE:HG12	1:D:182:LYS:HG2	1.92	0.51
1:A:66:VAL:O	1:A:67:GLN:HB2	2.10	0.50
1:B:200:PRO:HG3	1:D:204:ARG:CZ	2.42	0.50
1:C:109:LEU:N	1:C:110:PRO:CD	2.74	0.50
1:B:206:PRO:HA	1:D:200:PRO:HD2	1.93	0.50
1:C:66:VAL:O	1:C:67:GLN:HB2	2.12	0.49
1:D:164:LEU:HD22	1:D:175:ILE:HD12	1.94	0.49
1:A:82:TYR:CE1	1:A:89:ARG:HG2	2.48	0.49
1:C:128:ILE:O	1:C:133:PHE:HB2	2.12	0.49
1:B:47:ASP:O	1:B:49:TYR:HD1	1.96	0.49
1:C:127:LYS:O	1:C:131:ARG:HB3	2.13	0.48
1:B:66:VAL:O	1:B:67:GLN:HB2	2.14	0.48
1:A:84:LYS:HG2	1:A:88:GLU:OE1	2.13	0.48
1:A:109:LEU:CB	1:A:110:PRO:HD3	2.42	0.48
1:C:178:PHE:O	1:C:182:LYS:HG3	2.13	0.48
1:C:208:MET:HG3	1:C:209:ASP:H	1.79	0.48
1:A:95:TYR:HE1	1:B:51:MET:CE	2.27	0.47
1:D:49:TYR:HB3	1:D:57:MET:HE2	1.96	0.47
1:A:102:LEU:HD21	1:A:128:ILE:HG12	1.98	0.46
1:B:109:LEU:HD21	1:B:121:LEU:HD13	1.97	0.46
1:C:109:LEU:HD23	1:C:109:LEU:HA	1.61	0.46
1:A:121:LEU:C	1:A:121:LEU:HD23	2.36	0.46
1:A:86:ILE:HD12	1:A:87:LYS:HG2	1.97	0.46
1:A:120:LYS:HE3	1:A:120:LYS:HB2	1.47	0.46
1:B:145:GLN:CD	1:B:151:ASN:HB2	2.36	0.45
1:D:198:LEU:O	1:D:204:ARG:NH2	2.40	0.45
1:D:116:GLU:HB3	1:D:120:LYS:HZ2	1.80	0.45
1:B:63:MET:HE1	1:B:74:TYR:CE1	2.52	0.45
1:D:112:CYS:SG	1:D:120:LYS:HD3	2.57	0.45
1:A:9:TYR:HE1	1:A:56:PRO:HD3	1.82	0.44
1:A:87:LYS:HD2	1:B:63:MET:CE	2.47	0.44
1:B:182:LYS:O	1:B:186:THR:HG23	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:TYR:HB2	1:A:16:MET:CE	2.47	0.44
1:B:84:LYS:HG3	1:B:88:GLU:OE1	2.16	0.44
1:B:120:LYS:HA	1:B:120:LYS:HD3	1.51	0.44
1:B:4:LYS:HB2	1:B:4:LYS:HE2	1.85	0.44
1:A:95:TYR:HE1	1:B:51:MET:HE1	1.82	0.43
1:B:125:LYS:HG2	1:B:174:LEU:HD11	2.00	0.43
1:A:86:ILE:HD12	1:B:74:TYR:CE1	2.53	0.43
1:C:82:TYR:CD1	1:C:89:ARG:HG2	2.54	0.43
1:D:112:CYS:HB2	1:D:120:LYS:HE2	2.01	0.43
1:C:28:VAL:HG21	1:C:79:TYR:CZ	2.53	0.43
1:B:102:LEU:HD21	1:B:128:ILE:HG12	2.01	0.43
1:C:208:MET:HG3	1:C:209:ASP:N	2.33	0.43
1:D:161:VAL:HG21	1:D:188:ILE:HB	2.01	0.43
1:A:28:VAL:HG21	1:A:79:TYR:CZ	2.53	0.43
1:C:96:ILE:HA	1:C:99:ILE:HD12	2.01	0.42
1:B:17:GLU:HG2	1:B:166:TYR:OH	2.19	0.42
1:D:72:LEU:HB3	1:D:155:ARG:NH2	2.35	0.42
1:D:126:GLU:HG2	1:D:130:ASN:OD1	2.19	0.42
1:C:35:ILE:HD11	1:C:55:VAL:HG11	2.01	0.42
1:D:28:VAL:HG21	1:D:79:TYR:CZ	2.54	0.42
1:B:145:GLN:HG2	1:B:151:ASN:CG	2.40	0.42
1:D:148:LEU:HD11	1:D:160:LEU:HD23	2.01	0.42
1:C:49:TYR:CE1	1:C:64:LYS:HE3	2.54	0.42
1:C:174:LEU:HA	1:C:174:LEU:HD23	1.87	0.42
1:A:109:LEU:N	1:A:110:PRO:HD2	2.34	0.41
1:B:47:ASP:HB3	1:B:49:TYR:CD1	2.55	0.41
1:B:47:ASP:HB3	1:B:49:TYR:CE1	2.55	0.41
1:D:28:VAL:HG21	1:D:79:TYR:CE2	2.55	0.41
1:A:86:ILE:CD1	1:A:87:LYS:HG2	2.50	0.41
1:D:174:LEU:HA	1:D:174:LEU:HD23	1.78	0.41
1:A:121:LEU:HD23	1:A:125:LYS:HD3	2.03	0.41
1:C:82:TYR:O	1:C:89:ARG:HG3	2.21	0.41
1:D:4:LYS:HE3	1:D:4:LYS:HB3	1.90	0.41
1:A:9:TYR:CE1	1:A:56:PRO:HD3	2.55	0.41
1:A:174:LEU:HA	1:A:174:LEU:HD23	1.85	0.41
1:C:127:LYS:NZ	1:C:131:ARG:NH1	2.69	0.41
1:D:127:LYS:O	1:D:131:ARG:HB3	2.21	0.41
1:A:94:MET:SD	1:B:66:VAL:HG22	2.61	0.41
1:D:148:LEU:HD11	1:D:160:LEU:CD2	2.50	0.40
1:D:171:ASP:HB3	1:D:174:LEU:CD1	2.51	0.40
1:A:28:VAL:HG21	1:A:79:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/222 (89%)	192 (98%)	5 (2%)	0	100	100
1	B	208/222 (94%)	201 (97%)	7 (3%)	0	100	100
1	C	199/222 (90%)	190 (96%)	9 (4%)	0	100	100
1	D	205/222 (92%)	199 (97%)	5 (2%)	1 (0%)	29	34
All	All	809/888 (91%)	782 (97%)	26 (3%)	1 (0%)	51	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	114	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/196 (90%)	171 (97%)	5 (3%)	43	56
1	B	186/196 (95%)	177 (95%)	9 (5%)	25	33
1	C	177/196 (90%)	171 (97%)	6 (3%)	37	48
1	D	183/196 (93%)	178 (97%)	5 (3%)	44	57
All	All	722/784 (92%)	697 (96%)	25 (4%)	36	47

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LYS
1	A	36	LYS
1	A	43	LYS
1	A	82	TYR
1	A	120	LYS
1	B	4	LYS
1	B	10	PHE
1	B	43	LYS
1	B	82	TYR
1	B	84	LYS
1	B	120	LYS
1	B	142	SER
1	B	208	MET
1	B	211	LYS
1	C	6	LYS
1	C	82	TYR
1	C	109	LEU
1	C	112	CYS
1	C	123	LEU
1	C	126	GLU
1	D	6	LYS
1	D	43	LYS
1	D	82	TYR
1	D	173	SER
1	D	196	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	9AI	B	401	1	5,5,5	1.02	1 (20%)	2,4,4	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9AI	B	401	1	-	1/1/3/3	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	9AI	C1-N1	2.12	1.38	1.31

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	401	9AI	C4-C5-C6-N1

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/222 (90%)	0.47	11 (5%) 25 22	35, 52, 72, 94	0
1	B	210/222 (94%)	0.30	7 (3%) 46 43	39, 53, 78, 101	0
1	C	203/222 (91%)	0.57	13 (6%) 19 16	31, 46, 78, 94	0
1	D	207/222 (93%)	0.45	10 (4%) 30 28	33, 51, 85, 96	0
All	All	821/888 (92%)	0.45	41 (4%) 28 26	31, 51, 80, 101	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	110	PRO	6.7
1	D	10	PHE	4.6
1	A	117	LYS	4.3
1	D	110	PRO	4.2
1	C	121	LEU	4.1
1	D	38	ALA	4.0
1	C	2	ALA	3.8
1	B	211	LYS	3.8
1	A	109	LEU	3.8
1	C	108	LEU	3.6
1	B	212	SER	3.5
1	D	41	LEU	3.4
1	A	110	PRO	3.4
1	C	111	VAL	3.3
1	A	10	PHE	3.3
1	C	107	LEU	3.3
1	A	118	ASP	3.3
1	A	41	LEU	3.2
1	B	213	LEU	3.1
1	A	123	LEU	3.0
1	B	37	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	170	LEU	2.9
1	D	114	PRO	2.8
1	D	37	SER	2.8
1	C	208	MET	2.8
1	A	120	LYS	2.7
1	C	3	GLU	2.6
1	D	112	CYS	2.6
1	B	71	ILE	2.5
1	D	35	ILE	2.4
1	B	43	LYS	2.4
1	C	123	LEU	2.4
1	A	3	GLU	2.4
1	D	113	PRO	2.2
1	A	42	ASP	2.2
1	A	174	LEU	2.2
1	B	44	LEU	2.2
1	C	10	PHE	2.1
1	D	53	GLN	2.1
1	C	127	LYS	2.0
1	C	103	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	9AI	B	401	6/6	0.74	0.18	61,73,88,88	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.