



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 11:44 am BST

PDB ID : 4BJ6
Title : Crystal structure Rif2 in complex with the C-terminal domain of Rap1 (Rap1-RCT)
Authors : Shi, T.; Bunker, R.D.; Gut, H.; Scrima, A.; Thoma, N.H.
Deposited on : 2013-04-16
Resolution : 3.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1191 (3.30-3.22)
Clashscore	141614	1251 (3.30-3.22)
Ramachandran outliers	138981	1229 (3.30-3.22)
Sidechain outliers	138945	1228 (3.30-3.22)
RSRZ outliers	127900	1154 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	 0% 68% 12% 20%
1	B	365	 2% 66% 12% 21%
2	C	202	 0% 66% 8% 25%
2	E	202	 0% 69% 6% 25%
3	D	13	 8% 69% 31%
3	F	13	 15% 62% 38%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	1390	-	-	-	X
4	SO4	A	1391	-	-	-	X
4	SO4	E	1827	-	-	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAP1-INTERACTING FACTOR 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	2362	1529	378	442	13	0	0	0
1	B	290	2332	1510	371	438	13	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	GLY	-	expression tag	UNP Q06208
A	32	GLY	-	expression tag	UNP Q06208
A	33	GLY	-	expression tag	UNP Q06208
A	34	ARG	-	expression tag	UNP Q06208
B	31	GLY	-	expression tag	UNP Q06208
B	32	GLY	-	expression tag	UNP Q06208
B	33	GLY	-	expression tag	UNP Q06208
B	34	ARG	-	expression tag	UNP Q06208

- Molecule 2 is a protein called DNA-BINDING PROTEIN RAP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	151	1227	779	204	237	7	0	0	0
2	E	151	1227	779	204	237	7	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	626	GLY	-	expression tag	UNP P11938
E	626	GLY	-	expression tag	UNP P11938

- Molecule 3 is a protein called RAP1-INTERACTING FACTOR 2.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	13	Total	C	N	O	0	0	0
			84	56	14	14			
3	F	13	Total	C	N	O	0	0	0
			88	59	15	14			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 2: DNA-BINDING PROTEIN RAP1



- Molecule 3: RAP1-INTERACTING FACTOR 2



- Molecule 3: RAP1-INTERACTING FACTOR 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	108.25Å 112.68Å 140.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.79 – 3.26 48.79 – 3.26	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.79-3.26) 99.6 (48.79-3.26)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 3.25Å)	Xtrriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.172 , 0.195 0.184 , 0.206	Depositor DCC
R_{free} test set	1361 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	100.3	Xtrriage
Anisotropy	0.389	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 83.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.014 for k,h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7370	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2408	0.70	0/3257
1	B	0.52	0/2377	0.69	0/3216
2	C	0.46	0/1258	0.61	0/1700
2	E	0.48	0/1258	0.61	0/1700
3	D	0.54	0/84	0.76	0/116
3	F	0.55	0/88	0.76	0/120
All	All	0.50	0/7473	0.67	0/10109

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2362	0	2353	17	0
1	B	2332	0	2310	20	0
2	C	1227	0	1171	7	0
2	E	1227	0	1171	5	0
3	D	84	0	77	2	0
3	F	88	0	88	3	0
4	A	15	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	20	0	0	0	0
4	C	5	0	0	0	0
4	E	10	0	0	0	0
All	All	7370	0	7170	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:VAL:HA	1:B:259:SER:HB3	1.70	0.73
1:A:256:VAL:HA	1:A:259:SER:HB3	1.70	0.72
1:B:354:THR:HG23	1:B:357:GLU:HB2	1.81	0.62
1:A:312:LEU:O	1:A:316:GLN:HB2	2.01	0.61
1:B:312:LEU:O	1:B:316:GLN:HB2	2.02	0.59
1:A:354:THR:HG23	1:A:357:GLU:H	1.68	0.57
1:B:259:SER:O	1:B:263:SER:HB2	2.05	0.57
1:A:199:ASN:HB3	1:A:202:GLU:HG2	1.87	0.57
1:B:199:ASN:HB3	1:B:202:GLU:HG2	1.87	0.56
1:B:330:PHE:CE1	1:B:334:PHE:HB2	2.40	0.55
1:A:207:ILE:HD12	1:A:248:LYS:HE2	1.90	0.54
2:C:761:ASP:HA	3:D:45:VAL:HB	1.88	0.54
1:B:145:MET:HA	1:B:145:MET:HE2	1.88	0.53
1:A:240:GLU:HB2	1:A:247:PHE:HE2	1.74	0.53
1:B:379:LYS:HB3	1:B:382:GLN:HG2	1.89	0.52
1:A:379:LYS:HB3	1:A:382:GLN:HG2	1.91	0.52
1:B:207:ILE:HD12	1:B:248:LYS:HE2	1.91	0.52
1:B:240:GLU:HB2	1:B:247:PHE:HE2	1.75	0.52
1:A:145:MET:HA	1:A:145:MET:HE2	1.92	0.51
1:B:119:ILE:HG12	1:B:380:LYS:HA	1.93	0.50
1:A:330:PHE:CE1	1:A:334:PHE:HB2	2.46	0.50
1:A:119:ILE:HG12	1:A:380:LYS:HA	1.93	0.50
1:B:379:LYS:O	1:B:382:GLN:HB2	2.13	0.49
1:A:379:LYS:O	1:A:382:GLN:HB2	2.13	0.48
1:B:139:ASN:HB3	1:B:206:HIS:CD2	2.49	0.48
2:E:799:ASP:HB3	2:E:802:GLN:HE21	1.79	0.48
2:E:761:ASP:HA	3:F:45:VAL:HB	1.95	0.47
2:C:764:VAL:HG23	2:C:767:ARG:HD2	1.96	0.46
2:C:799:ASP:HB3	2:C:802:GLN:HE21	1.79	0.46
1:A:66:VAL:HA	1:A:354:THR:OG1	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:ILE:HG23	1:A:248:LYS:HD3	1.98	0.46
1:B:384:LEU:HD12	2:C:706:LEU:HD12	1.98	0.45
1:B:299:PHE:O	1:B:306:LEU:HD22	2.16	0.45
1:A:299:PHE:O	1:A:306:LEU:HD22	2.17	0.45
1:B:207:ILE:HG23	1:B:248:LYS:HD3	1.98	0.44
2:E:734:GLU:HG3	3:F:39:LEU:HD13	2.01	0.43
2:C:730:PRO:HA	3:D:44:LEU:HB2	2.00	0.42
1:A:346:THR:O	1:A:350:VAL:HB	2.19	0.42
1:B:75:LYS:HD2	1:B:350:VAL:HA	2.02	0.42
2:E:768:TYR:HA	2:E:786:ILE:HD11	2.02	0.42
1:B:312:LEU:HD22	1:B:359:LEU:HD13	2.02	0.41
2:C:768:TYR:HA	2:C:786:ILE:HD11	2.01	0.41
1:B:139:ASN:HB3	1:B:206:HIS:HD2	1.85	0.41
1:B:346:THR:O	1:B:350:VAL:HB	2.20	0.41
2:C:758:LEU:HD13	2:C:764:VAL:HG13	2.03	0.41
2:E:730:PRO:HA	3:F:44:LEU:HB2	2.03	0.41
1:A:354:THR:HG22	1:A:357:GLU:HB2	2.02	0.40
1:A:312:LEU:HD22	1:A:359:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/365 (79%)	276 (96%)	12 (4%)	1 (0%)	41	72
1	B	286/365 (78%)	272 (95%)	13 (4%)	1 (0%)	41	72
2	C	149/202 (74%)	144 (97%)	3 (2%)	2 (1%)	12	41
2	E	149/202 (74%)	145 (97%)	2 (1%)	2 (1%)	12	41
3	D	11/13 (85%)	11 (100%)	0	0	100	100
3	F	11/13 (85%)	11 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	895/1160 (77%)	859 (96%)	30 (3%)	6 (1%)	22 56

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	732	GLN
2	C	781	PRO
2	E	781	PRO
2	C	732	GLN
1	A	387	LYS
1	B	387	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	271/346 (78%)	252 (93%)	19 (7%)	15 43
1	B	266/346 (77%)	245 (92%)	21 (8%)	12 37
2	C	140/188 (74%)	135 (96%)	5 (4%)	35 63
2	E	140/188 (74%)	137 (98%)	3 (2%)	53 75
3	D	7/13 (54%)	5 (71%)	2 (29%)	0 1
3	F	8/13 (62%)	6 (75%)	2 (25%)	0 2
All	All	832/1094 (76%)	780 (94%)	52 (6%)	18 47

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	ARG
1	A	69	VAL
1	A	92	VAL
1	A	94	SER
1	A	96	SER
1	A	114	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	127	ARG
1	A	139	ASN
1	A	147	ASP
1	A	200	ASN
1	A	213	GLU
1	A	214	GLN
1	A	215	LEU
1	A	219	SER
1	A	220	THR
1	A	224	ILE
1	A	275	THR
1	A	345	TYR
1	A	365	ARG
1	B	69	VAL
1	B	92	VAL
1	B	96	SER
1	B	114	ASP
1	B	127	ARG
1	B	135	LEU
1	B	137	ASP
1	B	147	ASP
1	B	200	ASN
1	B	213	GLU
1	B	214	GLN
1	B	215	LEU
1	B	219	SER
1	B	220	THR
1	B	224	ILE
1	B	256	VAL
1	B	275	THR
1	B	320	LYS
1	B	345	TYR
1	B	354	THR
1	B	384	LEU
2	C	727	ASP
2	C	757	CYS
2	C	765	PHE
2	C	807	VAL
2	C	815	MET
3	D	42	LEU
3	D	47	ILE
2	E	727	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	765	PHE
2	E	807	VAL
3	F	42	LEU
3	F	47	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	ASN
1	A	206	HIS
1	A	244	ASN
1	A	323	ASN
1	A	370	ASN
1	B	139	ASN
1	B	206	HIS
1	B	244	ASN
1	B	323	ASN
1	B	370	ASN
2	C	709	HIS
2	C	749	ASN
2	C	778	ASN
2	C	802	GLN
2	E	709	HIS
2	E	715	GLN
2	E	749	ASN
2	E	778	ASN
2	E	802	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	1389	-	4,4,4	0.37	0	6,6,6	0.34	0
4	SO4	A	1391	-	4,4,4	0.15	0	6,6,6	0.15	0
4	SO4	B	1390	-	4,4,4	0.27	0	6,6,6	0.34	0
4	SO4	E	1826	-	4,4,4	0.29	0	6,6,6	0.42	0
4	SO4	E	1827	-	4,4,4	0.15	0	6,6,6	0.08	0
4	SO4	B	1392	-	4,4,4	0.22	0	6,6,6	0.15	0
4	SO4	C	1826	-	4,4,4	0.12	0	6,6,6	0.25	0
4	SO4	B	1391	-	4,4,4	0.17	0	6,6,6	0.11	0
4	SO4	B	1389	-	4,4,4	0.26	0	6,6,6	0.45	0
4	SO4	A	1390	-	4,4,4	0.20	0	6,6,6	0.16	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/365 (80%)	-0.06	4 (1%) 75 74	62, 94, 145, 190	0
1	B	290/365 (79%)	0.15	8 (2%) 53 50	72, 106, 176, 196	0
2	C	151/202 (74%)	0.15	1 (0%) 87 88	87, 116, 177, 199	0
2	E	151/202 (74%)	0.15	3 (1%) 65 63	73, 106, 166, 187	0
3	D	13/13 (100%)	0.82	1 (7%) 13 12	126, 148, 164, 165	0
3	F	13/13 (100%)	0.70	2 (15%) 2 2	116, 140, 160, 163	0
All	All	911/1160 (78%)	0.10	19 (2%) 63 61	62, 105, 168, 199	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	675	SER	5.1
1	B	64	GLU	4.7
1	B	220	THR	3.5
1	B	221	SER	3.4
3	F	39	LEU	3.2
1	B	219	SER	2.9
3	D	44	LEU	2.7
1	A	302	GLY	2.6
1	A	219	SER	2.6
1	B	70	PHE	2.6
1	A	61	PRO	2.5
1	B	362	ILE	2.5
3	F	42	LEU	2.4
1	B	299	PHE	2.4
2	E	675	SER	2.3
2	E	821	PHE	2.2
2	E	823	GLU	2.1
1	A	60	SER	2.1
1	B	150	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	1391	5/5	0.53	0.69	235,235,235,235	0
4	SO4	B	1391	5/5	0.67	0.33	213,214,214,214	0
4	SO4	E	1827	5/5	0.70	0.73	230,230,230,230	0
4	SO4	A	1390	5/5	0.75	0.46	187,187,187,188	0
4	SO4	B	1392	5/5	0.84	0.33	174,175,176,176	0
4	SO4	C	1826	5/5	0.85	0.19	181,181,182,183	0
4	SO4	E	1826	5/5	0.88	0.22	135,136,138,138	0
4	SO4	B	1390	5/5	0.95	0.22	113,113,118,119	0
4	SO4	B	1389	5/5	0.99	0.20	79,80,82,84	0
4	SO4	A	1389	5/5	0.99	0.21	74,77,78,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.