



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 23, 2017 – 11:42 AM EST

PDB ID : 6BMT  
Title : Crystal Structure of a Recombinant form of Human Myeloperoxidase Bound to an Inhibitor from *Staphylococcus delphini*  
Authors : Ploscariu, N.T.; Geisbrecht, B.V.  
Deposited on : 2017-11-15  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	745	 2% 73% 23%
2	B	105	 2% 60% 7% 33%

## 2 Entry composition [\(i\)](#)

There are 7 unique types of molecules in this entry. The entry contains 5335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myeloperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	573	4584	2889	838	825	32	0	0	0


- Molecule 2 is a protein called Hypothetical Protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	70	559	345	97	116	1	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
4	A	1	43	34	1	4	4	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
5	A	1	1	1	0	0

- Molecule 6 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>N<sub>1</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	14	8	1	5	0	0
6	A	1	14	8	1	5	0	0
6	A	1	14	8	1	5	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	102	102	102	0	0
7	B	3	3	3	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.63Å 90.67Å 125.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.34 – 2.40 45.34 – 2.40	Depositor EDS
% Data completeness (in resolution range)	94.3 (45.34-2.40) 93.5 (45.34-2.40)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.48 (at 2.39Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.185 , 0.218 0.182 , 0.217	Depositor DCC
$R_{free}$ test set	1875 reflections (5.23%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.3	Xtrriage
Anisotropy	0.425	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 34.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5335	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CA, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/4695	0.70	2/6376 (0.0%)
2	B	0.55	0/566	0.62	0/765
All	All	0.55	0/5261	0.69	2/7141 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	354	ARG	NE-CZ-NH2	-5.33	117.63	120.30
1	A	341	MET	CA-CB-CG	5.05	121.89	113.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4584	0	4539	29	0
2	B	559	0	544	6	0
3	A	1	0	0	0	0
4	A	43	0	30	4	0
5	A	1	0	0	0	0
6	A	42	0	39	13	0
7	A	102	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	3	0	0	0	0
All	All	5335	0	5152	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:391:ASN:HD21	6:A:806:NAG:C1	1.01	1.61
1:A:355:ASN:HD21	6:A:805:NAG:C1	0.97	1.58
1:A:483:ASN:HD21	6:A:804:NAG:C1	0.87	1.51
1:A:483:ASN:ND2	6:A:804:NAG:C1	1.73	1.43
1:A:355:ASN:ND2	6:A:805:NAG:C1	1.81	1.39
1:A:391:ASN:ND2	6:A:806:NAG:C1	1.82	1.38
1:A:408:GLU:OE2	4:A:802:HEM:HMB3	1.47	1.14
1:A:408:GLU:OE2	4:A:802:HEM:CMB	2.16	0.93
1:A:483:ASN:ND2	6:A:804:NAG:O5	2.18	0.74
1:A:408:GLU:CD	4:A:802:HEM:HMB3	2.10	0.72
1:A:355:ASN:CG	6:A:805:NAG:C1	2.57	0.71
1:A:391:ASN:ND2	6:A:806:NAG:O5	2.14	0.70
1:A:324:ILE:HG23	1:A:325:THR:HG23	1.76	0.66
1:A:483:ASN:CG	6:A:804:NAG:C1	2.62	0.64
1:A:391:ASN:CG	6:A:806:NAG:C1	2.61	0.64
4:A:802:HEM:HHD	4:A:802:HEM:HBC2	1.83	0.61
1:A:355:ASN:ND2	6:A:805:NAG:O5	2.30	0.60
1:A:404:THR:HG21	2:B:30:THR:HG21	1.89	0.54
1:A:583:LEU:HB3	1:A:584:PRO:HD3	1.92	0.52
1:A:394:ALA:HB2	6:A:806:NAG:H61	1.92	0.51
1:A:720:SER:HB3	1:A:726:ASP:HB3	1.92	0.51
1:A:213:TRP:O	1:A:287:GLN:NE2	2.46	0.49
1:A:370:GLN:OE1	1:A:373:GLY:HA2	2.16	0.46
1:A:268:GLU:HB2	1:A:269:PRO:HD2	1.97	0.45
2:B:28:GLN:N	2:B:28:GLN:OE1	2.49	0.45
1:A:387:CYS:HA	1:A:532:PHE:O	2.16	0.45
2:B:30:THR:HG22	2:B:38:HIS:HB3	1.98	0.45
1:A:672:LEU:O	1:A:676:ASP:HB2	2.17	0.44
1:A:391:ASN:HB2	1:A:535:TRP:NE1	2.32	0.43
1:A:480:ARG:O	1:A:481:SER:HB3	2.19	0.43
1:A:193:ARG:HG2	1:A:193:ARG:HH11	1.84	0.43
2:B:94:SER:O	2:B:95:LYS:CB	2.68	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:48:LEU:HD13	2:B:86:ALA:CB	2.50	0.42
2:B:94:SER:O	2:B:95:LYS:HB3	2.20	0.41
1:A:177:THR:O	1:A:190:ALA:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	569/745 (76%)	549 (96%)	18 (3%)	2 (0%)	38	54
2	B	68/105 (65%)	63 (93%)	5 (7%)	0	100	100
All	All	637/850 (75%)	612 (96%)	23 (4%)	2 (0%)	44	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	240	ASP
1	A	207	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/643 (78%)	501 (100%)	1 (0%)	94	98

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	64/88 (73%)	64 (100%)	0	100	100
All	All	566/731 (77%)	565 (100%)	1 (0%)	94	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	341	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	355	ASN
1	A	391	ASN
1	A	416	HIS
1	A	483	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all  $Z$  scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	HEM	A	802	1	28,50,50	1.94	4 (14%)	17,82,82	2.06	4 (23%)
6	NAG	A	804	-	14,14,15	0.29	0	15,19,21	0.57	0
6	NAG	A	805	-	14,14,15	0.28	0	15,19,21	0.58	0
6	NAG	A	806	-	14,14,15	0.28	0	15,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	802	1	-	0/6/54/54	0/0/8/8
6	NAG	A	804	-	-	0/6/23/26	0/1/1/1
6	NAG	A	805	-	-	0/6/23/26	0/1/1/1
6	NAG	A	806	-	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	802	HEM	C3B-C2B	-7.06	1.31	1.40
4	A	802	HEM	C3C-C2C	-4.33	1.34	1.40
4	A	802	HEM	C1B-NB	-3.29	1.32	1.36
4	A	802	HEM	C4D-ND	-2.28	1.34	1.36

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	802	HEM	CBA-CAA-C2A	-2.57	107.57	112.48
4	A	802	HEM	C4A-C3A-C2A	2.08	108.44	107.00
4	A	802	HEM	CAA-CBA-CGA	3.83	119.21	112.66
4	A	802	HEM	CAD-CBD-CGD	5.57	122.17	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	802	HEM	4	0
6	A	804	NAG	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	805	NAG	4	0
6	A	806	NAG	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	573/745 (76%)	-0.20	13 (2%) 61 58	39, 53, 75, 106	0
2	B	70/105 (66%)	-0.03	2 (2%) 52 50	50, 63, 91, 98	0
All	All	643/850 (75%)	-0.18	15 (2%) 61 58	39, 55, 78, 106	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	744	ALA	7.3
1	A	321	GLY	6.8
1	A	324	ILE	6.5
1	A	278	GLY	4.9
1	A	320	PRO	3.4
1	A	521	PRO	3.2
2	B	58	ASN	3.2
1	A	169	GLU	2.9
2	B	62	GLN	2.5
1	A	288	GLN	2.5
1	A	170	GLN	2.1
1	A	392	ARG	2.1
1	A	322	SER	2.1
1	A	256	GLY	2.1
1	A	259	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	A	806	14/15	0.84	0.17	1.62	60,71,79,86	0
6	NAG	A	805	14/15	0.90	0.16	1.01	52,67,79,79	0
3	CA	A	801	1/1	1.00	0.18	0.07	45,45,45,45	0
4	HEM	A	802	43/43	0.93	0.15	-0.78	50,57,63,67	0
5	CL	A	803	1/1	0.99	0.10	-1.56	49,49,49,49	0
6	NAG	A	804	14/15	0.83	0.17	-	71,81,87,88	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.