



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2023 – 12:01 PM EDT

PDB ID : 3BNE
Title : Lipoyxygenase-1 (Soybean) I553A Mutant
Authors : Tomchick, D.R.
Deposited on : 2007-12-14
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	839	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Seed lipoxygenase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	818	6651	4257	1124	1249	21	0	18	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	GLU	SER	engineered mutation	UNP P08170
A	553	ALA	ILE	engineered mutation	UNP P08170

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	692	Total	O	0	0
			692	692		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Seed lipoxygenase-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.11Å 92.95Å 49.42Å 90.00° 90.76° 90.00°	Depositor
Resolution (Å)	29.72 – 1.40 29.72 – 1.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.72-1.40) 99.4 (29.72-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.2.0009	Depositor
R, R_{free}	0.166 , 0.186 0.170 , 0.191	Depositor DCC
R_{free} test set	5011 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.005 for k,h,-l 0.006 for -k,-h,-l 0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7344	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/6816	0.78	9/9261 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	402	ARG	NE-CZ-NH2	-7.03	116.78	120.30
1	A	588	ARG	NE-CZ-NH1	6.13	123.36	120.30
1	A	341	MET	CG-SD-CE	-6.00	90.60	100.20
1	A	543	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	A	712	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	A	435	LEU	CB-CG-CD2	5.38	120.15	111.00
1	A	543	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	588	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	613	ASP	CB-CG-OD1	5.05	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6651	0	6591	37	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	692	0	0	6	0
All	All	7344	0	6591	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:495:GLN:HG2	3:A:1443:HOH:O	1.57	1.01
1:A:55:LYS:HG2	1:A:160[B]:GLU:HG3	1.41	0.98
1:A:55:LYS:HG2	1:A:160[B]:GLU:CG	2.13	0.78
1:A:160[B]:GLU:H	1:A:160[B]:GLU:CD	1.87	0.76
1:A:552:ILE:HG21	1:A:750[A]:VAL:HG21	1.71	0.72
1:A:495:GLN:CG	3:A:1443:HOH:O	2.27	0.71
1:A:713:ARG:CZ	3:A:1564:HOH:O	2.39	0.71
1:A:254:ALA:O	1:A:257:ILE:HG12	1.98	0.64
1:A:270:PHE:HD2	1:A:746[B]:ILE:HD13	1.65	0.62
1:A:301:ARG:HG3	1:A:323:HIS:NE2	2.17	0.60
1:A:160[B]:GLU:CD	1:A:160[B]:GLU:N	2.55	0.59
1:A:270:PHE:HD2	1:A:746[B]:ILE:CD1	2.15	0.59
1:A:41[B]:SER:HB3	1:A:63:PHE:CE2	2.40	0.56
1:A:160[A]:GLU:OE1	1:A:160[A]:GLU:HA	2.11	0.51
1:A:55:LYS:CG	1:A:160[B]:GLU:HG3	2.29	0.50
1:A:497:MET:SD	1:A:567:SER:HA	2.52	0.50
1:A:196:ASP:OD2	1:A:226[A]:VAL:HG22	2.11	0.50
1:A:354[B]:VAL:CG2	1:A:700:TYR:HB2	2.41	0.50
1:A:795:LYS:HE2	1:A:799:ARG:NH2	2.28	0.49
1:A:270:PHE:CD2	1:A:746[B]:ILE:HD13	2.46	0.49
1:A:382:LYS:HB2	1:A:382:LYS:HZ3	1.78	0.48
1:A:573:ASN:OD1	3:A:1486:HOH:O	2.20	0.48
1:A:767:ARG:NE	3:A:1562:HOH:O	2.39	0.47
1:A:750[B]:VAL:O	1:A:754:LEU:HG	2.15	0.46
1:A:495:GLN:HG3	1:A:557:PHE:CZ	2.51	0.46
1:A:379:GLN:OE1	1:A:410:HIS:HE1	1.99	0.45
1:A:65:GLU:HG2	1:A:82:ASN:O	2.16	0.44
1:A:586:ILE:HD11	1:A:601:VAL:HG12	2.00	0.43
1:A:308:ILE:CD1	1:A:321:GLY:HA2	2.48	0.43
1:A:88:ASP:OD1	1:A:88:ASP:C	2.58	0.43
1:A:304:ILE:HG13	1:A:323:HIS:HA	2.01	0.42
1:A:418:ARG:NH1	1:A:458:ASP:OD1	2.48	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:767:ARG:NH2	3:A:1562:HOH:O	2.51	0.42
1:A:649:TRP:HB2	1:A:669:LEU:HD22	2.02	0.42
1:A:593:LYS:HD2	1:A:600:GLY:O	2.20	0.42
1:A:307:ILE:CD1	1:A:325:LEU:HD11	2.50	0.41
1:A:301:ARG:CG	1:A:323:HIS:NE2	2.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	830/839 (99%)	806 (97%)	24 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	731/729 (100%)	727 (100%)	4 (0%)	88 74

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	308	ILE
1	A	535	ASN
1	A	608	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	234	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	818/839 (97%)	0.63	76 (9%) 8 8	10, 21, 31, 48	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	THR	13.6
1	A	71	LEU	13.4
1	A	117	ILE	8.8
1	A	72	PRO	8.6
1	A	456	ALA	8.0
1	A	116	ALA	6.6
1	A	53	HIS	6.5
1	A	70	SER	6.5
1	A	69	THR	6.5
1	A	67	ILE	5.4
1	A	662	ASP	5.1
1	A	461	ALA	5.0
1	A	18	ASN	5.0
1	A	308	ILE	4.5
1	A	459	LEU	4.5
1	A	52	ALA	4.4
1	A	595	PRO	4.3
1	A	31	ASN	4.3
1	A	88	ASP	4.2
1	A	598	PRO	4.1
1	A	457	GLY	3.8
1	A	455	SER	3.7
1	A	74	LEU	3.6
1	A	424	ASN	3.6
1	A	682	ILE	3.5
1	A	246	LEU	3.5
1	A	32	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	91	MET	3.3
1	A	597	THR	3.2
1	A	425	SER	3.2
1	A	93	ILE	3.1
1	A	54	GLY	3.0
1	A	160[A]	GLU	3.0
1	A	486	VAL	3.0
1	A	721	PRO	3.0
1	A	458	ASP	2.9
1	A	257	ILE	2.9
1	A	289	VAL	2.9
1	A	302	ASP	2.9
1	A	75	GLY	2.8
1	A	19	GLU	2.8
1	A	7	LYS	2.8
1	A	596	SER	2.7
1	A	487	ILE	2.6
1	A	634	ALA	2.6
1	A	683	ILE	2.6
1	A	226[A]	VAL	2.6
1	A	138[A]	LYS	2.5
1	A	719	GLY	2.4
1	A	460	SER	2.4
1	A	378	ASP	2.4
1	A	104	GLN	2.4
1	A	68	ASN	2.4
1	A	472	GLU	2.4
1	A	86	GLU	2.4
1	A	109	LEU	2.4
1	A	51	ASP	2.3
1	A	115	GLU	2.3
1	A	532	TYR	2.3
1	A	303	VAL	2.3
1	A	59	GLY	2.3
1	A	284	HIS	2.2
1	A	528	LEU	2.2
1	A	9	LYS	2.2
1	A	275	ASP	2.1
1	A	717	GLU	2.1
1	A	637	ASP	2.1
1	A	485	TYR	2.1
1	A	309	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	236	GLU	2.0
1	A	511	VAL	2.0
1	A	776	SER	2.0
1	A	837	ILE	2.0
1	A	602	ARG	2.0
1	A	122	THR	2.0
1	A	76	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FE	A	840	1/1	1.00	0.06	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.