



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 09:11 pm BST

PDB ID : 4BRK  
Title : Legionella pneumophila NTPDase1 N302Y variant crystal form III (closed) in complex with MG UMPPNP  
Authors : Zebisch, M.; Schaefer, P.; Lauble, P.; Straeter, N.  
Deposited on : 2013-06-04  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	 2% 89% 8%
1	B	368	 % 89% 9%

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	359	2950	1884	476	574	16	0	21	0
1	B	360	2982	1902	482	583	15	0	20	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	-	expression tag	UNP Q5ZUA2
A	394	LEU	-	expression tag	UNP Q5ZUA2
A	395	GLU	-	expression tag	UNP Q5ZUA2
A	396	HIS	-	expression tag	UNP Q5ZUA2
A	397	HIS	-	expression tag	UNP Q5ZUA2
A	398	HIS	-	expression tag	UNP Q5ZUA2
A	399	HIS	-	expression tag	UNP Q5ZUA2
A	400	HIS	-	expression tag	UNP Q5ZUA2
A	401	HIS	-	expression tag	UNP Q5ZUA2
A	137	ASP	GLU	conflict	UNP Q5ZUA2
A	149	VAL	ALA	conflict	UNP Q5ZUA2
A	302	TYR	ASN	engineered mutation	UNP Q5ZUA2
B	34	MET	-	expression tag	UNP Q5ZUA2
B	394	LEU	-	expression tag	UNP Q5ZUA2
B	395	GLU	-	expression tag	UNP Q5ZUA2
B	396	HIS	-	expression tag	UNP Q5ZUA2
B	397	HIS	-	expression tag	UNP Q5ZUA2
B	398	HIS	-	expression tag	UNP Q5ZUA2
B	399	HIS	-	expression tag	UNP Q5ZUA2
B	400	HIS	-	expression tag	UNP Q5ZUA2
B	401	HIS	-	expression tag	UNP Q5ZUA2
B	137	ASP	GLU	conflict	UNP Q5ZUA2
B	149	VAL	ALA	conflict	UNP Q5ZUA2
B	302	TYR	ASN	engineered mutation	UNP Q5ZUA2





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
			Total	C	N	O			S	
4	A	1	Total	12	6	1	4	1	0	0
4	B	1	Total	12	6	1	4	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			
5	A	1	Total	6	3	3	0	0
5	B	1	Total	6	3	3	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Cl 1	0	0
6	A	2	Total 2	Cl 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	412	Total 415	O 415	0	3
7	B	354	Total 356	O 356	0	2


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I



- Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.60Å 82.97Å 109.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.43 – 1.50 27.41 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (27.43-1.50) 99.5 (27.41-1.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.129 , 0.178 0.126 , 0.177	Depositor DCC
$R_{free}$ test set	1183 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.3	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6802	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNP, GOL, MG, MES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.03	1/3089 (0.0%)	0.94	10/4211 (0.2%)
1	B	0.99	2/3113 (0.1%)	0.96	5/4244 (0.1%)
All	All	1.01	3/6202 (0.0%)	0.95	15/8455 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	122	ARG	CD-NE	-5.69	1.36	1.46
1	B	122	ARG	CD-NE	-5.05	1.37	1.46
1	B	384	TRP	CD2-CE2	5.05	1.47	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	122	ARG	NE-CZ-NH1	17.78	129.19	120.30
1	B	122	ARG	NE-CZ-NH2	-15.97	112.31	120.30
1	A	122	ARG	NE-CZ-NH2	-12.18	114.21	120.30
1	A	122	ARG	NE-CZ-NH1	10.04	125.32	120.30
1	B	234	MET	CG-SD-CE	8.29	113.46	100.20
1	A	234	MET	CG-SD-CE	8.21	113.34	100.20
1	A	96	ASP	CB-CG-OD1	6.67	124.30	118.30
1	B	122	ARG	CD-NE-CZ	6.22	132.31	123.60
1	A	65	ASP	CB-CG-OD1	6.12	123.81	118.30
1	A	392[A]	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	392[B]	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	323	GLN	CA-CB-CG	5.36	125.18	113.40
1	A	323	GLN	N-CA-CB	-5.17	101.29	110.60
1	A	122	ARG	CG-CD-NE	-5.06	101.17	111.80
1	B	96	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2950	0	2850	22	0
1	B	2982	0	2864	25	0
2	A	29	0	12	0	0
2	B	29	0	12	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	12	0	13	0	0
4	B	12	0	13	3	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	2	0	0	0	0
6	B	1	0	0	0	0
7	A	415	0	0	13	0
7	B	356	0	0	10	0
All	All	6802	0	5780	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253[B]:ASP:HB3	7:A:2276:HOH:O	1.56	1.03
7:A:2260:HOH:O	4:B:1399:MES:H72	1.59	1.02
1:B:346[B]:TYR:OH	7:B:2329:HOH:O	1.76	1.00
1:A:213[A]:ASN:ND2	1:A:218:ASN:OD1	2.09	0.85
1:B:301[A]:SER:OG	7:B:2280:HOH:O	1.90	0.84
1:B:189:GLY:HA2	1:B:231[A]:GLN:NE2	1.96	0.80
7:A:2259:HOH:O	4:B:1399:MES:O2S	1.99	0.80
7:A:2260:HOH:O	4:B:1399:MES:C7	2.23	0.79
1:A:301[A]:SER:OG	7:A:2328:HOH:O	1.90	0.78

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:333[B]:GLN:OE1	1:A:336[B]:ILE:HG12	1.85	0.76
1:B:333[B]:GLN:NE2	7:B:2316:HOH:O	2.10	0.75
1:A:392[B]:ARG:NH2	7:A:2408:HOH:O	2.24	0.69
1:A:161:LEU:HD11	1:A:212:LEU:HD22	1.75	0.67
1:A:232:THR:CG2	7:A:2248:HOH:O	2.46	0.64
1:A:232:THR:HG22	7:A:2248:HOH:O	2.00	0.62
1:B:126:GLN:HG2	7:B:2117:HOH:O	1.98	0.61
1:B:210[B]:VAL:HG22	1:B:221:LEU:HB2	1.81	0.61
1:B:346[A]:TYR:HB2	7:B:2328:HOH:O	2.01	0.58
1:A:109:HIS:HB3	1:A:145:GLN:HB3	1.86	0.58
1:A:213[B]:ASN:ND2	7:A:2234:HOH:O	2.39	0.55
1:B:231[B]:GLN:NE2	1:B:357:TYR:OH	2.39	0.55
1:B:204:ILE:HD11	1:B:285:LEU:HD21	1.89	0.55
1:A:126:GLN:HG2	7:A:2068:HOH:O	2.06	0.53
1:A:80:LYS:HE3	7:A:2026:HOH:O	2.08	0.53
1:B:265:LYS:NZ	7:B:2252:HOH:O	2.32	0.53
1:B:189:GLY:HA2	1:B:231[A]:GLN:HE21	1.73	0.52
1:B:386:ILE:HG13	7:B:2028:HOH:O	2.12	0.50
1:A:323:GLN:NE2	7:A:2359:HOH:O	2.45	0.49
1:A:86:ALA:HB2	1:A:121[B]:MET:SD	2.53	0.48
1:B:37:ASN:HD21	1:B:108:ILE:HD12	1.78	0.48
1:B:224:HIS:HE1	1:B:281:GLN:OE1	1.96	0.47
1:B:136:LEU:HD23	1:B:136:LEU:C	2.35	0.47
1:B:64[A]:THR:HG21	7:B:2034:HOH:O	2.13	0.46
1:A:114:TYR:OH	1:A:392[A]:ARG:HD2	2.16	0.45
1:B:119:ALA:HB3	1:B:190:ALA:HB3	1.99	0.45
1:A:110:ASN:OD1	1:A:110:ASN:N	2.50	0.45
1:B:346[B]:TYR:CD1	2:B:1397:UNP:H2'	2.52	0.45
1:B:210[A]:VAL:HG13	1:B:221:LEU:HB2	1.99	0.44
1:B:211:GLU:CG	1:B:218[A]:ASN:HD21	2.31	0.44
1:A:136:LEU:C	1:A:136:LEU:HD23	2.39	0.43
1:A:126:GLN:HG3	7:A:2142:HOH:O	2.19	0.43
1:A:45:ILE:HG21	1:A:45:ILE:HD13	1.69	0.42
1:B:64[A]:THR:CG2	7:B:2034:HOH:O	2.67	0.42
1:B:161:LEU:HD13	1:B:210[B]:VAL:HG21	2.00	0.42
1:B:346[B]:TYR:CE1	2:B:1397:UNP:H2'	2.55	0.41
1:B:273:ASN:O	1:B:277:LYS:HD3	2.20	0.41
1:A:119:ALA:HB3	1:A:190:ALA:HB3	2.02	0.41
1:A:170[A]:LYS:HG3	1:A:382:LEU:HD13	2.01	0.41
1:B:70:PRO:HD3	7:B:2042:HOH:O	2.21	0.41
1:A:286:LEU:HA	1:A:286:LEU:HD23	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	378/368 (103%)	367 (97%)	11 (3%)	0	100	100
1	B	377/368 (102%)	366 (97%)	11 (3%)	0	100	100
All	All	755/736 (103%)	733 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/328 (103%)	335 (99%)	2 (1%)	86	74
1	B	340/328 (104%)	337 (99%)	3 (1%)	78	61
All	All	677/656 (103%)	672 (99%)	5 (1%)	81	69

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ARG
1	A	212	LEU
1	B	56	ARG
1	B	344	ASP
1	B	356	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	HIS
1	A	318	ASN
1	A	323	GLN
1	B	157	ASN
1	B	179	ASN
1	B	323	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	UNP	B	1397	3	25,30,30	1.78	9 (36%)	27,47,47	1.48	3 (11%)
4	MES	A	1397	-	12,12,12	1.89	1 (8%)	14,16,16	3.37	8 (57%)
4	MES	B	1399	-	12,12,12	1.81	1 (8%)	14,16,16	3.27	5 (35%)
5	GOL	A	1398	-	5,5,5	0.67	0	5,5,5	0.43	0
2	UNP	A	1395	3	25,30,30	1.70	6 (24%)	27,47,47	1.41	4 (14%)
5	GOL	B	1400	-	5,5,5	0.76	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UNP	B	1397	3	-	1/16/38/38	0/2/2/2
4	MES	A	1397	-	-	1/6/14/14	0/1/1/1
4	MES	B	1399	-	-	4/6/14/14	0/1/1/1
5	GOL	A	1398	-	-	0/4/4/4	-
2	UNP	A	1395	3	-	1/16/38/38	0/2/2/2
5	GOL	B	1400	-	-	0/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1397	MES	C8-S	-5.96	1.69	1.77
4	B	1399	MES	C8-S	-5.49	1.69	1.77
2	A	1395	UNP	O4'-C1'	3.77	1.46	1.41
2	B	1397	UNP	PB-O3B	3.33	1.51	1.46
2	A	1395	UNP	PB-N2B	2.98	1.71	1.63
2	A	1395	UNP	PB-O1B	-2.97	1.48	1.56
2	B	1397	UNP	PG-O2G	2.82	1.50	1.46
2	B	1397	UNP	PB-O1B	-2.79	1.49	1.56
2	B	1397	UNP	O4'-C1'	2.60	1.44	1.41
2	B	1397	UNP	C2-N3	-2.58	1.33	1.38
2	A	1395	UNP	PB-O3A	2.48	1.62	1.59
2	B	1397	UNP	PG-O3G	-2.44	1.50	1.56
2	B	1397	UNP	PG-N2B	2.28	1.69	1.63
2	A	1395	UNP	PG-O3G	-2.27	1.50	1.56
2	A	1395	UNP	PB-O3B	2.24	1.49	1.46
2	B	1397	UNP	C2'-C1'	-2.24	1.50	1.53
2	B	1397	UNP	PB-N2B	2.07	1.68	1.63

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1399	MES	O3S-S-C8	8.94	120.22	105.77
4	A	1397	MES	O1S-S-C8	8.08	116.64	106.92
2	B	1397	UNP	O2G-PG-N2B	-4.76	104.76	111.77
4	A	1397	MES	O2S-S-C8	-4.66	101.30	106.92
4	B	1399	MES	O3S-S-O2S	-4.34	100.68	111.27
4	A	1397	MES	C7-N4-C3	-4.04	100.92	111.23
2	A	1395	UNP	O2G-PG-N2B	-3.90	106.03	111.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1397	MES	C6-C5-N4	3.81	115.88	110.10
4	A	1397	MES	O1-C6-C5	-3.77	103.49	111.80
4	B	1399	MES	C2-C3-N4	3.59	115.55	110.10
2	B	1397	UNP	O1B-PB-O3B	3.07	116.35	109.92
4	B	1399	MES	C5-N4-C3	3.06	115.72	108.83
4	A	1397	MES	O3S-S-C8	2.94	110.52	105.77
4	B	1399	MES	O1-C6-C5	2.84	118.06	111.80
2	A	1395	UNP	C3'-C2'-C1'	2.84	105.26	100.98
4	A	1397	MES	O2S-S-O1S	-2.54	105.16	113.95
2	A	1395	UNP	O1B-PB-O3B	2.32	114.78	109.92
2	A	1395	UNP	O1G-PG-O3G	2.29	113.73	107.64
2	B	1397	UNP	O2'-C2'-C1'	-2.28	102.44	110.85
4	A	1397	MES	C7-N4-C5	2.01	116.37	111.23

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1397	UNP	PB-N2B-PG-O2G
4	B	1399	MES	N4-C7-C8-S
4	B	1399	MES	C7-C8-S-O1S
4	B	1399	MES	C7-C8-S-O3S
4	B	1399	MES	C7-C8-S-O2S
4	A	1397	MES	C7-C8-S-O2S
2	A	1395	UNP	PB-N2B-PG-O2G


There are no ring outliers.

2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1397	UNP	2	0
4	B	1399	MES	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	359/368 (97%)	-0.18	7 (1%) 66 71	7, 13, 32, 52	0
1	B	360/368 (97%)	-0.24	5 (1%) 75 79	7, 13, 29, 49	0
All	All	719/736 (97%)	-0.21	12 (1%) 70 75	7, 13, 31, 52	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	204	ILE	4.8
1	A	36	THR	4.4
1	B	36	THR	4.1
1	A	205	SER	3.8
1	B	66[A]	ASP	3.6
1	A	394	LEU	3.0
1	A	37	ASN	2.6
1	A	104	ALA	2.6
1	B	179	ASN	2.4
1	B	37	ASN	2.3
1	B	394	LEU	2.2
1	A	393	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MES	B	1399	12/12	0.95	0.12	18,26,32,32	12
4	MES	A	1397	12/12	0.96	0.09	14,19,25,29	12
5	GOL	B	1400	6/6	0.96	0.15	21,27,28,33	0
5	GOL	A	1398	6/6	0.98	0.05	11,14,16,17	0
2	UNP	A	1395	29/29	0.99	0.06	7,9,24,30	0
2	UNP	B	1397	29/29	0.99	0.07	6,12,36,50	0
6	CL	A	1400	1/1	0.99	0.08	15,15,15,15	1
6	CL	A	1399	1/1	1.00	0.04	11,11,11,11	0
3	MG	B	1398	1/1	1.00	0.08	8,8,8,8	0
6	CL	B	1396	1/1	1.00	0.09	14,14,14,14	0
3	MG	A	1396	1/1	1.00	0.09	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.