

Full wwPDB X-ray Structure Validation Report (i)

Sep 25, 2023 - 04:54 PM EDT

ım gut bac-

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	$1284 \ (2.56-2.52)$
Clashscore	141614	$1332 \ (2.56-2.52)$
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	269	^{2%} 92%	5%	·
1	В	269	2% 91%	7%	·
1	С	269	3% 89%	8%	•
1	D	269	% 90%	7%	•
1	Е	269	88%	9%	•



Mol	Chain	Length	Quality of chain	
1	F	269	4% 89%	7% ••
1	G	269	3% 90%	7% •
1	Н	269	<u>6%</u> 90%	5% 5%



$6 \mathrm{BRM}$

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 16403 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace			
1 1	969	Total	С	Ν	0	S	0	0	0				
	A	202	2054	1311	346	387	10	0	0	0			
1	В	263	Total	С	Ν	0	S	0	1	0			
	D	205	2073	1325	348	390	10	0	1	0			
1	С	262	Total	С	Ν	0	S	0	0	0			
	U			U	202	2054	1311	346	387	10	0	0	0
1	Л	261	Total	С	Ν	0	S	0	0	0			
	D	201	2037	1301	342	384	10		0	0			
1	F	F 961	Total	С	Ν	0	S	0	0	0			
	Ľ	201	2049	1308	345	386	10						
1	Б	261	Total	С	Ν	0	S	0	0	0			
	Г	201	2035	1299	340	386	10	0	0	0			
1	1	261	Total	С	Ν	0	S	0	0	0			
	G	201	2041	1305	342	384	10	0	0	0			
1	1 H	256	Total	С	Ν	0	S	0	0	0			
		Н	Н 256	1990	1271	332	377	10	0	0	U		

• Molecule 1 is a protein called Putative metal-dependent isothiocyanate hydrolase SaxA.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	260	SER	-	expression tag	UNP A0A0N7FW12
А	261	ALA	-	expression tag	UNP A0A0N7FW12
А	262	TRP	-	expression tag	UNP A0A0N7FW12
А	263	SER	-	expression tag	UNP A0A0N7FW12
А	264	HIS	-	expression tag	UNP A0A0N7FW12
А	265	PRO	-	expression tag	UNP A0A0N7FW12
А	266	GLN	-	expression tag	UNP A0A0N7FW12
А	267	PHE	-	expression tag	UNP A0A0N7FW12
А	268	GLU	-	expression tag	UNP A0A0N7FW12
А	269	LYS	-	expression tag	UNP A0A0N7FW12
В	260	SER	-	expression tag	UNP A0A0N7FW12
В	261	ALA	-	expression tag	UNP A0A0N7FW12
В	262	TRP	-	expression tag	UNP A0A0N7FW12



6BRM

Continuea from previous page								
Chain	Residue	Modelled	Actual	Comment	Reference			
B	263	SER	-	expression tag	UNP A0A0N7FW12			
B	264	HIS	-	expression tag	UNP A0A0N7FW12			
B	265	PRO	-	expression tag	UNP A0A0N7FW12			
B	266	GLN	-	expression tag	UNP A0A0N7FW12			
B	267	PHE	-	expression tag	UNP A0A0N7FW12			
B	268	GLU	-	expression tag	UNP A0A0N7FW12			
В	269	LYS	-	expression tag	UNP A0A0N7FW12			
C	260	SER	-	expression tag	UNP A0A0N7FW12			
С	261	ALA	-	expression tag	UNP A0A0N7FW12			
С	262	TRP	-	expression tag	UNP A0A0N7FW12			
С	263	SER	-	expression tag	UNP A0A0N7FW12			
С	264	HIS	-	expression tag	UNP A0A0N7FW12			
С	265	PRO	-	expression tag	UNP A0A0N7FW12			
С	266	GLN	-	expression tag	UNP A0A0N7FW12			
С	267	PHE	-	expression tag	UNP A0A0N7FW12			
С	268	GLU	-	expression tag	UNP A0A0N7FW12			
С	269	LYS	-	expression tag	UNP A0A0N7FW12			
D	260	SER	-	expression tag	UNP A0A0N7FW12			
D	261	ALA	-	expression tag	UNP A0A0N7FW12			
D	262	TRP	-	expression tag	UNP A0A0N7FW12			
D	263	SER	-	expression tag	UNP A0A0N7FW12			
D	264	HIS	-	expression tag	UNP A0A0N7FW12			
D	265	PRO	-	expression tag	UNP A0A0N7FW12			
D	266	GLN	-	expression tag	UNP A0A0N7FW12			
D	267	PHE	-	expression tag	UNP A0A0N7FW12			
D	268	GLU	-	expression tag	UNP A0A0N7FW12			
D	269	LYS	-	expression tag	UNP A0A0N7FW12			
Е	260	SER	-	expression tag	UNP A0A0N7FW12			
Е	261	ALA	-	expression tag	UNP A0A0N7FW12			
Е	262	TRP	-	expression tag	UNP A0A0N7FW12			
Е	263	SER	-	expression tag	UNP A0A0N7FW12			
Е	264	HIS	-	expression tag	UNP A0A0N7FW12			
Е	265	PRO	-	expression tag	UNP A0A0N7FW12			
Е	266	GLN	-	expression tag	UNP A0A0N7FW12			
Е	267	PHE	-	expression tag	UNP A0A0N7FW12			
Е	268	GLU	-	expression tag	UNP A0A0N7FW12			
Е	269	LYS	-	expression tag	UNP A0A0N7FW12			
F	260	SER	-	expression tag	UNP A0A0N7FW12			
F	261	ALA	-	expression tag	UNP A0A0N7FW12			
F	262	TRP	-	expression tag	UNP A0A0N7FW12			
F	263	SER	-	expression tag	UNP A0A0N7FW12			
F	264	HIS	-	expression tag	UNP A0A0N7FW12			

Cntia J fa



6BRM

Chain	Residue	Modelled	Actual	Comment	Reference			
F	265	PRO	-	expression tag	UNP A0A0N7FW12			
F	266	GLN	-	expression tag	UNP A0A0N7FW12			
F	267	PHE	-	expression tag	UNP A0A0N7FW12			
F	268	GLU	-	expression tag	UNP A0A0N7FW12			
F	269	LYS	-	expression tag	UNP A0A0N7FW12			
G	260	SER	-	expression tag	UNP A0A0N7FW12			
G	261	ALA	-	expression tag	UNP A0A0N7FW12			
G	262	TRP	-	expression tag	UNP A0A0N7FW12			
G	263	SER	-	expression tag	UNP A0A0N7FW12			
G	264	HIS	-	expression tag	UNP A0A0N7FW12			
G	265	PRO	-	expression tag	UNP A0A0N7FW12			
G	266	GLN	-	expression tag	UNP A0A0N7FW12			
G	267	PHE	-	expression tag	UNP A0A0N7FW12			
G	268	GLU	-	expression tag	UNP A0A0N7FW12			
G	269	LYS	-	expression tag	UNP A0A0N7FW12			
Н	260	SER	-	expression tag	UNP A0A0N7FW12			
Н	261	ALA	-	expression tag	UNP A0A0N7FW12			
Н	262	TRP	-	expression tag	UNP A0A0N7FW12			
Н	263	SER	-	expression tag	UNP A0A0N7FW12			
Н	264	HIS	-	expression tag	UNP A0A0N7FW12			
Н	265	PRO	-	expression tag	UNP A0A0N7FW12			
Н	266	GLN	-	expression tag	UNP A0A0N7FW12			
Н	267	PHE	-	expression tag	UNP A0A0N7FW12			
Н	268	GLU	-	expression tag	UNP A0A0N7FW12			
Н	269	LYS	-	expression tag	UNP A0A0N7FW12			

d fr mtin C

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Zn 2 2	0	0
2	В	2	Total Zn 2 2	0	0
2	С	2	Total Zn 2 2	0	0
2	D	2	Total Zn 2 2	0	0
2	Е	2	Total Zn 2 2	0	0
2	F	2	Total Zn 2 2	0	0
2	G	2	Total Zn 2 2	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	Н	2	Total Zn 2 2	0	0

• Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	Ε	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0
3	G	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 3 1 2 \end{array}$	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	5	Total O 5 5	0	0
4	В	7	Total O 7 7	0	0
4	С	6	Total O 6 6	0	0
4	D	5	$\begin{array}{cc} \text{Total} & \text{O} \\ 5 & 5 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Е	5	Total O 5 5	0	0
4	F	5	Total O 5 5	0	0
4	G	5	Total O 5 5	0	0
4	Н	4	Total O 4 4	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative metal-dependent isothiocyanate hydrolase SaxA









4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	179.16Å 79.53Å 90.41Å	Deperitor	
a, b, c, α , β , γ	90.00° 89.98° 90.00°	Depositor	
$\mathbf{P}_{\text{osolution}}(\hat{\mathbf{A}})$	49.84 - 2.55	Depositor	
Resolution (A)	49.84 - 2.55	EDS	
% Data completeness	97.6 (49.84-2.55)	Depositor	
(in resolution range)	98.5(49.84-2.55)	EDS	
R_{merge}	0.11	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$1.94 (at 2.54 \text{\AA})$	Xtriage	
Refinement program	REFMAC 5.8.0158	Depositor	
D D	0.206 , 0.243	Depositor	
$\mathbf{n}, \mathbf{n}_{free}$	0.225 , 0.264	DCC	
R_{free} test set	4116 reflections (5.00%)	wwPDB-VP	
Wilson B-factor $(Å^2)$	49.9	Xtriage	
Anisotropy	0.226	Xtriage	
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, 23.9	EDS	
L-test for $twinning^2$	$< L >=0.49, < L^2>=0.31$	Xtriage	
Estimated twinning fraction	0.011 for -h,-k,l	Xtriage	
Pepertod twinning fraction	0.657 for H, K, L	Depositor	
Reported twinning fraction	0.343 for -h,-k,l	Depositor	
Outliers	5 of 82280 reflections (0.006%)	Xtriage	
F_o, F_c correlation	0.94	EDS	
Total number of atoms	16403	wwPDB-VP	
Average B, all atoms $(Å^2)$	58.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 73.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7105e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for a centric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.36	0/2098	0.58	0/2850
1	В	0.36	0/2118	0.56	0/2877
1	С	0.36	0/2098	0.57	0/2850
1	D	0.36	0/2081	0.60	0/2828
1	Е	0.35	0/2093	0.57	0/2843
1	F	0.35	0/2077	0.56	0/2821
1	G	0.36	0/2085	0.57	0/2833
1	Н	0.36	0/2029	0.55	0/2755
All	All	0.36	0/16679	0.57	0/22657

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2054	0	2044	3	0
1	В	2073	0	2060	2	0
1	С	2054	0	2044	3	0
1	D	2037	0	2020	6	0
1	Е	2049	0	2042	4	0



6BRM

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2035	0	2013	8	0
1	G	2041	0	2031	7	0
1	Н	1990	0	1975	3	0
2	А	2	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
2	Е	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	Н	2	0	0	0	0
3	А	3	0	1	0	0
3	С	3	0	1	0	0
3	Е	3	0	1	0	0
3	G	3	0	1	0	0
4	А	5	0	0	0	0
4	В	7	0	0	0	0
4	С	6	0	0	0	0
4	D	5	0	0	0	0
4	Е	5	0	0	0	0
4	F	5	0	0	0	0
4	G	5	0	0	0	0
4	Н	4	0	0	0	0
All	All	16403	0	16233	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:244:ILE:HD12	1:H:248:ILE:HD11	1.89	0.54
1:G:6:ILE:HB	1:G:10:THR:HG23	1.91	0.53
1:D:192:ASN:HB2	1:D:198:ILE:HD13	1.91	0.53
1:G:65:LEU:HD12	1:G:98:ILE:HD11	1.90	0.53
1:B:153:HIS:ND1	1:B:154:HIS:O	2.24	0.53
1:F:244:ILE:HD12	1:F:248:ILE:HD11	1.91	0.52
1:B:1:MET:HE1	1:B:159:ILE:HD12	1.94	0.49
1:F:171:TYR:CZ	1:F:175:LEU:HD21	2.48	0.48
1:F:58:LEU:HD22	1:F:81:VAL:HG12	1.96	0.48
1:G:7:ARG:O	1:G:10:THR:HG22	2.13	0.48



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:C:185:LEU:HD13	1:C:200:MET:CE	2.44	0.46
1:F:22:ILE:HD12	1:F:22:ILE:N	2.31	0.46
1:A:240:LEU:HD23	1:A:245:GLU:HG2	1.97	0.46
1:F:105:ASN:HD21	1:G:154:HIS:HB2	1.81	0.46
1:C:150:VAL:HG22	1:C:160:TYR:CD1	2.52	0.45
1:D:127:GLN:HB3	1:D:168:VAL:HG21	1.97	0.45
1:D:7:ARG:O	1:D:10:THR:HG22	2.16	0.45
1:F:205:THR:HG21	1:F:248:ILE:HG12	1.99	0.45
1:H:87:ILE:HB	1:H:106:ILE:HG22	1.98	0.45
1:F:150:VAL:HG22	1:F:160:TYR:CD1	2.52	0.44
1:D:164:ASP:OD1	1:D:186:ASN:ND2	2.44	0.43
1:G:98:ILE:HG22	1:G:103:PHE:HB2	1.99	0.43
1:E:150:VAL:HG22	1:E:160:TYR:CD1	2.53	0.43
1:E:127:GLN:HE22	1:E:131:ASN:ND2	2.16	0.43
1:D:205:THR:CG2	1:D:248:ILE:HD12	2.48	0.43
1:A:116:VAL:HG11	1:D:119:LEU:HD12	2.02	0.42
1:F:105:ASN:HD21	1:G:154:HIS:CB	2.31	0.42
1:A:217:ILE:HD12	1:A:217:ILE:N	2.35	0.42
1:G:58:LEU:HD21	1:G:81:VAL:HG22	2.01	0.42
1:H:244:ILE:HD12	1:H:248:ILE:CD1	2.49	0.42
1:E:65:LEU:HD12	1:E:98:ILE:HD13	2.00	0.42
1:E:88:TYR:HB3	1:E:109:LEU:HD13	2.04	0.40
1:C:235:LEU:HD23	1:C:250:ILE:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	260/269~(97%)	242 (93%)	17 (6%)	1 (0%)	34 46
1	В	262/269~(97%)	244 (93%)	16 (6%)	2 (1%)	19 27



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	С	260/269~(97%)	250~(96%)	9~(4%)	1 (0%)	34 46
1	D	259/269~(96%)	241 (93%)	18 (7%)	0	100 100
1	Ε	259/269~(96%)	246 (95%)	13~(5%)	0	100 100
1	F	257/269~(96%)	239~(93%)	17 (7%)	1 (0%)	34 46
1	G	259/269~(96%)	241 (93%)	17 (7%)	1 (0%)	34 46
1	Н	250/269~(93%)	232 (93%)	18 (7%)	0	100 100
All	All	2066/2152~(96%)	1935 (94%)	125 (6%)	6~(0%)	41 51

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	224	SER
1	А	224	SER
1	С	194	LEU
1	F	224	SER
1	В	73	TRP
1	В	37	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	220/228~(96%)	211~(96%)	9~(4%)	30 41
1	В	222/228~(97%)	209~(94%)	13~(6%)	19 25
1	С	220/228~(96%)	204~(93%)	16 (7%)	14 18
1	D	217/228~(95%)	208~(96%)	9 (4%)	30 41
1	Ε	220/228~(96%)	203~(92%)	17 (8%)	13 16
1	F	217/228~(95%)	206~(95%)	11 (5%)	24 32
1	G	218/228~(96%)	208~(95%)	10 (5%)	27 36
1	Н	212/228~(93%)	202 (95%)	10 (5%)	26 35
All	All	1746/1824~(96%)	1651 (95%)	95(5%)	22 29



All	(95)	residues	with a	non-rotameric	sidechain	are listed	below:
-----	------	----------	--------	---------------	-----------	------------	--------

Mol	Chain	Res	Type
1	А	1	MET
1	А	97	LEU
1	А	109	LEU
1	А	149	LEU
1	А	156	GLU
1	А	177	ARG
1	А	194	LEU
1	А	210	LYS
1	А	227	HIS
1	В	3	LEU
1	В	7	ARG
1	В	24	PRO
1	В	40	ARG
1	В	44	ARG
1	В	65	LEU
1	В	109	LEU
1	В	152	THR
1	В	194	LEU
1	В	211	MET
1	В	230	LEU
1	В	232	ARG
1	В	257	MET
1	С	5	GLN
1	С	21	LEU
1	С	45	ASN
1	С	57	LEU
1	С	64	ILE
1	С	85	MET
1	С	119	LEU
1	С	125	ASP
1	С	136	ASP
1	С	152	THR
1	C	173	LYS
1	C	177	ARG
1	С	194	LEU
1	С	211	MET
1	C	217	ILE
1	С	230	LEU
1	D	7	ARG
1	D	21	LEU
1	D	42	HIS
1	D	44	ARG



Mol	Chain	Res	Type
1	D	57	LEU
1	D	109	LEU
1	D	110	LYS
1	D	149	LEU
1	D	203	GLU
1	Е	1	MET
1	Е	14	GLN
1	Е	22	ILE
1	Е	25	MET
1	Е	85	MET
1	Е	104	PHE
1	Е	109	LEU
1	Е	125	ASP
1	Е	128	HIS
1	Е	149	LEU
1	Е	157	LYS
1	Е	205	THR
1	Е	207	ARG
1	Е	211	MET
1	Е	212	LEU
1	Ε	227	HIS
1	Е	245	GLU
1	F	21	LEU
1	F	79	GLN
1	F	86	LEU
1	F	105	ASN
1	F	109	LEU
1	F	110	LYS
1	F	173	LYS
1	F	175	LEU
1	F	185	LEU
1	F	202	LYS
1	F	211	MET
1	G	50	LEU
1	G	85	MET
1	G	109	LEU
1	G	119	LEU
1	G	125	ASP
1	G	138	GLN
1	G	149	LEU
1	G	212	LEU
1	G	227	HIS



Mol	Chain	Res	Type
1	G	240	LEU
1	Н	11	LEU
1	Н	83	LYS
1	Н	85	MET
1	Н	109	LEU
1	Н	133	LEU
1	Н	159	ILE
1	Н	181	GLU
1	Н	212	LEU
1	Н	217	ILE
1	Н	230	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	138	GLN
1	В	101	GLN
1	С	131	ASN
1	D	192	ASN
1	Е	113	ASN
1	Е	131	ASN
1	Е	176	GLN
1	F	5	GLN
1	G	45	ASN
1	Н	14	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 16 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Trme		Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain Dag	T:1-	Bond lengths		Bond angles		
INIOI	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2									
3	FMT	С	303	-	2,2,2	0.70	0	$1,\!1,\!1$	0.71	0									
3	FMT	G	303	-	2,2,2	0.70	0	$1,\!1,\!1$	0.70	0									
3	FMT	А	303	-	2,2,2	0.73	0	$1,\!1,\!1$	0.70	0									
3	FMT	E	303	-	2,2,2	0.72	0	1,1,1	0.70	0									

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	262/269~(97%)	0.31	6 (2%) 60 67	39, 56, 69, 86	0
1	В	263/269~(97%)	0.30	6 (2%) 60 67	36, 56, 77, 101	0
1	С	262/269~(97%)	0.40	9 (3%) 45 52	36, 55, 85, 114	0
1	D	261/269~(97%)	0.24	4 (1%) 73 79	37, 52, 72, 96	0
1	E	261/269~(97%)	0.36	8 (3%) 49 56	39, 56, 83, 105	0
1	F	261/269~(97%)	0.38	10 (3%) 40 47	37, 58, 83, 111	0
1	G	261/269~(97%)	0.41	8 (3%) 49 56	45, 62, 77, 91	0
1	Н	256/269~(95%)	0.55	17 (6%) 18 21	43, 67, 87, 98	0
All	All	2087/2152~(96%)	0.37	68 (3%) 46 53	36, 58, 82, 114	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	55	GLU	5.8
1	Н	37	GLY	5.5
1	G	6	ILE	5.3
1	G	58	LEU	5.0
1	А	55	GLU	4.0
1	G	54	VAL	3.9
1	F	39	ALA	3.7
1	В	258	ALA	3.7
1	Н	183	VAL	3.5
1	G	52	VAL	3.5
1	F	48	VAL	3.4
1	В	26	LEU	3.3
1	G	57	LEU	3.3
1	А	57	LEU	3.3
1	Е	50	LEU	3.3
1	Н	32	TRP	3.3



Mol	Chain	Res	Type	RSRZ
1	Н	248	ILE	3.2
1	В	27	ALA	3.1
1	В	263	SER	3.0
1	Н	55	GLU	2.9
1	С	49	ALA	2.9
1	С	260	SER	2.8
1	С	258	ALA	2.8
1	А	58	LEU	2.8
1	F	30	GLU	2.8
1	Н	190	ALA	2.8
1	Е	258	ALA	2.7
1	Н	229	LEU	2.7
1	С	42	HIS	2.7
1	F	41	PRO	2.7
1	В	262	TRP	2.7
1	Н	138	GLN	2.6
1	Е	20	PHE	2.6
1	Е	22	ILE	2.5
1	Н	249	LEU	2.5
1	А	6	ILE	2.5
1	С	261	ALA	2.4
1	D	43	LEU	2.4
1	D	42	HIS	2.4
1	Н	218	VAL	2.4
1	D	79	GLN	2.4
1	С	56	ASP	2.4
1	Н	54	VAL	2.4
1	С	51	PRO	2.4
1	Е	35	PHE	2.4
1	F	108	VAL	2.3
1	Е	55	GLU	2.3
1	В	248	ILE	2.3
1	А	215	ALA	2.3
1	F	26	LEU	2.3
1	С	243	GLY	2.3
1	Е	219	ALA	2.3
1	Н	232	ARG	2.3
1	Н	258	ALA	2.2
1	F	98	ILE	2.2
1	D	142	LEU	2.2
1	F	246	ASP	2.1
1	Н	257	MET	2.1



Mol	Chain	Res	Type	RSRZ
1	Н	194	LEU	2.1
1	G	258	ALA	2.1
1	F	102	GLY	2.1
1	F	189	TYR	2.1
1	А	97	LEU	2.1
1	Н	79	GLN	2.1
1	С	219	ALA	2.0
1	Е	260	SER	2.0
1	G	12	VAL	2.0
1	Н	26	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
3	FMT	А	303	3/3	0.84	0.21	$51,\!51,\!54,\!55$	0
3	FMT	G	303	3/3	0.88	0.30	83,83,84,86	0
3	FMT	С	303	3/3	0.90	0.14	$67,\!67,\!68,\!69$	0
3	FMT	Е	303	3/3	0.92	0.12	$64,\!64,\!66,\!67$	0
2	ZN	В	301	1/1	0.96	0.16	48,48,48,48	0
2	ZN	D	302	1/1	0.97	0.14	41,41,41,41	0
2	ZN	G	302	1/1	0.97	0.12	$54,\!54,\!54,\!54$	0
2	ZN	F	301	1/1	0.98	0.16	$52,\!52,\!52,\!52$	0
2	ZN	Н	302	1/1	0.98	0.12	59, 59, 59, 59, 59	0
2	ZN	В	302	1/1	0.99	0.12	$52,\!52,\!52,\!52$	0
2	ZN	F	302	1/1	0.99	0.13	44,44,44,44	0
2	ZN	G	301	1/1	0.99	0.17	52,52,52,52	0
2	ZN	C	301	1/1	0.99	0.13	43,43,43,43	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	ZN	Н	301	1/1	0.99	0.12	$55,\!55,\!55,\!55$	0
2	ZN	С	302	1/1	0.99	0.14	47,47,47,47	0
2	ZN	D	301	1/1	0.99	0.20	$38,\!38,\!38,\!38$	0
2	ZN	А	301	1/1	0.99	0.15	45,45,45,45	0
2	ZN	Е	301	1/1	0.99	0.14	49,49,49,49	0
2	ZN	Е	302	1/1	0.99	0.13	$51,\!51,\!51,\!51$	0
2	ZN	А	302	1/1	1.00	0.15	47,47,47,47	0

Continued from previous page...

6.5 Other polymers (i)

There are no such residues in this entry.

