



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 05:59 pm BST

PDB ID : 2BSH  
Title : Crystal structure of the type III secretion chaperone SycT from *Yersinia enterocolitica* (crystal form 2)  
Authors : Buttner, C.R.; Cornelis, G.R.; Heinz, D.W.; Niemann, H.H.  
Deposited on : 2005-05-21  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	125	 10% 65% 27% • •
1	B	125	 6% 70% 18% 6% 6%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SYCT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	120	993	632	168	190	1	2	0	0	0
1	B	117	948	602	161	181	1	3	0	0	1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	78	Total	O	0	0
			78	78		
2	B	61	Total	O	0	0
			61	61		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SYCT



- Molecule 1: SYCT



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.04Å 92.04Å 55.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	79.81 – 1.90 46.02 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (79.81-1.90) 99.9 (46.02-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.90 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.174 , 0.220 0.174 , 0.219	Depositor DCC
$R_{free}$ test set	1085 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtrriage
Anisotropy	0.338	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2080	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.12	4/1016 (0.4%)	0.95	1/1376 (0.1%)
1	B	1.02	3/967 (0.3%)	0.93	1/1307 (0.1%)
All	All	1.07	7/1983 (0.4%)	0.94	2/2683 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	24	GLU	CD-OE2	10.98	1.37	1.25
1	A	24	GLU	CD-OE1	9.09	1.35	1.25
1	A	57	ASP	CG-OD2	7.57	1.42	1.25
1	B	96	GLU	CG-CD	6.50	1.61	1.51
1	A	84	TRP	CB-CG	6.07	1.61	1.50
1	B	96	GLU	CB-CG	5.64	1.62	1.52
1	B	96	GLU	CD-OE2	5.49	1.31	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	THR	N-CA-C	-6.83	92.56	111.00
1	B	13	PHE	CB-CA-C	5.11	120.62	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	119	PHE	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	993	0	938	32	0
1	B	948	0	898	27	0
2	A	78	0	0	1	0
2	B	61	0	0	2	0
All	All	2080	0	1836	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:PHE:HE2	1:A:20:HIS:HB2	1.11	1.16
1:A:13:PHE:CE2	1:A:20:HIS:HB2	1.99	0.98
1:A:3:THR:O	1:A:3:THR:CG2	2.22	0.88
1:A:3:THR:HG23	1:A:3:THR:O	1.71	0.87
1:B:32:VAL:HG22	1:B:33:ASP:OD1	1.79	0.81
1:B:32:VAL:HG13	1:B:36:ILE:HB	1.62	0.81
1:A:121:PRO:HB2	1:A:122:ASN:OXT	1.86	0.74
1:A:12:LEU:HA	1:A:94:ILE:HD11	1.70	0.74
1:B:24:GLU:CD	1:B:24:GLU:H	1.91	0.73
1:B:30:PHE:HZ	1:B:101:LEU:HD11	1.56	0.70
1:B:33:ASP:OD2	1:B:35:HIS:ND1	2.25	0.70
1:B:23:ASN:HB2	1:B:24:GLU:OE2	1.94	0.68
1:A:30:PHE:CE1	1:A:101:LEU:HD21	2.30	0.67
1:A:6:THR:O	1:A:10:GLN:HG3	1.96	0.65
1:B:15:LYS:HG2	1:B:98:GLN:OE1	1.99	0.62
1:B:113:GLU:OE1	1:B:113:GLU:HA	1.98	0.62
1:A:30:PHE:CZ	1:A:40:ILE:HD11	2.37	0.59
1:A:4:THR:H	1:A:7:GLU:HB2	1.66	0.59
1:A:19:ASN:HB2	1:A:20:HIS:CD2	2.37	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:LYS:HE2	1:B:98:GLN:OE1	2.02	0.58
1:A:30:PHE:HE2	1:A:40:ILE:CD1	2.17	0.58
1:A:109:GLU:O	1:A:113:GLU:HG2	2.04	0.57
1:A:30:PHE:CE2	1:A:40:ILE:CD1	2.88	0.56
1:B:33:ASP:OD2	1:B:35:HIS:CE1	2.58	0.56
1:B:31:GLU:HG2	1:B:37:GLN:HE21	1.70	0.56
1:A:106:ASP:O	1:A:110:GLN:HG3	2.06	0.56
1:A:30:PHE:HE2	1:A:40:ILE:HD12	1.72	0.54
1:A:32:VAL:O	1:A:34:GLY:N	2.41	0.54
1:B:30:PHE:CZ	1:B:101:LEU:HD11	2.39	0.53
1:A:117:THR:H	1:A:121:PRO:HD3	1.73	0.52
1:A:18:LEU:HB3	1:A:20:HIS:ND1	2.24	0.52
1:A:23:ASN:ND2	2:A:2008:HOH:O	2.42	0.52
1:B:10:GLN:O	1:B:14:LEU:HD12	2.11	0.49
1:A:30:PHE:CE2	1:A:40:ILE:HD11	2.48	0.49
1:B:30:PHE:CZ	1:B:101:LEU:HD21	2.47	0.49
1:A:3:THR:N	1:A:4:THR:HA	2.27	0.49
1:A:9:MSE:HA	1:A:9:MSE:HE2	1.96	0.48
1:B:32:VAL:HG13	1:B:36:ILE:CB	2.38	0.48
1:B:15:LYS:CG	1:B:98:GLN:OE1	2.60	0.48
1:A:22:VAL:HA	1:A:27:VAL:O	2.14	0.48
1:B:57:ASP:OD1	1:B:76:ARG:NH1	2.47	0.48
1:A:93:ASP:OD2	1:A:95:ASP:HB2	2.14	0.47
1:A:114:PRO:O	1:A:117:THR:HG22	2.16	0.46
1:A:32:VAL:HB	1:A:36:ILE:HB	1.98	0.45
1:A:30:PHE:CD2	1:A:30:PHE:N	2.85	0.45
1:B:32:VAL:O	1:B:33:ASP:CG	2.55	0.45
1:B:13:PHE:CG	1:B:20:HIS:HB3	2.52	0.45
1:B:2:GLN:HE21	1:B:45:GLN:HG3	1.82	0.45
1:A:56:ALA:O	1:A:57:ASP:HB2	2.16	0.44
1:A:30:PHE:HZ	1:A:40:ILE:HD11	1.83	0.44
1:B:13:PHE:CD2	1:B:20:HIS:HB3	2.52	0.44
1:B:31:GLU:CG	1:B:37:GLN:HE21	2.30	0.43
1:A:60:THR:H	1:A:120:GLN:HE22	1.64	0.43
1:A:4:THR:OG1	1:A:7:GLU:HG3	2.19	0.43
1:B:29:THR:HG22	1:B:30:PHE:N	2.35	0.41
1:B:88:SER:HB2	2:B:2049:HOH:O	2.20	0.41
1:B:92:LEU:CD1	1:B:96:GLU:HG3	2.50	0.41
1:B:96:GLU:HG2	2:B:2046:HOH:O	2.20	0.41
1:B:58:LEU:HB3	1:B:59:PRO:HD2	2.03	0.41

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/125 (94%)	111 (94%)	5 (4%)	2 (2%)	9	2
1	B	115/125 (92%)	110 (96%)	4 (4%)	1 (1%)	17	7
All	All	233/250 (93%)	221 (95%)	9 (4%)	3 (1%)	12	4

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	ASP
1	A	120	GLN
1	B	113	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/108 (101%)	107 (98%)	2 (2%)	59	55
1	B	102/108 (94%)	96 (94%)	6 (6%)	19	10
All	All	211/216 (98%)	203 (96%)	8 (4%)	33	24

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	20	HIS
1	B	14	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	18	LEU
1	B	19	ASN
1	B	24	GLU
1	B	32	VAL
1	B	45	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	46	GLN
1	A	120	GLN
1	B	2	GLN
1	B	11	GLN
1	B	20	HIS
1	B	25	ASN
1	B	37	GLN
1	B	45	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	118/125 (94%)	0.46	12 (10%) 6 8	8, 16, 50, 60	0
1	B	114/125 (91%)	0.25	7 (6%) 21 24	14, 21, 45, 53	0
All	All	232/250 (92%)	0.36	19 (8%) 11 13	8, 19, 47, 60	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	114	PRO	7.1
1	A	18	LEU	6.7
1	A	121	PRO	6.3
1	A	13	PHE	4.8
1	A	19	ASN	4.8
1	A	122	ASN	4.8
1	A	30	PHE	4.4
1	A	17	GLY	4.4
1	A	35	HIS	4.2
1	A	20	HIS	4.1
1	B	30	PHE	3.8
1	B	20	HIS	3.7
1	A	16	LEU	3.4
1	B	18	LEU	3.1
1	A	33	ASP	3.0
1	A	3	THR	2.8
1	B	19	ASN	2.8
1	B	33	ASP	2.8
1	B	113	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.