

wwPDB X-ray Structure Validation Summary Report (i)

Dec 13, 2023 – 03:28 pm GMT

PDB ID : 4BS0

Title: Crystal Structure of Kemp Eliminase HG3.17 E47N,N300D Complexed with

Transition State Analog 6-Nitrobenzotriazole

Authors: Blomberg, R.; Kries, H.; Pinkas, D.M.; Mittl, P.R.E.; Gruetter, M.G.; Privett,

H.K.; Mayo, S.; Hilvert, D.

Deposited on : 2013-06-06

Resolution : 1.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{-}467$

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	318	88%	6%	6%
1	В	318	6% 85%	9%	6%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 15025 atoms, of which 6983 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called KEMP ELIMINASE HG3.17.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace		
1	A	300	Total 5032	C 1600	H 2491	N 443	O 484	S 14	0	33	1
1	В	300	Total 9036	C 2890	H 4480	N 780	O 864	S 22	0	300	1

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P23360
A	0	ALA	-	- expression tag	
A	1	GLU	-	expression tag	UNP P23360
A	304	GLY	-	expression tag	UNP P23360
A	305	SER	-	expression tag	UNP P23360
A	306	ILE	-	expression tag	UNP P23360
A	307	GLU	-	expression tag	UNP P23360
A	308	GLY	-	expression tag	UNP P23360
A	309	ARG	-	expression tag	UNP P23360
A	310	GLY	-	expression tag	UNP P23360
A	311	HIS	-	expression tag	UNP P23360
A	312	HIS	-	expression tag	UNP P23360
A	313	HIS	-	expression tag	UNP P23360
A	314	HIS	-	expression tag	UNP P23360
A	315	HIS	-	expression tag	UNP P23360
A	316	HIS	-	expression tag	UNP P23360
A	6	ILE	VAL	engineered mutation	UNP P23360
A	37	LYS	GLN	engineered mutation	UNP P23360
A	42	MET	GLN	engineered mutation	UNP P23360
A	44	TRP	THR	engineered mutation	UNP P23360
A	50	GLN	LYS	engineered mutation	UNP P23360
A	81	GLY	ARG	engineered mutation	UNP P23360
A	82	ALA	GLY	engineered mutation	UNP P23360
A	83	GLY	HIS	engineered mutation	UNP P23360
A	84	CYS	THR	engineered mutation	UNP P23360

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	89	ASN	SER	engineered mutation	UNP P23360
A	90	PHE	GLN	engineered mutation	UNP P23360
A	105	ILE	THR	engineered mutation	UNP P23360
A	125	THR	ALA	engineered mutation	UNP P23360
A	130	GLY	ASN	engineered mutation	UNP P23360
A	142	ASN	THR	engineered mutation	UNP P23360
A	172	MET	ASN	engineered mutation	UNP P23360
A	208	MET	THR	engineered mutation	UNP P23360
A	234	SER	ALA	engineered mutation	UNP P23360
A	236	LEU	THR	engineered mutation	UNP P23360
A	237	MET	GLU	engineered mutation	UNP P23360
A	267	MET	TRP	engineered mutation	UNP P23360
A	275	ALA	TRP	engineered mutation	UNP P23360
A	276	PHE	ARG	engineered mutation	UNP P23360
A	279	SER	THR	engineered mutation	UNP P23360
В	-1	MET	-	expression tag	UNP P23360
В	0	ALA	-	expression tag	UNP P23360
В	1	GLU	-	expression tag	UNP P23360
В	304	GLY	-	expression tag	UNP P23360
В	305	SER	-	expression tag	UNP P23360
В	306	ILE	-	expression tag	UNP P23360
В	307	GLU	-	expression tag	UNP P23360
В	308	GLY	-	expression tag	UNP P23360
В	309	ARG	-	expression tag	UNP P23360
В	310	GLY	-	expression tag	UNP P23360
В	311	HIS	-	expression tag	UNP P23360
В	312	HIS	_	expression tag	UNP P23360
В	313	HIS	-	expression tag	UNP P23360
В	314	HIS	-	expression tag	UNP P23360
В	315	HIS	-	expression tag	UNP P23360
В	316	HIS	-	expression tag	UNP P23360
В	6	ILE	VAL	engineered mutation	UNP P23360
В	37	LYS	GLN	engineered mutation	UNP P23360
В	42	MET	GLN	engineered mutation	UNP P23360
В	44	TRP	THR	engineered mutation	UNP P23360
В	50	GLN	LYS	engineered mutation	UNP P23360
В	81	GLY	ARG	engineered mutation	UNP P23360
В	82	ALA	GLY	engineered mutation	UNP P23360
В	83	GLY	HIS	engineered mutation	UNP P23360
В	84	CYS	THR	engineered mutation	UNP P23360
В	89	ASN	SER	engineered mutation	UNP P23360
В	90	PHE	GLN	engineered mutation	UNP P23360


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	105	ILE	THR	engineered mutation	UNP P23360
В	125	THR	ALA	engineered mutation	UNP P23360
В	130	GLY	ASN	engineered mutation	UNP P23360
В	142	ASN	THR	engineered mutation	UNP P23360
В	172	MET	ASN	engineered mutation	UNP P23360
В	208	MET	THR	engineered mutation	UNP P23360
В	234	SER	ALA	engineered mutation	UNP P23360
В	236	LEU	THR	engineered mutation	UNP P23360
В	237	MET	GLU	engineered mutation	UNP P23360
В	267	MET	TRP	engineered mutation	UNP P23360
В	275	ALA	TRP	engineered mutation	UNP P23360
В	276	PHE	ARG	engineered mutation	UNP P23360
В	279	SER	THR	engineered mutation	UNP P23360


 \bullet Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0

 \bullet Molecule 3 is 6-NITROBENZOTRIAZOLE (three-letter code: 6NT) (formula: $\mathrm{C_6H_4N_4O_2}).$





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
9	Λ	1	Total	С	Н	N	О	0	0	
3	3 A	1	16	6	4	4	2	0		
2	D	1	Total	С	Н	N	О	0	1	
)	3 B	1	32	12	8	8	4		1	

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	461	Total O 472 472	0	11
4	В	371	Total O 417 417	0	46



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: KEMP ELIMINASE HG3.17





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	76.08Å 77.95Å 98.28Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.34 - 1.09	Depositor
Resolution (A)	28.34 - 1.02	EDS
% Data completeness	93.1 (28.34-1.09)	Depositor
(in resolution range)	81.2 (28.34-1.02)	EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.17 (at 1.02Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
D.D.	0.124 , 0.145	Depositor
R, R_{free}	0.126 , 0.146	DCC
R_{free} test set	12038 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	10.0	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.41, 51.8	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	15025	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 6NT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.49	0/2608	0.67	0/3549	
1	В	0.41	0/4656	0.61	0/6350	
All	All	0.44	0/7264	0.63	0/9899	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2541	2491	2468	20	0
1	В	4556	4480	4478	29	2
2	A	10	0	0	1	0
2	В	10	0	0	0	0
3	A	12	4	4	0	0
3	В	24	8	8	0	0
4	A	472	0	0	14	3
4	В	417	0	0	12	1
All	All	8042	6983	6958	45	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:272[B]:PRO:HG3	4:A:2429:HOH:O	1.50	1.07
1:B:149[A]:GLY:C	4:B:2203[A]:HOH:O	1.99	0.98
1:A:303:GLN:N	4:A:2461:HOH:O	2.00	0.94
1:B:150[A]:GLU:N	4:B:2203[A]:HOH:O	2.01	0.93
1:B:220[B]:GLN:OE1	4:B:2333:HOH:O	1.94	0.84

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:37[B]:LYS:HZ2	4:A:2112:HOH:O[3_654]	1.48	0.12
4:A:2217:HOH:O	4:B:2008:HOH:O[3_644]	2.17	0.03
1:B:110[A]:ASN:OD1	4:A:2358:HOH:O[2_554]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Perce	Percentiles	
1	A	331/318 (104%)	326 (98%)	5 (2%)	0	100	100	
1	В	596/318 (187%)	588 (99%)	8 (1%)	0	100	100	
All	All	927/636 (146%)	914 (99%)	13 (1%)	0	100	100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	275/255 (108%)	271 (98%)	4 (2%)	65 27		
1	В	484/255 (190%)	478 (99%)	6 (1%)	71 36		
All	All	759/510 (149%)	749 (99%)	10 (1%)	71 32		

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	62[B]	ASN
1	В	87[A]	TRP
1	В	87[B]	TRP
1	A	87	TRP
1	В	18[A]	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	58	GLN
1	A	74	GLN
1	A	189	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuna	Chain	Pag	Res Link		Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	SO4	A	1304	-	4,4,4	0.15	0	6,6,6	0.11	0	
3	6NT	В	1305[B]	_	8,13,13	1.24	0	10,18,18	2.19	1 (10%)	
2	SO4	В	1304	-	4,4,4	0.21	0	6,6,6	0.68	0	
2	SO4	A	1303	-	4,4,4	0.15	0	6,6,6	0.42	0	
3	6NT	A	1305	_	8,13,13	1.36	1 (12%)	10,18,18	2.54	5 (50%)	
3	6NT	В	1305[A]	_	8,13,13	1.32	0	10,18,18	2.08	1 (10%)	
2	SO4	В	1303	_	4,4,4	0.14	0	6,6,6	0.09	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	6NT	В	1305[A]	-	-	0/2/4/4	0/2/2/2
3	6NT	В	1305[B]	-	-	0/2/4/4	0/2/2/2
3	6NT	A	1305	-	-	0/2/4/4	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
3	A	1305	6NT	O11-NO1	3.08	1.28	1.22

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	В	1305[B]	6NT	N3-N2-N1	-6.04	103.42	111.25
3	В	1305[A]	6NT	N3-N2-N1	-5.30	104.38	111.25
3	A	1305	6NT	C5-C4-C3A	-4.88	113.80	118.53
3	A	1305	6NT	C6-C5-C4	3.73	126.31	120.61
3	A	1305	6NT	C4-C5-NO1	-3.53	115.63	118.75

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1304	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	300/318 (94%)	0.01	4 (1%) 77 73	6, 14, 22, 28	0
1	В	300/318 (94%)	0.38	19 (6%) 20 20	6, 9, 16, 20	0
All	All	$600/636 \ (94\%)$	0.20	23 (3%) 40 37	6, 11, 21, 28	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	В	181[A]	TYR	5.2	
1	В	44[A]	TRP	4.1	
1	В	259[A]	GLN	3.3	
1	В	141[A]	GLN	3.2	
1	В	102[A]	ASN	3.1	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	SO4	A	1303	5/5	0.86	0.23	35,36,37,38	5
2	SO4	A	1304	5/5	0.87	0.17	30,30,30,31	5
2	SO4	В	1303	5/5	0.93	0.13	40,40,42,42	5
2	SO4	В	1304	5/5	0.97	0.10	23,24,25,25	5
3	6NT	В	1305[A]	12/12	0.99	0.11	7,7,10,10	16
3	6NT	В	1305[B]	12/12	0.99	0.11	7,8,10,10	16
3	6NT	A	1305	12/12	1.00	0.10	7,9,11,11	0

6.5 Other polymers (i)

There are no such residues in this entry.

