



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 10, 2023 – 07:29 pm GMT

PDB ID : 8BSL
Title : Human GLS in complex with compound 12
Authors : Debreczeni, J.E.
Deposited on : 2022-11-25
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.3
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	 6% 78% 7% 15%
1	B	478	 6% 77% 7% 16%
1	C	478	 9% 76% 8% 16%
1	D	478	 5% 80% 5% 15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutaminase kidney isoform, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	408	3167	2017	534	588	28	0	0	0
1	B	403	3139	2005	526	580	28	0	0	0
1	C	403	3123	1995	523	577	28	0	0	0
1	D	405	3154	2009	532	585	28	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	GLY	-	expression tag	UNP O94925
A	122	SER	-	expression tag	UNP O94925
B	121	GLY	-	expression tag	UNP O94925
B	122	SER	-	expression tag	UNP O94925
C	121	GLY	-	expression tag	UNP O94925
C	122	SER	-	expression tag	UNP O94925
D	121	GLY	-	expression tag	UNP O94925
D	122	SER	-	expression tag	UNP O94925

- Molecule 2 is {N}-[5-[[[(3 {R})-1-(5-azanyl-1,3,4-thiadiazol-2-yl)pyrrolidin-3-yl]amino]-1,3,4-thiadiazol-2-yl]-2-phenyl-ethanamide (three-letter code: R90) (formula: C₁₆H₁₈N₈OS₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	16	8	1	2		
2	D	1	Total	C	N	O	S	0	0
			27	16	8	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	176	Total	O	0	0
			176	176		
3	B	216	Total	O	0	0
			216	216		
3	C	156	Total	O	0	0
			156	156		
3	D	223	Total	O	0	0
			223	223		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutaminase kidney isoform, mitochondrial





- Molecule 1: Glutaminase kidney isoform, mitochondrial






- Molecule 1: Glutaminase kidney isoform, mitochondrial





- Molecule 1: Glutaminase kidney isoform, mitochondrial



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.99Å 139.03Å 177.72Å 90.00° 96.01° 90.00°	Depositor
Resolution (Å)	47.64 – 2.38 47.64 – 2.38	Depositor EDS
% Data completeness (in resolution range)	95.5 (47.64-2.38) 95.5 (47.64-2.38)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.37Å)	Xtrriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.209 , 0.246 0.202 , 0.238	Depositor DCC
R_{free} test set	4678 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtrriage
Anisotropy	0.476	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.028 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13408	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R90

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3237	0.56	0/4371
1	B	0.41	0/3208	0.59	0/4328
1	C	0.38	0/3192	0.57	0/4309
1	D	0.39	0/3223	0.58	0/4350
All	All	0.39	0/12860	0.58	0/17358

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3167	0	3130	22	0
1	B	3139	0	3118	21	0
1	C	3123	0	3088	23	0
1	D	3154	0	3127	13	0
2	A	27	0	0	2	0
2	D	27	0	0	0	0
3	A	176	0	0	0	0
3	B	216	0	0	0	0
3	C	156	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	223	0	0	0	0
All	All	13408	0	12463	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:533:LEU:O	1:A:544:ARG:NH2	1.98	0.97
1:B:234:LEU:HD22	1:B:520:ASP:HB3	1.59	0.84
1:A:234:LEU:HD22	1:A:520:ASP:HB3	1.63	0.80
1:D:234:LEU:HD22	1:D:520:ASP:HB3	1.71	0.72
1:A:537:ALA:HB2	1:C:450:PRO:HG2	1.72	0.71
1:B:537:ALA:HB2	1:D:450:PRO:HG2	1.71	0.71
1:B:450:PRO:HG2	1:D:537:ALA:HB2	1.73	0.71
1:C:234:LEU:HD22	1:C:520:ASP:HB3	1.75	0.69
1:A:278:THR:HA	1:A:424:CYS:HB2	1.80	0.62
1:A:450:PRO:HG2	1:C:537:ALA:HB2	1.81	0.62
1:B:340:VAL:HG23	1:B:395:LEU:HD13	1.83	0.60
1:A:193:VAL:HG12	1:A:193:VAL:O	2.02	0.58
1:C:302:THR:OG1	1:C:455:ASN:ND2	2.32	0.58
1:A:156:VAL:HG12	1:A:193:VAL:HG13	1.86	0.57
1:A:477:GLY:O	1:A:529:ASN:HB2	2.05	0.57
1:A:340:VAL:HG23	1:A:395:LEU:HD13	1.87	0.56
1:B:265:VAL:HG22	1:B:498:MET:HE3	1.88	0.56
1:B:479:PRO:HD2	1:B:491:VAL:O	2.08	0.54
1:A:322:PHE:N	2:A:601:R90:N5	2.54	0.53
1:C:379:GLN:HA	1:C:382:ARG:HD3	1.89	0.53
1:D:432:ALA:HB1	1:D:441:PRO:HG2	1.90	0.53
1:B:427:ALA:HB3	1:B:499:MET:HG2	1.91	0.53
1:C:378:PHE:CZ	1:C:382:ARG:HD2	2.45	0.51
1:B:349:VAL:O	1:B:354:LYS:HE3	2.11	0.51
1:A:312:GLU:HG2	1:C:470:GLY:HA3	1.94	0.50
1:C:432:ALA:HB1	1:C:441:PRO:HG3	1.94	0.50
1:B:265:VAL:HG22	1:B:498:MET:CE	2.43	0.49
1:A:148:ALA:HA	1:A:154:ILE:HG12	1.95	0.48
1:B:148:ALA:HA	1:B:154:ILE:HG12	1.95	0.47
1:B:278:THR:HA	1:B:424:CYS:HB2	1.96	0.47
1:C:261:TRP:CE3	1:C:502:SER:HB2	2.49	0.47
1:D:340:VAL:HG23	1:D:395:LEU:HD13	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:340:VAL:HG23	1:C:395:LEU:HD13	1.97	0.47
1:C:382:ARG:NH2	1:C:412:ASP:OD1	2.48	0.47
1:A:321:LEU:HB3	2:A:601:R90:N6	2.31	0.46
1:A:292:LYS:HD3	1:A:417:LEU:HD13	1.98	0.46
1:B:292:LYS:HD3	1:B:417:LEU:HD13	1.98	0.46
1:C:479:PRO:HD2	1:C:491:VAL:O	2.16	0.45
1:B:529:ASN:ND2	1:D:529:ASN:HD21	2.14	0.45
1:D:479:PRO:HD2	1:D:491:VAL:O	2.16	0.45
1:B:332:PRO:CD	1:B:459:LEU:HD13	2.46	0.45
1:D:148:ALA:HA	1:D:154:ILE:HG12	1.99	0.45
1:C:220:VAL:HG21	1:C:495:VAL:HA	1.97	0.45
1:A:313:PRO:HG3	1:A:462:SER:HB2	1.97	0.45
1:D:432:ALA:CB	1:D:441:PRO:HG2	2.47	0.44
1:B:258:PRO:HB3	1:B:504:PRO:HG3	1.99	0.44
1:B:185:THR:O	1:B:188:THR:OG1	2.36	0.44
1:A:349:VAL:O	1:A:354:LYS:HE3	2.17	0.44
1:C:477:GLY:O	1:C:529:ASN:HB2	2.19	0.43
1:D:248:ASP:HA	1:D:254:ALA:HB2	2.00	0.43
1:A:435:ALA:HB2	1:A:491:VAL:HG13	2.01	0.43
1:C:148:ALA:HA	1:C:154:ILE:HG12	2.00	0.43
1:B:462:SER:O	1:B:469:SER:HB3	2.19	0.43
1:C:378:PHE:CE1	1:C:416:GLN:HG3	2.54	0.43
1:A:479:PRO:HG3	1:C:530:TYR:CE1	2.54	0.43
1:D:220:VAL:HG21	1:D:495:VAL:HA	2.01	0.43
1:C:278:THR:HA	1:C:424:CYS:HB2	2.02	0.42
1:C:332:PRO:HD2	1:C:459:LEU:HD13	2.01	0.42
1:A:461:HIS:CE1	1:A:529:ASN:HD22	2.37	0.42
1:C:258:PRO:HB3	1:C:504:PRO:HG3	2.00	0.42
1:C:432:ALA:CB	1:C:441:PRO:HG3	2.50	0.42
1:A:331:ASN:O	1:A:337:GLY:HA3	2.20	0.41
1:B:195:LEU:HD23	1:B:199:LEU:HD23	2.01	0.41
1:D:331:ASN:O	1:D:337:GLY:HA3	2.20	0.41
1:A:515:ILE:HD13	1:A:515:ILE:HA	1.97	0.41
1:B:331:ASN:O	1:B:337:GLY:HA3	2.21	0.41
1:C:332:PRO:CD	1:C:459:LEU:HD13	2.51	0.41
1:A:479:PRO:HD2	1:A:491:VAL:O	2.21	0.41
1:B:498:MET:HE1	1:B:517:PHE:CZ	2.56	0.40
1:B:529:ASN:HD21	1:D:529:ASN:HD21	1.69	0.40
1:C:346:LYS:HG3	1:C:357:TYR:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/478 (84%)	393 (97%)	10 (2%)	1 (0%)	47	61
1	B	397/478 (83%)	391 (98%)	6 (2%)	0	100	100
1	C	397/478 (83%)	388 (98%)	9 (2%)	0	100	100
1	D	401/478 (84%)	391 (98%)	10 (2%)	0	100	100
All	All	1599/1912 (84%)	1563 (98%)	35 (2%)	1 (0%)	51	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	350/415 (84%)	348 (99%)	2 (1%)	86	93
1	B	348/415 (84%)	345 (99%)	3 (1%)	78	89
1	C	345/415 (83%)	340 (99%)	5 (1%)	67	81
1	D	350/415 (84%)	347 (99%)	3 (1%)	78	89
All	All	1393/1660 (84%)	1380 (99%)	13 (1%)	78	89

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	361	PHE
1	A	463	CYS
1	B	311	LYS
1	B	361	PHE
1	B	529	ASN
1	C	152	GLU
1	C	275	THR
1	C	361	PHE
1	C	484	VAL
1	C	508	MET
1	D	275	THR
1	D	361	PHE
1	D	471	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	461	HIS
1	B	529	ASN
1	C	455	ASN
1	D	529	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	R90	A	601	-	21,30,30	0.90	1 (4%)	21,41,41	0.69	1 (4%)
2	R90	D	601	-	21,30,30	0.87	2 (9%)	21,41,41	0.82	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R90	A	601	-	-	0/8/25/25	0/4/4/4
2	R90	D	601	-	-	0/8/25/25	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	R90	C14-N4	2.69	1.36	1.32
2	D	601	R90	C13-C10	2.28	1.57	1.53
2	D	601	R90	C14-N4	2.21	1.35	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	601	R90	C12-C11-C10	-2.74	100.10	104.08
2	A	601	R90	C12-C11-C10	-2.14	100.98	104.08

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	R90	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/478 (85%)	0.37	31 (7%) 13 15	31, 45, 81, 98	0
1	B	403/478 (84%)	0.31	28 (6%) 16 18	26, 37, 72, 88	0
1	C	403/478 (84%)	0.41	43 (10%) 6 6	31, 44, 85, 103	0
1	D	405/478 (84%)	0.37	25 (6%) 20 22	27, 40, 76, 95	0
All	All	1619/1912 (84%)	0.36	127 (7%) 13 14	26, 42, 79, 103	0

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	PRO	7.6
1	C	195	LEU	7.1
1	C	156	VAL	6.2
1	C	186	LEU	6.2
1	B	186	LEU	6.1
1	A	191	ASP	6.1
1	A	190	SER	6.1
1	C	145	TYR	5.8
1	B	154	ILE	5.5
1	B	145	TYR	5.5
1	C	154	ILE	5.5
1	A	136	LEU	5.4
1	C	152	GLU	5.3
1	B	200	PHE	4.8
1	B	195	LEU	4.7
1	D	148	ALA	4.7
1	A	255	LYS	4.6
1	C	144	PHE	4.5
1	C	199	LEU	4.5
1	C	148	ALA	4.5
1	C	147	ILE	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	142	LEU	4.4
1	B	194	MET	4.3
1	C	187	GLN	4.2
1	C	200	PHE	4.2
1	D	193	VAL	4.2
1	B	193	VAL	4.2
1	B	146	THR	4.2
1	D	255	LYS	4.2
1	D	191	ASP	4.1
1	A	150	GLY	4.1
1	A	154	ILE	4.1
1	A	138	SER	4.1
1	A	142	LEU	4.0
1	D	146	THR	4.0
1	D	189	THR	4.0
1	D	142	LEU	4.0
1	D	151	GLN	4.0
1	C	160	ILE	3.9
1	A	193	VAL	3.9
1	C	137	PRO	3.9
1	C	185	THR	3.9
1	C	146	THR	3.9
1	C	149	GLU	3.8
1	B	148	ALA	3.8
1	B	188	THR	3.8
1	A	151	GLN	3.7
1	B	149	GLU	3.7
1	B	142	LEU	3.7
1	D	190	SER	3.6
1	C	546	GLY	3.6
1	B	144	PHE	3.6
1	C	157	HIS	3.5
1	C	215	PHE	3.5
1	D	199	LEU	3.4
1	A	148	ALA	3.4
1	B	156	VAL	3.4
1	D	192	GLY	3.3
1	D	156	VAL	3.3
1	A	186	LEU	3.3
1	C	153	LYS	3.3
1	B	153	LYS	3.2
1	B	152	GLU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	145	TYR	3.1
1	C	251	PRO	3.1
1	A	157	HIS	3.1
1	D	154	ILE	3.1
1	A	152	GLU	3.0
1	C	159	PHE	3.0
1	A	144	PHE	3.0
1	C	158	LYS	3.0
1	A	192	GLY	2.9
1	C	150	GLY	2.9
1	B	151	GLN	2.9
1	A	143	LEU	2.9
1	B	143	LEU	2.9
1	D	200	PHE	2.9
1	C	151	GLN	2.9
1	D	144	PHE	2.9
1	A	546	GLY	2.9
1	C	194	MET	2.9
1	B	196	ASP	2.9
1	D	186	LEU	2.8
1	A	145	TYR	2.8
1	D	157	HIS	2.8
1	C	155	PRO	2.8
1	C	143	LEU	2.8
1	C	161	THR	2.8
1	C	138	SER	2.7
1	C	252	GLN	2.7
1	B	216	ARG	2.7
1	A	200	PHE	2.7
1	B	150	GLY	2.6
1	C	136	LEU	2.6
1	C	141	ASP	2.6
1	B	546	GLY	2.6
1	C	249	TYR	2.5
1	A	189	THR	2.5
1	A	147	ILE	2.5
1	D	149	GLU	2.5
1	B	147	ILE	2.5
1	D	197	LYS	2.5
1	B	155	PRO	2.4
1	C	205	GLN	2.4
1	D	139	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	196	ASP	2.4
1	A	197	LYS	2.4
1	A	216	ARG	2.4
1	A	158	LYS	2.3
1	C	182	LEU	2.3
1	D	138	SER	2.3
1	B	215	PHE	2.3
1	D	544	ARG	2.3
1	C	319	ASN	2.3
1	A	199	LEU	2.3
1	B	158	LYS	2.3
1	B	253	LEU	2.2
1	B	157	HIS	2.2
1	A	156	VAL	2.2
1	A	403	GLU	2.2
1	C	256	PHE	2.1
1	A	205	GLN	2.1
1	D	150	GLY	2.1
1	C	250	ILE	2.1
1	C	403	GLU	2.1
1	D	215	PHE	2.1
1	A	184	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	R90	A	601	27/27	0.89	0.22	58,59,67,67	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	R90	D	601	27/27	0.92	0.20	53,58,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.