



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 7, 2023 – 10:31 pm BST

PDB ID : 8BU6
Title : Structure of DDB1 bound to DS55-engaged CDK12-cyclin K
Authors : Kozicka, Z.; Kempf, G.; Focht, V.; Thoma, N.H.
Deposited on : 2022-11-30
Resolution : 3.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1291 (3.52-3.40)
Clashscore	141614	1372 (3.52-3.40)
Ramachandran outliers	138981	1337 (3.52-3.40)
Sidechain outliers	138945	1338 (3.52-3.40)
RSRZ outliers	127900	1205 (3.52-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	840	
1	D	840	
1	G	840	
2	B	344	
2	E	344	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	344	<p>24% 78% 16% • 6%</p>
3	C	271	<p>12% 87% • 8%</p>
3	F	271	<p>9% 85% 7% 8%</p>
3	I	271	<p>15% 87% • 8%</p>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 67260 atoms, of which 33551 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA damage-binding protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	G	826	12936	4105	6449	1094	1252	36	6449	0	0
1	D	827	12957	4111	6462	1095	1253	36	6462	0	0
1	A	826	12938	4106	6452	1093	1251	36	6452	0	0

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-3	GLY	-	expression tag	UNP Q16531
G	-2	GLY	-	expression tag	UNP Q16531
G	-1	GLY	-	expression tag	UNP Q16531
G	0	ARG	-	expression tag	UNP Q16531
G	700	GLY	-	linker	UNP Q16531
G	701	ASN	-	linker	UNP Q16531
G	702	GLY	-	linker	UNP Q16531
G	703	ASN	-	linker	UNP Q16531
G	704	SER	-	linker	UNP Q16531
G	705	GLY	-	linker	UNP Q16531
G	706	GLU	-	linker	UNP Q16531
G	707	ILE	-	linker	UNP Q16531
G	708	GLN	-	linker	UNP Q16531
D	-3	GLY	-	expression tag	UNP Q16531
D	-2	GLY	-	expression tag	UNP Q16531
D	-1	GLY	-	expression tag	UNP Q16531
D	0	ARG	-	expression tag	UNP Q16531
D	700	GLY	-	linker	UNP Q16531
D	701	ASN	-	linker	UNP Q16531
D	702	GLY	-	linker	UNP Q16531
D	703	ASN	-	linker	UNP Q16531
D	704	SER	-	linker	UNP Q16531
D	705	GLY	-	linker	UNP Q16531

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	706	GLU	-	linker	UNP Q16531
D	707	ILE	-	linker	UNP Q16531
D	708	GLN	-	linker	UNP Q16531
A	-3	GLY	-	expression tag	UNP Q16531
A	-2	GLY	-	expression tag	UNP Q16531
A	-1	GLY	-	expression tag	UNP Q16531
A	0	ARG	-	expression tag	UNP Q16531
A	700	GLY	-	linker	UNP Q16531
A	701	ASN	-	linker	UNP Q16531
A	702	GLY	-	linker	UNP Q16531
A	703	ASN	-	linker	UNP Q16531
A	704	SER	-	linker	UNP Q16531
A	705	GLY	-	linker	UNP Q16531
A	706	GLU	-	linker	UNP Q16531
A	707	ILE	-	linker	UNP Q16531
A	708	GLN	-	linker	UNP Q16531

- Molecule 2 is a protein called Cyclin-dependent kinase 12.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S			
2	H	324	Total	C	H	N	O	P	S	2657	0	0
			5297	1692	2657	446	485	1	16			
2	E	324	Total	C	H	N	O	P	S	2658	0	0
			5298	1692	2658	446	485	1	16			
2	B	325	Total	C	H	N	O	P	S	2663	0	0
			5309	1695	2663	447	486	1	17			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	709	GLY	-	expression tag	UNP Q9NYV4
H	710	GLY	-	expression tag	UNP Q9NYV4
H	711	GLY	-	expression tag	UNP Q9NYV4
H	965	ARG	LYS	engineered mutation	UNP Q9NYV4
H	1052	GLN	-	expression tag	UNP Q9NYV4
E	709	GLY	-	expression tag	UNP Q9NYV4
E	710	GLY	-	expression tag	UNP Q9NYV4
E	711	GLY	-	expression tag	UNP Q9NYV4
E	965	ARG	LYS	engineered mutation	UNP Q9NYV4
E	1052	GLN	-	expression tag	UNP Q9NYV4
B	709	GLY	-	expression tag	UNP Q9NYV4
B	710	GLY	-	expression tag	UNP Q9NYV4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	711	GLY	-	expression tag	UNP Q9NYV4
B	965	ARG	LYS	engineered mutation	UNP Q9NYV4
B	1052	GLN	-	expression tag	UNP Q9NYV4

- Molecule 3 is a protein called Cyclin-K.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	I	248	Total	C	H	N	O	S	2048	0	0
			4111	1341	2048	346	363	13			
3	F	248	Total	C	H	N	O	S	2048	0	0
			4111	1341	2048	346	363	13			
3	C	248	Total	C	H	N	O	S	2048	0	0
			4111	1341	2048	346	363	13			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
I	-3	GLY	-	expression tag	UNP O75909
I	-2	GLY	-	expression tag	UNP O75909
I	-1	GLY	-	expression tag	UNP O75909
I	0	ARG	-	expression tag	UNP O75909
F	-3	GLY	-	expression tag	UNP O75909
F	-2	GLY	-	expression tag	UNP O75909
F	-1	GLY	-	expression tag	UNP O75909
F	0	ARG	-	expression tag	UNP O75909
C	-3	GLY	-	expression tag	UNP O75909
C	-2	GLY	-	expression tag	UNP O75909
C	-1	GLY	-	expression tag	UNP O75909
C	0	ARG	-	expression tag	UNP O75909

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	G	1	Total	O	S	0	0
			5	4	1		
4	I	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is {N}-(1 {H}-benzimidazol-2-ylmethyl)-9-(1-methylpyrazol-4-yl)-2-morpholin-4-yl-purin-6-amine (three-letter code: RQE) (formula: C₂₁H₂₂N₁₀O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
			Total	C	H	N			O	
5	H	1	Total	54	21	22	10	1	22	0
5	E	1	Total	54	21	22	10	1	22	0
5	B	1	Total	54	21	22	10	1	22	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA damage-binding protein 1





- Molecule 1: DNA damage-binding protein 1







• Molecule 1: DNA damage-binding protein 1




• Molecule 2: Cyclin-dependent kinase 12







- Molecule 2: Cyclin-dependent kinase 12




- Molecule 2: Cyclin-dependent kinase 12




- Molecule 3: Cyclin-K



• Molecule 3: Cyclin-K



• Molecule 3: Cyclin-K



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	249.27Å 249.27Å 220.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	60.68 – 3.45 215.88 – 3.45	Depositor EDS
% Data completeness (in resolution range)	73.8 (60.68-3.45) 73.9 (215.88-3.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 3.41Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.218 , 0.249 0.233 , 0.257	Depositor DCC
R_{free} test set	3856 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	139.6	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 101.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	0.067 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	67260	wwPDB-VP
Average B, all atoms (Å ²)	165.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, TPO, RQE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.33	0/6603	0.60	0/8930
1	D	0.33	0/6612	0.60	0/8942
1	G	0.33	0/6604	0.61	0/8931
2	B	0.39	1/2693 (0.0%)	0.62	1/3630 (0.0%)
2	E	0.34	0/2687	0.61	0/3622
2	H	0.34	0/2687	0.60	0/3622
3	C	0.33	0/2120	0.56	0/2868
3	F	0.34	0/2120	0.56	0/2868
3	I	0.33	0/2120	0.56	0/2868
All	All	0.34	1/34246 (0.0%)	0.60	1/46281 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	888	GLU	CG-CD	-6.75	1.41	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	888	GLU	CG-CD-OE2	-5.71	106.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6486	6452	6454	67	0
1	D	6495	6462	6464	48	1
1	G	6487	6449	6451	48	0
2	B	2646	2663	2663	33	1
2	E	2640	2658	2658	30	0
2	H	2640	2657	2657	34	0
3	C	2063	2048	2048	7	0
3	F	2063	2048	2048	11	0
3	I	2063	2048	2048	5	0
4	A	5	0	0	0	0
4	C	5	0	0	0	0
4	F	10	0	0	0	0
4	G	5	0	0	0	0
4	I	5	0	0	0	0
5	B	32	22	0	0	0
5	E	32	22	0	0	0
5	H	32	22	0	0	0
All	All	33709	33551	33491	274	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 274 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:858:ARG:NH2	2:B:893:TPO:O1P	2.04	0.89
2:E:829:VAL:HG12	2:E:1033:LEU:HD12	1.55	0.88
2:B:828:LEU:HB3	2:B:1034:PRO:HG3	1.68	0.76
2:H:835:HIS:HD2	2:H:1029:ALA:HB1	1.49	0.75
1:A:7:VAL:HG13	1:A:1091:GLY:HA3	1.71	0.73

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:368:GLU:O	2:B:888:GLU:OE2[4_565]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	822/840 (98%)	800 (97%)	21 (3%)	1 (0%)	51	84
1	D	823/840 (98%)	805 (98%)	17 (2%)	1 (0%)	51	84
1	G	822/840 (98%)	801 (97%)	21 (3%)	0	100	100
2	B	322/344 (94%)	314 (98%)	8 (2%)	0	100	100
2	E	321/344 (93%)	309 (96%)	11 (3%)	1 (0%)	41	75
2	H	321/344 (93%)	311 (97%)	9 (3%)	1 (0%)	41	75
3	C	246/271 (91%)	242 (98%)	4 (2%)	0	100	100
3	F	246/271 (91%)	242 (98%)	4 (2%)	0	100	100
3	I	246/271 (91%)	242 (98%)	4 (2%)	0	100	100
All	All	4169/4365 (96%)	4066 (98%)	99 (2%)	4 (0%)	51	84

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	895	LYS
2	H	911	GLU
1	D	772	SER
1	A	980	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	721/728 (99%)	716 (99%)	5 (1%)	84	93
1	D	722/728 (99%)	719 (100%)	3 (0%)	91	97
1	G	721/728 (99%)	712 (99%)	9 (1%)	71	87
2	B	292/308 (95%)	287 (98%)	5 (2%)	60	82
2	E	291/308 (94%)	284 (98%)	7 (2%)	49	76
2	H	291/308 (94%)	285 (98%)	6 (2%)	53	78
3	C	223/242 (92%)	222 (100%)	1 (0%)	91	97
3	F	223/242 (92%)	221 (99%)	2 (1%)	78	91
3	I	223/242 (92%)	221 (99%)	2 (1%)	78	91
All	All	3707/3834 (97%)	3667 (99%)	40 (1%)	73	88

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	231	TYR
2	B	896	VAL
1	A	7	VAL
1	A	930	VAL
2	B	1025	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	835	HIS
2	E	864	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TPO	E	893	2	8,10,11	1.89	2 (25%)	10,14,16	1.07	0
2	TPO	H	893	2	8,10,11	1.04	0	10,14,16	0.94	0
2	TPO	B	893	2	8,10,11	1.67	1 (12%)	10,14,16	1.23	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	E	893	2	-	5/9/11/13	-
2	TPO	H	893	2	-	0/9/11/13	-
2	TPO	B	893	2	-	1/9/11/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	893	TPO	P-O1P	3.50	1.61	1.50
2	B	893	TPO	P-O1P	3.36	1.61	1.50
2	E	893	TPO	CB-CA	2.34	1.59	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	893	TPO	CG2-CB-CA	-2.67	107.90	113.16

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	893	TPO	N-CA-CB-CG2
2	E	893	TPO	N-CA-CB-OG1
2	E	893	TPO	C-CA-CB-CG2
2	E	893	TPO	O-C-CA-CB
2	E	893	TPO	CG2-CB-OG1-P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	893	TPO	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	C	301	-	4,4,4	0.14	0	6,6,6	0.11	0
4	SO4	A	1201	-	4,4,4	0.14	0	6,6,6	0.26	0
5	RQE	H	1101	-	30,37,37	1.04	2 (6%)	27,53,53	1.00	2 (7%)
4	SO4	I	301	-	4,4,4	0.16	0	6,6,6	0.23	0
5	RQE	E	1101	-	30,37,37	0.97	2 (6%)	27,53,53	1.00	2 (7%)
4	SO4	F	302	-	4,4,4	0.13	0	6,6,6	0.08	0
5	RQE	B	1101	-	30,37,37	0.94	1 (3%)	27,53,53	1.10	3 (11%)
4	SO4	G	1201	-	4,4,4	0.17	0	6,6,6	0.16	0
4	SO4	F	301	-	4,4,4	0.13	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	RQE	E	1101	-	-	7/7/21/21	0/6/6/6
5	RQE	H	1101	-	-	5/7/21/21	0/6/6/6
5	RQE	B	1101	-	-	7/7/21/21	0/6/6/6

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	1101	RQE	C3-N3	-3.21	1.40	1.44
5	B	1101	RQE	C6-N8	2.31	1.36	1.32
5	E	1101	RQE	C3-N3	-2.16	1.41	1.44
5	H	1101	RQE	C4-N4	-2.09	1.31	1.34
5	E	1101	RQE	C6-N8	2.07	1.35	1.32

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	1101	RQE	C5-C6-N8	-2.98	118.33	120.81
5	H	1101	RQE	C5-C6-N8	-2.80	118.48	120.81
5	B	1101	RQE	N8-C15-N9	2.39	120.05	117.11
5	B	1101	RQE	C12-C11-C10	-2.31	116.76	120.08
5	H	1101	RQE	C12-C11-C10	-2.21	116.91	120.08

There are no chirality outliers.



5 of 19 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	H	1101	RQE	C8-C7-N5-C6
5	H	1101	RQE	N10-C15-N9-C16
5	H	1101	RQE	N10-C15-N9-C19
5	H	1101	RQE	N8-C15-N9-C16
5	H	1101	RQE	N8-C15-N9-C19

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9	
1	A	826/840 (98%)	0.87	147 (17%)	1 2	112, 150, 228, 315	0
1	D	827/840 (98%)	0.86	133 (16%)	1 2	103, 142, 218, 327	0
1	G	826/840 (98%)	0.90	152 (18%)	1 2	103, 147, 223, 337	0
2	B	324/344 (94%)	1.11	60 (18%)	1 2	95, 131, 210, 287	0
2	E	323/344 (93%)	1.08	64 (19%)	1 1	109, 143, 221, 270	0
2	H	323/344 (93%)	1.17	83 (25%)	0 0	113, 153, 218, 284	0
3	C	248/271 (91%)	0.94	32 (12%)	3 5	102, 129, 176, 242	0
3	F	248/271 (91%)	0.91	25 (10%)	7 9	96, 120, 170, 227	0
3	I	248/271 (91%)	0.89	41 (16%)	1 2	111, 143, 192, 247	0
All	All	4193/4365 (96%)	0.94	737 (17%)	1 2	95, 143, 220, 337	0

The worst 5 of 737 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	1034	PRO	13.2
1	G	1016	ASN	9.3
2	H	720	GLY	9.3
2	E	1035	HIS	9.0
2	B	1039	CYS	7.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TPO	E	893	11/12	0.61	0.18	185,192,232,235	6
2	TPO	H	893	11/12	0.74	0.19	190,198,235,241	5
2	TPO	B	893	11/12	0.77	0.20	145,162,191,196	6

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	F	301	5/5	0.79	0.17	152,159,165,209	0
4	SO4	A	1201	5/5	0.81	0.19	164,165,175,222	0
4	SO4	G	1201	5/5	0.82	0.23	140,146,151,191	0
5	RQE	E	1101	32/32	0.90	0.41	91,128,155,157	22
5	RQE	H	1101	32/32	0.91	0.41	116,129,158,162	22
4	SO4	I	301	5/5	0.91	0.20	142,143,146,175	0
4	SO4	C	301	5/5	0.92	0.14	154,154,163,195	0
5	RQE	B	1101	32/32	0.92	0.38	104,116,141,145	22
4	SO4	F	302	5/5	0.97	0.19	133,135,138,156	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.