



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 30, 2022 – 12:29 am GMT

PDB ID : 2BX3  
Title : Crystal Structure of SARS Coronavirus Main Proteinase (P43212)  
Authors : Verschueren, K.H.G.; Mesters, J.R.; Hilgenfeld, R.  
Deposited on : 2005-07-22  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.31.3  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	306	75% 21% . .

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	300	2413	1529	405	455	24	0	12	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	259	Total	O	0	0
			259	259		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase nsp5

Chain A:  75% 21%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.67Å 69.67Å 100.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.80 – 2.00 40.68 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.9 (56.80-2.00) 97.5 (40.68-1.74)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.48 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.175 , 0.257 0.188 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.7	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2672	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.89	2/2486 (0.1%)	0.84	0/3375

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	145	CYS	CB-SG	-6.84	1.70	1.82
1	A	22	CYS	CB-SG	-6.24	1.71	1.82

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	LYS	Peptide
1	A	138	GLY	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2413	0	2358	59	0
2	A	259	0	0	18	2
All	All	2672	0	2358	59	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:CYS:SG	1:A:61:LYS:NZ	2.09	1.24
1:A:19:GLN:HE21	1:A:26:THR:HG21	1.08	1.15
1:A:6[B]:MET:SD	2:A:2255:HOH:O	2.05	1.13
1:A:45[B]:THR:OG1	2:A:2056:HOH:O	1.83	0.94
1:A:298[B]:ARG:NH1	2:A:2255:HOH:O	2.17	0.77
1:A:19:GLN:NE2	1:A:26:THR:HG21	1.94	0.76
1:A:186:VAL:H	1:A:192:GLN:HE22	1.33	0.74
1:A:137:LYS:O	2:A:2139:HOH:O	2.09	0.70
1:A:3:PHE:CZ	1:A:300:CYS:SG	2.85	0.69
1:A:19:GLN:HE21	1:A:26:THR:CG2	1.98	0.68
1:A:163:HIS:CE1	1:A:172:HIS:HB3	2.32	0.65
1:A:47[A]:GLU:OE2	2:A:2059:HOH:O	2.14	0.64
1:A:240:GLU:OE2	2:A:2208:HOH:O	2.14	0.64
1:A:298[B]:ARG:HH21	1:A:298[B]:ARG:HG2	1.63	0.63
1:A:123:SER:HB2	2:A:2130:HOH:O	1.98	0.62
1:A:118[B]:TYR:HE2	2:A:2130:HOH:O	1.83	0.62
1:A:224:THR:O	2:A:2191:HOH:O	2.16	0.59
1:A:298[B]:ARG:HH21	1:A:298[B]:ARG:CG	2.15	0.59
1:A:138:GLY:O	1:A:140:PHE:N	2.36	0.55
1:A:118[A]:TYR:CD1	1:A:118[A]:TYR:O	2.60	0.55
1:A:298[B]:ARG:HG2	1:A:298[B]:ARG:NH2	2.21	0.55
1:A:40:ARG:HA	1:A:87:LEU:HG	1.88	0.54
1:A:61:LYS:HE2	1:A:65:SER:OG	2.08	0.54
1:A:117:CYS:O	1:A:144:SER:HB3	2.08	0.54
1:A:288[B]:GLU:HG2	1:A:291:PHE:CE2	2.43	0.53
1:A:117:CYS:O	1:A:144:SER:CB	2.57	0.53
1:A:45[B]:THR:HG22	1:A:48:ASP:OD2	2.09	0.53
1:A:227:LEU:HD11	1:A:242:LEU:HD23	1.92	0.51
1:A:144:SER:HA	2:A:2141:HOH:O	2.09	0.51
1:A:19:GLN:NE2	1:A:119:ASN:HB3	2.26	0.50
1:A:165[A]:MET:SD	1:A:186:VAL:O	2.70	0.49
1:A:277:ASN:CG	1:A:277:ASN:O	2.51	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:ASN:ND2	2:A:2232:HOH:O	2.45	0.49
1:A:139:SER:HA	1:A:172:HIS:NE2	2.27	0.48
1:A:256:GLN:NE2	2:A:2221:HOH:O	2.46	0.48
1:A:245:ASP:HB3	2:A:2214:HOH:O	2.13	0.48
1:A:60:ARG:HH11	1:A:60:ARG:HG3	1.79	0.47
1:A:118[A]:TYR:O	1:A:119:ASN:C	2.52	0.47
1:A:113:SER:O	1:A:149:GLY:HA2	2.15	0.46
1:A:123:SER:CB	2:A:2130:HOH:O	2.59	0.46
1:A:108:PRO:HD3	2:A:2123:HOH:O	2.16	0.45
1:A:106:ILE:HD11	1:A:112:PHE:CE1	2.52	0.45
1:A:106:ILE:CD1	1:A:112:PHE:CD1	3.00	0.45
1:A:56:ASP:OD1	1:A:60:ARG:NH1	2.50	0.45
1:A:96:PRO:HA	2:A:2109:HOH:O	2.17	0.45
1:A:220:LEU:HD21	1:A:259:ILE:HD13	1.99	0.44
1:A:298[A]:ARG:NH2	2:A:2257:HOH:O	2.50	0.44
1:A:17:MET:HG3	1:A:117:CYS:SG	2.57	0.44
1:A:102:LYS:HZ3	1:A:104:VAL:CG1	2.31	0.44
1:A:256:GLN:HG3	2:A:2220:HOH:O	2.18	0.43
1:A:86:LEU:HG	1:A:179:GLY:HA2	2.00	0.43
1:A:222:ARG:HA	1:A:222:ARG:HD2	1.83	0.42
1:A:233:VAL:O	1:A:234:ALA:C	2.58	0.42
1:A:167:LEU:HB3	1:A:168:PRO:CD	2.50	0.41
1:A:244:GLN:O	1:A:247:VAL:HB	2.21	0.41
1:A:164:HIS:CD2	1:A:175:THR:HG23	2.56	0.41
1:A:220:LEU:HD23	1:A:220:LEU:HA	1.95	0.40
1:A:31:TRP:CD2	1:A:95:ASN:HB2	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2109:HOH:O	2:A:2221:HOH:O[6_565]	1.90	0.30
2:A:2084:HOH:O	2:A:2228:HOH:O[4_455]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	310/306 (101%)	299 (96%)	7 (2%)	4 (1%)	12 6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	SER
1	A	142	ASN
1	A	154	TYR
1	A	251	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/263 (103%)	260 (96%)	10 (4%)	34 32

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6[A]	MET
1	A	6[B]	MET
1	A	27	LEU
1	A	118[A]	TYR
1	A	118[B]	TYR
1	A	123	SER
1	A	222	ARG
1	A	284	SER
1	A	286	ILE
1	A	289	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	41	HIS
1	A	127	GLN
1	A	142	ASN
1	A	192	GLN
1	A	214	ASN
1	A	246	HIS
1	A	256	GLN
1	A	274	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.